I am grateful to Dr. Wilfrid G. Harding, medical officer of health, London Borough of Camden, for his interest and encouragement.—I am, etc.,

R. ROUTLEDGE

Shenfield, Essex

- 1 Axelsson, B., *Odontologisk Revy*, 195, 6, No. 1.
- 2 Murdock, H. G., *Journal of the American Medical Association*, 1946, 131, 1103.
- 3 Harris, D. K., *British Journal of Industrial Medicine*, 1953, 10, 25.
- 4 Ehrlicher, H., in *Kongress für Arbeitsschutz 1961*. Darmstadt, Steinkopf, 1962.
- 5 Karpov, B. D., *Trudy Leningradskogo Sanitarnogigienicheskogo Meditsinskogo Instituta*, 1953, 14, 43.
- 6 Karpov, B. D., *Gigiena i Sanitarya*, 1954, No. 10, 25.
- 7 Raines, L. A., *Gigiena Truda i Professional'nye Zabolevaniya*, 1957, 1, 56.
- 8 Harris, D. K., *British Journal of Industrial Medicine*, 1959, 16, 221.

Making Hospital Geriatrics Work

SIR,—It is interesting to read that one of the consequences of Drs. H. M. Hodkinson and P. M. Jeffery's approach to "Making Hospital Geriatrics Work" (2 December, p. 536) is the maintenance of the morale and professional satisfaction of "ancillary staff." May I, as a physiotherapist working in this particular arena, put a different point of view?

Paramedical staff involved in the geriatric service frequently experience feelings of impotence and frustration. This is one of the major causes for poor recruitment in this field of medicine and I do not believe that the situation can be remedied by increasing patient turnover.

Rehabilitation of the elderly patient is a lengthy and detailed procedure. Generally it is not so much the acute medical problems that concern the therapist but, to quote Drs. Hodkinson and Jefferys, "the unavoidable [case load] of patients with greater disability." If such patients are to return once again as viable members of their community and maintain optimum independence there, then much time needs to be given by all those involved in their reablement. In these situations, the therapist will need medical support over a long period rather than medical "pressure" to carry out such a programme effectively. Drs. Hodkinson and Jefferys may be able to concede "less than complete therapeutic results," but in rehabilitation of older patients there should be no such compromise. The process cannot be hurried. The rate of turnover may be slowed down in consequence, but the alternative for the patient is breakdown at home and eventual hospital readmission, and for the therapist the knowledge that one has been prevented from giving one's professional best.

After three months in hospital, in Drs. Hodkinson and Jefferys's view, discharge becomes very much harder because of "institutionalization" and "the withering of the patient's ties with the community." One might venture to suggest that perhaps it is those with the greatest disability who tend to have the longest stay in hospital. If a patient has adequate incentive and his reablement programme is both realistic and appropriate, it has not been my experience that "institutionalization" occurs, however prolonged his hospital stay may be. Moreover—and this is essential—if relatives, friends, and supporting social service personnel are involved from the beginning,

need there be this inevitable "withering" of community ties? Rehabilitation after all, is an exercise in vivo and not in vitro.—I am, etc.,

MARGARET HAWKER
Superintendent Physiotherapist

Geriatric Unit,
Edgware General Hospital,
Edgware, Middlesex

Sickle-cell Anaemia

SIR,—Sickle-cell anaemia was unknown to medicine until 1905, when James B. Herrick of Chicago, an experienced haematologist in a country with a large negro population, seems to have got the shock of his life on seeing the sickling phenomenon in the blood film of a negro with severe anaemia.¹ I have been informed by a haematologist that of course sickle-cell anaemia was well known before 1905. However, at the beginning of the century there were numerous books on haematology with coloured illustrations of the bloods of all known anaemias, and at least in British, American, French, and German books there is no mention of sickle-cell anaemia or thalassaemia, with its almost equally arresting blood picture. These two words seem to have been unknown to the official nomenclatures of those countries. Can anyone refer me to any evidence in black and white in any language that these two diseases were known to exist before 1905?—I am, etc.,

H. GALL

Richmond upon Thames,
Surrey

¹ Herrick, J. B., *Archives of Internal Medicine*, 1910, 6, 517.

Nitrazepam: Another Interesting Syndrome

SIR,—A 55-year-old woman for many years accustomed to taking five 100 mg tablets of butobarbitone during the night was admitted to hospital for treatment for her alcoholism. She was immediately changed to nitrazepam 10 mg at bedtime, something I had been unable to achieve in her home environment. Six weeks later for the first time, and after her discharge home, she complained of pins and needles in both hands and that the fingers were clumsy and losing their power. Her symptoms were worse in the morning and gradually improved as the day progressed. On examination there was no evidence of peripheral neuritis, and no other obvious cause was found. The severity of her symptoms increased, and when she was woken during the night by the tingling in her hands she resorted to taking another nitrazepam tablet to obtain sleep.

Her hands became useless in the mornings; she had to stop knitting and could not hold a needle to sew on a button. By late evening her hands were almost back to normal. One night she omitted her nitrazepam because she had to see her son off at 5 a.m. She noticed for the first time in weeks that her hands felt normal. She took her nitrazepam at 5 a.m., and within two hours she woke with her hands tingling and feeling numb once more. Nitrazepam was immediately stopped and she has had no recurrence of the symptoms since.

I am drawing attention to this case because when my patient was wakened by her symptoms she took another dose of nitrazepam,

thereby inadvertently aggravating them. I did not elicit the history of improvement as the day wore on until afterwards. It would appear that when a patient complains of acroparaesthesiae for which no other cause can be found an inquiry into the taking of nitrazepam should be made.—I am, etc.,

HAMISH MACLEAN

Gairloch, Rossshire

Internal Fixation of Secondary Deposits in Long Bones

SIR,—We would like to question the finding of Campbell,¹ quoted by Mr. Malcolm Fidler (10 February, p. 341), that malignant tissue is not disseminated along the shaft of a long bone by the insertion of a Küntscher nail.

It is our clinical experience to have discovered, in some cases, further neoplastic deposits at the opposite end of the bone to the point of entry of the internal splint, some months after the operation and local radiotherapy. Because of this it has been our policy for some years now to irradiate (preferably with megavoltage) the whole length of the shaft postoperatively.—We are, etc.,

W. H. BOND

A. J. BANKS

WILLIAM G. JONES

Queen Elizabeth Hospital,
Birmingham

¹ Campbell, C. J., in *Palliative Care of the Cancer Patient*, ed. R. C. Hickey, p. 313. Boston, Little, Brown and Co., 1967.

Anthrax

SIR,—Dr. Robert Lamb's article on anthrax (20 January, p. 157) was most valuable for doctors working in the U.K. but did not cover the full spectrum of the disease as seen elsewhere.

During my service a good many years ago as a pathologist in the Gold Coast (now Ghana) it was not very uncommon to encounter anthrax in the necropsy room, although a malignant pustule was never seen. The story was usually that the body had been found under a tree, supplemented occasionally by evidence that the deceased was one of many people who had eaten the meat of a cow which had been killed when moribund. The ratio of reported fatalities to incidents was usually unity, and it seemed likely that the common mode of infection was by ingestion and that the infectivity of the organism by this route was very low.

Soon after my arrival at the Accra laboratory I was taught, by Dr. George Robinson, that it was a good plan to remove the skull-cap before opening the trunk when doing a necropsy. A bulging purple dura with haemorrhagic meningitis was suggestive of anthrax, which could be confirmed by puncturing the membranes and withdrawing a drop of cerebrospinal fluid for examination; in this way further opening of an infected body could be avoided. My curiosity once led me to do a full necropsy on one of these patients, in whom large masses of mesenteric lymph nodes, which looked like ripe Victoria plums, convinced me that the gut had indeed been the portal of entry of the infection.

On one occasion only was I able to diag-

nose anthrax during life by finding the organism in the cerebrospinal fluid, but the patient died despite treatment with penicillin.—I am, etc.,

M. H. HUGHES

Public Health Laboratory,
Winchester

Glucagon and Growth Hormone

SIR,—I read with interest your leading article on the effect of glucagon on growth hormone release (27 January, p. 188).

In a study on 28 children who were investigated for possible growth hormone deficiency¹ we found that twenty-three showed a marked rise in serum growth hormone level during a glucagon test. In 20 of these patients a significant rise took place within two minutes of injecting 15 µg of glucagon intravenously. The mean level at two minutes was double the level before glucagon administration. In stress tests, such as administration of insulin, there is usually a latent period of 15-30 minutes before growth hormone levels rise. We therefore consider that glucagon has a pharmacological effect either directly on the adenohypophysis or secondary to a rise in the level of cyclic adenosine monophosphate.

Glucagon given intravenously appears to be without effect on adult males, whereas intravenous insulin and heavy exercise produce growth hormone release in these subjects. It is possible that more growth hormone is readily available in growing children and hence is more easily released by lesser stimuli than in the adult whose growth has ceased.—I am, etc.,

L. STIMMLER

Guy's Hospital Medical School,
London S.E.1

¹ Snodgrass, G., and Stimmler, L., *Journal of Clinical Endocrinology and Metabolism*, 1972, 34, 410.

Ethics in Epidemiological Research

SIR,—Your leading article (27 January, p. 187) on the Medical Research Council's statement on "Responsibility in the Use of Medical Information for Research" (p. 213) states, quite rightly, that "the need to communicate confidential information is particularly evident in epidemiological research" (though "need to obtain" would have been a happier phrase, and "desire to obtain" even happier). However, in its statement the M.R.C. acknowledges "the help and advice received from its Committee on General Epidemiology, which was responsible for drafting this statement" (my italics). The impression is certainly given that what is printed is a statement written by the committee and then formally approved by the M.R.C.; and if this is not the case, perhaps we may be told, at least in general terms, who it was who prepared a modified version of the committee's draft, and with whose help. There is no mention of lay persons with experience in the handling of confidential information—say, lawyers or journalists—let alone members of the general public being asked to help in any way with the drafting of the statement, though the M.R.C. itself received assistance from "the chief medical officers of the Health Departments, the British Medical Association, and the directors of certain of its own establish-

ments." Were there no priests available, no patients, no common or garden jurymen?

Like the other persons involved, the members of the committee, I have no doubt, are all most able and conscientious men (and I am not here concerned with the intrinsic merits of their statement). But it does seem curious, as it is mainly epidemiologists who want this confidential information, that the rules that shall govern access to it should be formulated primarily by epidemiologists. I suggest that the M.R.C. was mistaken not to keep in mind that justice, in the drawing up of a code of conduct as in a law court, must not merely be done but must be seen to be done; or to put it a shade differently, that men should never be the sole legislators in respect of a cause to which they are very interested parties.

The M.R.C. has kept this in mind in relation to its new Standing Committee on the Use of Medical Information for Research, which is to contain representatives of the public. That committee, however, is to carry out its functions in accordance with the principles contained in the present statement. If it is thought appropriate that members of the public should help to interpret the principles, why was it not thought proper that they should help to enunciate them? After all, it is confidential data concerning individual members of the general public that we are talking about, and neither the M.R.C. nor epidemiologists nor anybody else, apart from the personal physicians involved, can claim a prescriptive right to any kind of access to it. Eminence in no way blurs the distinction between a right and a privilege. I suggest the M.R.C. should think again and give its standing committee a different and a wider mandate.—I am, etc.,

S. BRADSHAW

Hereford

An Integrated Child Health Service

SIR,—I read with interest the article by Dr. F. N. Bamford and Professor J. A. Davis (*Supplement*, 20 January, p. 20), particularly those aspects which relate to school doctors. In support of the suggestions made by the authors and to illustrate how the role can be filled by a married woman doctor working on a part-time basis, I write to describe the arrangements for school work which apply here in an expanding town of about 12,000 inhabitants.

Early in 1968 the group practice for this area and the local authority invited me to do local school work in liaison with the practice as part of a local authority pilot scheme for attachment of local general practitioners to health clinics. I was to be employed by the local authority on a sessional basis during school terms, aided by a health assistant and a clerk. I had just returned from abroad with young children of my own and had done paediatric work previously at home and overseas. With one weekly session, I began. Despite policy changes and administrative difficulties I have remained "the school doctor" here ever since and have recently been able to combine this with a limited amount of hospital paediatrics.

This arrangement provides continuity of medical service in the same schools, with obvious advantages. With head teacher and doctor well acquainted, problems can often

be raised and sorted out without delay or referral; familiarity with teaching staff and children's backgrounds clarifies situations more readily. By agreed arrangement with the local general practitioners, referrals are made directly to consultants in the more specialized fields—unless special considerations arise. Follow-up, in application to school problems, is assured, which is particularly helpful for the handicapped or deprived child. I am not involved in school buildings or kitchen hygiene, although occasionally it has been possible to help in these fields if help or advice has been sought.

Only 5-year-old children are seen routinely in this county now; older children attend by a system of selection. It is therefore mainly with the infant and primary schools that the arrangements described above are concerned, and it is there that they seem most appreciated by head teachers, staff, and parents. Indeed parents (usually mothers) attend "school medicals" almost unfailingly and with interest—sometimes with over-enthusiasm, when sick children have been brought from bed at home to attend.

After five years in this field I am sure that school medical work should not be limited to screening, immunization, filling in forms, and collecting statistics, as some might have it now. It can offer much more both to the child and to the doctor and so could play a useful and practical part in an integrated health service. Further, bearing in mind the present efforts to attract married women doctors back into medicine, I suggest that school medical work could be an ideal field for the doctor with paediatric experience and young children of her own. The hours of work are convenient and the doctor is closely in touch with children and children's problems. At the risk of an outcry by women's lib. and male doctors in school medical work now, may one not envisage the school doctor of the future as a married woman, with children, employed on a part-time contract, attached to a paediatric unit or a local group practice or both, trained in paediatrics, and having an interest in and concern for children—both other people's and her own?—I am, etc.,

HELENOR PRATT

Marlow, Bucks

SIR,—In their interesting paper Dr. F. N. Bamford and Professor J. A. Davis (*Supplement*, 20 January, p. 20), referring to the duties of school doctors, note that "they are expected to give advice about school buildings and kitchen hygiene, to examine teachers to ascertain their medical fitness for employment and supperannuation, and to take part in health education."

The authors state that it is wrong that school doctors should be diverted by the problems of environmental health and occupational medicine from their most essential study of child development. This is such an extraordinary statement that it suggests that the authors have a limited idea of what school medicine entails. It is not possible, in my opinion, to separate the consideration of school buildings and kitchen hygiene from the general health and welfare of the child in the school and, from personal experience, I consider that unsatisfactory conditions can