operation. The tea goes into the drainage bag, but the pleasure of it going down is not lightly to be dismissed. Total cystectomy, which in my experience is followed by at least 4-5 days ileus, is now my most common reason for gastrostomy. To have a tube down your nose for so long is intolerable, and if removed prematurely it must be replaced. With a gastrostomy the nurse has only to empty the bag when full and measure the contents and to insert or remove a spigot. Ward sisters always say a gastrostomy causes less work.

I have seen no complication clearly and directly attributable to gastrostomy. I believe gastrostomy increases safety by efficiently and quickly removing swallowed air which would otherwise pass into the small bowel. The advantages here are obvious—particularly when a bad aerophagist is seen blowing up the bag like a balloon. I am aware of three disadvantages to gastrostomy. Firstly, though it can be performed under local anaesthetic at any time it is preferable to insert the tube at the time of operation. Thus gastrostomy may be used more often than is strictly necessary. Secondly, I prefer to leave the tube in situ for 10 days. This means that the patient could spend an extra day or two in hospital. Thirdly, if the patient drinks large amounts immediately after operation there may be a fairly large loss of potassium. This has to be borne in mind when planning the intravenous therapy.

I believe that there is no longer any case for adding nasogastric aspiration to the discomforts of operation.—I am, etc.,

D. M. ESSENHUGH

Department of Urology,
Newcastle General Hospital,
Newcastle upon Tyne

Dangerous Patients

SIR,—There is little doubt that it is that part of the Mental Health Act, 1959, dealing with restricted patients which causes psychiatrists working in conventional hospitals most trouble and concern. Doctors treating mentally disturbed offenders have an intense dislike of restriction orders and it is a truism that these orders restrict doctors as well as their patients.

In your forthright and perceptive leading article (3 February, p. 247) on the Aarvold Report,¹ you imply that all patients subjected to special restrictions as imposed by sections 60/65 are dangerous. In fact this is not always so. Indeed, one of my patients was placed on a restriction order, without limit of time, as the result of being found guilty of a minor offence of theft. This man was classified as severely subnormal and was very simple and quiet, but was rather gullible and had been induced by his co-offenders to stand watch while they actually stole some material. However, he had been convicted by the same court on two previous occasions and one can only assume that he was placed on a restriction order because the court thought him to be a nuisance to society rather than a menace. This in practice is becoming so commonplace that one is reluctant to recommend a section 60 order in certain cases because of the fear that the court will add on a section 65 order. It should therefore be possible for the doctor to state in his report to the court

whether or not he is prepared to treat the offender in his hospital under section 65, and also he should be able to advise the judge as to the length of time the restriction order should remain in force, if the court feels that a restriction order is essential for the protection of the public.

With regard to patients granted conditional discharge, the position can become quite ludicrous. For instance, one of my patients classified as subnormal was placed on conditional discharge in September 1971. In November 1972 he pleaded guilty to various offences and was sentenced to 12 months imprisonment. Both I and the prison medical officer reported to the court that he could no longer be considered to be subnormal within the meaning of the Mental Health Act, and he was consequently dealt with by the normal processes of law. However, this man is still on conditional discharge and so after he has served his sentence he could be returned to hospital. We shall then have the ridiculous position of the responsible medical officer having to treat a person as subnormal in spite of the fact that he has previously reported that he is no longer suffering from this condition. In cases such as this, I believe that the conditionally discharged patient should be discharged absolutely.

Last year I drew attention² to the divergence of opinion with regard to the interpretation of a part of the Mental Health Act. In this letter I have pointed out further aspects of the working of the Act which require clarification. No doubt other psychiatrists will have formed their own views about sections of the Mental Health Act which require alteration. I therefore feel that now is an opportune time for a complete review of the Mental Health Act.—I am, etc.,

A. I. ROITH

Monyhull Hospital,
Birmingham

¹ Home Office Report on the Review of Procedures for the Discharge and Supervision of Psychiatric Patients Subject to Special Restrictions, Cmnd. 5191. London, H.M.S.O., 1973.

² Roith, A. I., *Lancet*, 1972, 1, 1389.

Working of the Mental Health Act

SIR,—The letter from Mr. J. A. Cooke (3 February, p. 292) is not clear on all details, and I wonder if I might put the following points, which I would consider axiomatic.

(1) Though the four specific types of mental disorder described in section 4 of the Mental Health Act 1959 and used for orders under section 26 are separate and distinguishable, there is nothing to stop a clinician using any two of them at the same time. Indeed, clinicians regularly combine "subnormality" and "psychopathic disorder"—not only for the purpose of section 26 but also for that of section 60.

(2) Ever since the Act came into being clinicians have regarded mental disorder resulting from organic disease of the brain, including brain damage, as constituting "mental illness," a term which, as Mr. Cooke says, is undefined in the Act and therefore designed to be pliable.

(3) If this condition secondary to brain damage or other organic disease is mainly manifested by "seriously irresponsible conduct," I can see no reason why the section 26 or section 60 medical recommendation should not be on the ground of "mental ill-

ness combined with psychopathic disorder." The order would indicate that the patient had been made mentally ill by the organic disease but that his illness was mainly appearing in the form of a disturbance of personality rather than as notable dementia.

I am sure in fact that no mental disorder secondary to such organic disease can occur without there being a degree of dementia which would make the use of the term "mental illness" still more justifiable.

(4) Provided that the diagnosis of "mental illness" preceded that of "psychopathic disorder" on the recommendation, the clause under "psychopathic disorder" whereby a person cannot be detained beyond the age of 21 would not apply.

I hope that what I am saying is in accord with the rather obscure last paragraph of Mr. Cooke's letter. Indeed, I think it could be argued that a simple classification of "mental illness" in the sort of case in question could not be used without the addition of the words "or psychopathic disorder," even when personality disturbance was the main feature of the condition. If, however, the clinician has doubts, I cannot see why he need stop using both. In this connexion, I have just seen a woman in her late sixties who, after a stroke, has become an alcoholic, a compulsive stealer of drugs and generally disinhibited in behaviour, grossly egocentric, and quite insouciant. It is likely that she will soon need to be in hospital under section 26 of the Act and I have no doubt that the classification of "mental illness" will properly apply to her.—I am, etc.,

SEYMOUR SPENCER

Oxford

Myocardial Infarction and Pulmonary Thromboembolism

SIR,—Dr. Helen H. Tucker and her colleagues (6 January, p. 10) report a low clinical incidence of pulmonary embolism in their group of patients who were mobilized in the first few days after their myocardial infarction. Their reported incidence is probably an underestimate.¹ Immobility is the commonest single cause of venous thrombosis.² While early mobilization would be expected to reduce the incidence of this condition, it has been well documented³ that most thrombotic episodes occur during the first few hours of immobility.

In Dr. Tucker's series, as in others, there was a high mortality rate in patients who developed atrial fibrillation after myocardial infarction. This may be because a pulmonary embolus caused the arrhythmia. A cardiac arrhythmia may be the only manifestation of pulmonary embolization.⁴ We have studied six patients whose principal objective abnormality was a cardiac arrhythmia and in whom the presence of pulmonary embolism was demonstrated angiographically.

Unexplained dyspnoea is used as an indication for the treatment of cardiac failure by Dr. Tucker and her colleagues. It is likely that at least in some of these cases pulmonary embolism accounts for this symptom. It is well known that pulmonary emboli do not necessarily cause any radiographic or electrocardiographic abnormality and that the clinical diagnosis of venous thrombosis is very inaccurate.

There is a tendency to attribute any cardiorespiratory abnormality occurring in

patients with preexisting heart disease to a deterioration in the underlying cardiac condition. We would suggest that, if there is no positive evidence for this explanation, then pulmonary embolism should be considered as a possible cause.—We are, etc.,

W. J. WINDEBANK
F. MORAN

Royal Infirmary,
Glasgow

- ¹ Sasahara, A. A., Cannilla, J. E., Morse, R. L., Sidd, J. J., and Tremblay, G. M., *American Journal of Cardiology*, 1967, 20, 10.
- ² Gibbs, N. M., *British Journal of Surgery*, 1957, 45, 209.
- ³ Flanc, C., Kakkar, V. V., and Clarke, M. B., *British Journal of Surgery*, 1968, 55, 742.
- ⁴ Littmann, D., in *Pulmonary Embolic Disease*, ed. A. A. Sasahara and M. Stein, p. 186. New York, Grune and Stratton, 1965.

Fractured Lippes Loop and Pregnancy

SIR,—We have used the Lippes loop for eight years and have experience of about 1,000 cases. Our first case of fracture of the loop in utero occurred in December 1972. At about the same time there appeared two reports of similar observations.^{1,2}

The Lippes loop was inserted in 1966 after a third induced abortion, the patient then being 37 years old. A yearly check-up showed no abnormality, and the loop caused no complaints. In November 1972 menstruation ceased, and in December pregnancy became obvious. The patient applied for interruption. Before evacuation of the uterus we attempted to remove the loop, but when the indicator suture was pulled out only a small part of the loop was found to be attached to it. A radiograph showed that three fragments of the loop were still in the uterus (see fig.). These parts were removed under general anaesthesia after the cervix had been dilated. The fractured parts fitted together.

After several years of use chemical changes in an intrauterine device may produce loss of rigidity that may lead to fracture. Fracture may affect the contraceptive action, as was evidently the case in this patient, who became pregnant at the age of 42 after six years of uneventful use of the device.—We are, etc.,

Z. DOMÁNY
M. HANCSÓK

Budapest, Hungary

- ¹ Last, P. A., *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1972, 79, 190.
- ² Lang, J. R., *British Medical Journal*, 1972, 4, 795.

Childhood Leukaemia and Pregnancy Viraemia

SIR,—In answer to Dr. C. M. D. Edmonds (27 January, p. 235), given an expectation of 0.40 deaths per survey, no deaths would be found in about 67% of such surveys, one death in 27%, two deaths in 5%, and over two deaths in the remaining 1%; these percentages are given as individual terms of the Poisson distribution. Thus to find no deaths in one such survey is hardly unusual.

But where the expectancy of a death is 0.15 per survey, two deaths would be found in less than 1% of surveys; in other words, to find two deaths in the one and only survey is significant at the 1% level.

The accumulation of large numbers is not necessary, in all circumstances, to suggest statistical or commonsense significance. To win a major pools dividend once is, perhaps, fortuitous: to win twice, suspicious.—I am, etc.,

R. H. LINDENBAUM

Oxford

AHF-related Protein and Precipitation Reactions

SIR,—Dr. H. Ekert and others (30 December, p. 789) raised the question of the specificity of the antiserum we used when we investigated patients with von Willebrand's disease (5 August, p. 317). We did not find IgM or lipoprotein by means of specific antisera in the purified antihemophilic factor (AHF) which we used for immunization. The antiserum used was absorbed with one-tenth volume of normal plasma. The anti-AHF antibody was present in such large amounts that this absorption did not appreciably affect its titre. No difference was observed when the patients were tested with unabsorbed antiserum or antiserum absorbed with one-tenth volume of von Willebrand plasma.

We have recently had the opportunity of comparing our antiserum with those of other workers in this field (Drs. Bouma, of Utrecht; Meyer, of Paris; Kernoff, of Oxford; Prentice, of Glasgow; and Trobisch, of Marburg). All antisera gave lines of complete identity in double diffusion tests, indicating that one and the same von Willebrand antigen is determined by all. There is no doubt, however, that the antigen is heterogeneous, as demonstrated by cross-immuno-

electrophoresis, and quantitative estimation by the Laurell technique must be carefully performed. An improved method is, however, under way.

We did not regularly test our patients with an inhibitory coagulation test, as the results of such methods are somewhat inconsistent.—We are, etc.,

LARS HOLMBERG
INGA MARIE NILSSON

Allmänna, Sjukhuset,
Malmö, Sweden

Sympathomimetic Amines and Antidepressant Agents

SIR,—The very interesting and important research described by Dr. A. J. Boakes and others (10 February, p. 311) is reassuring in its suggestion that two monoamine oxidase inhibitors do not harmfully potentiate the effects on the blood pressure and pulse rate of four sympathomimetic amines. However, the authors' finding that imipramine produces a greater potentiation of these effects than do the monoamine oxidase inhibitors appears to be contrary to clinical experience.

Possibly this apparent anomaly results from the experimental design. One would expect clinical pharmacologists, forewarned by their knowledge of the risks of drug interaction, to be scared by lying down for three hours or so while four amines in succession are infused into them immediately after they have taken antidepressives. Their fear could magnify any cardiovascular actions of the amines under these conditions. Habituation would be likely to reduce the emotional reactions and, if I interpret the methodology rightly, each subject had infusions on four occasions. The authors do not say which they gave first, imipramine or the monoamine oxidase inhibitors, but I think it must have been the tricyclic drug because they mention it first in their description of the method and most clinicians believe that to give tricyclics before monoamine oxidase inhibitors is safer than vice versa.

If this sequence was indeed the one in which the experiment proceeded, it could perhaps explain also the statistically significant increase of resting blood pressure which the authors observed during the preliminary uncomplicated administration of imipramine and which likewise is the reverse of clinical experience. One would surmise that while the subjects were taking this drug in the early stages of their ordeal they were particularly apprehensive about what was in store for them.

I realize that Dr. Boakes and his colleagues probably dismissed additions to their design as making it so vast and complicated that they could not have put it into practice, but if they had given normal saline without amines as a fifth substance during each infusion they might have been able to distinguish pharmacological from psychological effects of interaction, whereas they could have thrown some light on the concurrence of imipramine medication and hypertension by a double-blind administration of a placebo instead of an antidepressive to another group of subjects undergoing the whole procedure.

I doubt whether from this research we need seriously suspect imipramine of inter-