- Clark, T. J. H. (1972). *Lancet*, 1, 1361.
 Greig, W. R., Maxwell, J. D., Boyle, J. A., Lindsay, R. M., and Browning, M. C. K. (1969). *Postgraduate Medical Journal*, 45, 307.
 Lal, S., Harris, D. M., Bhalla, K. K., Singhal, S. N., and Butler, A. G. (1972). *British Medical Journal*, 3, 314.
 Linder, W. R. (1964). *Archives of Internal Medicine*, 113, 655.
 Murphy, B. E. P. (1967). *Journal of Clinical Endocrinology and Metabolism*, 23, 973.

- Raffle, E. J., and Frain-Bell, W. (1967). *British Journal of Dermatology*, 79, 487.
 Scoggins, R. B., and Kliman, B. (1965). *New England Journal of Medicine*, 273, 831.
 Westerhof, L., van Ditmars, M. J., der Kinderen, P. J., Thijssen, J. H. H., and Schwartz, F. (1970). *British Medical Journal*, 4, 534.
 Westerhof, L., van Ditmars, M. J., der Kinderen, P. J., Thijssen, J. H. H., and Schwartz, F. (1972). *British Medical Journal*, 2, 195.

MEDICAL MEMORANDA

Behçet's Disease with Decreased Fibrinolysis and Superior Vena Caval Occlusion

T. CHAJEK, M. FAINARU

British Medical Journal, 1973, 1, 782-783

Thrombophlebitis is a recognized manifestation of Behçet's disease but thrombotic occlusion of large vessels is rare (Haim *et al.*, 1971; Enoch *et al.*, 1968; Kansu *et al.*, 1972). We report here two cases with superior vena caval obstruction and defective fibrinolysis which were treated successfully with fibrinolytic agents.

Case 1

A 30-year-old Arab man was admitted to hospital because of increasing swelling of the face and neck over the past six weeks. He had suffered from recurrent oral and scrotal ulcers for two years, and 18 months previously he had developed superficial thrombophlebitis of the left leg. Three months after that he had noticed engorgement of the superficial veins of his left upper limb. Ten months later a painful swelling in the right cervical region developed, spreading rapidly to the entire neck and causing puffiness of the face.

On admission the classical signs of superior vena caval obstruction, with prominent venous collaterals over the chest and abdomen, were present. Ophthalmological examination showed evidence of uveitis and pronounced dilatation of the retinal veins. The right supraclavicular space was occupied by a dusky, painful mass, and there were several superficial oral and scrotal ulcers. The heart, lungs, abdomen, and nervous system were normal. Total serum protein was 7.2g/100 ml, with a slight increase in alpha₂ globulin fraction (16.8%). No cryoglobulin was detected. A chest radiograph showed widening of the superior mediastinum. A right brachial venogram showed widening and obstruction of the subclavian vein, very dilated axillary and brachial veins, and prominent venous collaterals (see fig.). Bone marrow aspirate showed mild eosinophilia and plasmacytosis. Biopsy of material from a scrotal ulcer showed nonspecific inflammation. Lymph node and liver biopsy were normal. The supraclavicular mass was found on exploration to be a thrombosis of the large veins, which extended into the mediastinum. No tumour or inflammatory process was found. Biopsy of material from one of the thrombotic veins showed chronic inflammation of the wall with a large organized adherent thrombus.

The results of coagulation studies are shown in the table. The clotting time, prothrombin time, partial thromboplastin time, factor II, V, VII, and X, and the platelet aggregation were all within normal limits on admission.

Department of Medicine B, Hadassah University Hospital, Jerusalem, Israel

T. CHAJEK, M.D., Resident
 M. FAINARU, M.D., Second Assistant

While in hospital the patient developed erythema nodosum-like lesions on both arms, red papules and pustules after each needle prick, and thrombophlebitis after every venepuncture. He was given Varidase 12 tablets daily by mouth (each tablet containing streptokinase 10,000 U and streptodornase 2,500 U). After 48 hours the general condition greatly improved, the swelling of the head and neck decreased, and the supraclavicular mass disappeared. After 14 days Varidase treatment was discontinued for one week: the patient's condition deteriorated and fibrinolysis decreased. When treatment was resumed there was again a clinical remission and plasma fibrinolytic activity increased. The patient was discharged on a maintenance dose of 12 tablets of Varidase daily, and 16 months later he was well and showed no signs of thrombophlebitis.

Right brachial venogram in case 1.

Case 2

A 22-year-old Arab man was admitted because of fever and progressive swelling of the face of four weeks' duration. For the past four years he had had recurrent aphthous stomatitis, scrotal ulcers, and conjunctivitis accompanied by febrile episodes. Nine months before admission he had noticed periodic symmetrical swelling of the neck and dilatation of the superficial veins of the chest and abdomen. He presented the classical signs of superior vena caval occlusion. He had superficial ulcers, 5-7 mm in diameter, on both lips, tongue, and scrotum. Chest radiography showed widening of the right superior mediastinum with some opacity of the right lung parenchyma and a small amount of fluid in the right pleural cavity. The results of relevant coagulation studies are given in the table. The other tests were within normal limits.

Treatment was with intravenous fluids. Severe thrombophlebitis developed at the site of one of the infusions, and extended into the axilla. Streptokinase 300,000 U infused over the first 24 hours produced a dramatic improvement of the thrombophlebitis and correction of the fibrinolytic defect (see table). Treatment was continued with oral Varidase, and six months later he was well, except for occasional superficial oral ulcers, and the euglobulin lysis time was within normal limits.

Results of Coagulation Studies

	Fibrinogen (mg/100 ml)	Factor VIII (%)	Euglobulin Lysis Time (hours)
Normal range	Case 1 200-400	50-150	3-4
On admission	550	1,090	>24
On Varidase treatment (14 days)	480	—	2-3
Without treatment (for seven days)	450	1,000	>8
On Varidase treatment .. .	460	290	2-7
On admission	Case 2 580	780	>24
24 hours after treatment with intravenous streptokinase ..	202	375	0-5
On oral Varidase treatment ..	456	280	3-5

Comment

Few studies of the clotting mechanism in Behçet's disease have been reported. Cunliffe and Menon (1969) investigated a case with deep thrombophlebitis and repeated pulmonary emboli. They found that despite a normal coagulation and fibrinolytic mechanism the patient improved on treatment with phenformin and ethyloestrenol, which enhanced fibrinolysis. In the present two patients plasma fibrinolytic activity was strikingly

decreased and the levels of fibrinogen and factor VIII increased. We do not think these coagulation defects have been described in Behçet's disease before. In both cases giving streptokinase reduced the euglobulin lysis time to normal and greatly improved the thrombophlebitis. Phlebitis and decreased fibrinolysis recurred in the first case when treatment was discontinued.

The aetiology of Behçet's disease is still unknown, but vasculitis seems to be the underlying cause of most of its diverse manifestations (Dowling, 1961). The recurrent thrombophlebitis may be secondary to or enhanced by decreased fibrinolysis. Because of the successful treatment with streptokinase in the cases reported here we suggest that fibrinolytic agents may be indicated in cases of Behçet's disease with superficial or deep thrombophlebitis.

References

- Cunliffe, W. J., and Menon, I. S. (1969). *Lancet*, 1, 1239.
Dowling, G. B. (1961). *Proceedings of the Royal Society of Medicine*, 54, 101.
Enoch, B. A., Khoo, T. C. L., Castillo Olivares, J. L., Grainger, R. G., and Henry, L. (1968). *Postgraduate Medical Journal*, 44, 453.
Haim, S., Barzilai, D., and Hazani, E. (1971). *British Journal of Dermatology*, 84, 283.
Kansu, E., et al. (1972). *Quarterly Journal of Medicine*, 162, 141.

Facial Palsy as the Sole Presenting Symptom of Parotid Carcinoma

M. J. M. BLACK

British Medical Journal, 1973, 1, 783

Facial palsy is by no means an unusual feature of an advanced carcinoma of the parotid salivary gland. The two most common presenting features of the disease are, however, the presence of a mass and pain. Facial palsy may be associated with these symptoms, and the significance of this palsy with regard to the diagnosis of malignancy is well recognized.

Facial palsy may, however, precede the appearance of other features by many months. Frazell (1954) stated "In a few patients seen by us, such palsy was the initial and only complaint for weeks or months prior to discovery of the tumour."

It is our experience, in the combined plastic surgery radiotherapy clinic, that this mode of presentation is more common than we had previously appreciated. From 1969 to 1971 three cases (see below) out of 21 (14%) of parotid carcinoma presented in this way.

Case Histories

Case 1.—An 80-year-old woman presented with a history of weakness of the left side of the face for one week, associated with some pain in the region of the left side of the mandible. An almost complete facial palsy was present but there was no evidence of causation. Bell's palsy was diagnosed. The palsy did not recover. Thirty-three months later she was referred for treatment of corneal exposure problems as a result of the complete palsy. At examination a small ulcer was found near the lobule of the left ear, with an underlying mass. Biopsy showed a poorly differentiated carcinoma of parotid gland origin. A cervical lymph node contained metastatic tumour. Lateral tarsorrhaphy was performed and radiotherapy started.

Case 2.—A woman aged 41 was under regular review after the treatment of basal cell carcinomata of the skin. A right-sided

facial palsy developed and was diagnosed as Bell's palsy. Sixteen months later a parotid mass was first apparent. Parotidectomy was performed and a parotid carcinoma of squamous cell type was found.

Case 3.—A 58-year-old woman presented with a left-sided facial palsy which was diagnosed as a Bell's palsy. It did not recover. Seven months later an ill-defined firm mass was found between the jaw and mastoid process. A biopsy specimen showed normal parotid gland tissue. Plain x-ray examination of the region and parotid sialography showed nothing abnormal. Ten months later a further mass became apparent and was then associated with pain in the left side of the face. Exploration showed a tumour of the parotid gland of the squamous cell type. Radiotherapy was given.

Comment

The first two cases are typical of the problem, the delay between diagnosis of facial palsy and eventual discovery of the parotid neoplasm being 33 months and 16 months respectively. The third case shows how difficult it may be to confirm the diagnosis even when suspicion has been aroused. It is our belief that at the time of biopsy the clinically apparent mass was normal parotid gland tissue displaced laterally by a more deeply lying neoplasm.

In case 1 pain was mentioned at the time of initial examination. Perhaps its significance should have been realized. It seems doubtful, however, if any more direct evidence of a neoplasm could have been found at that time.

When a patient presents with a facial palsy without other diagnostic features it is more than likely that the diagnosis of Bell's palsy will be made. In many cases this must be diagnosis by exclusion. If a parotid neoplasm is the underlying cause it will escape detection until other features become apparent.

The purpose of this report is to increase the awareness of the possibility of parotid neoplasm as a cause of the apparent Bell's palsy, and earlier diagnosis may well be achieved by critical follow-up of all cases diagnosed initially as Bell's palsy.

In cases where the palsy recovers, the label of Bell's palsy could be justified. When it does not recover, the label should be "facial palsy—cause not yet diagnosed."

I would like to thank Mr. W. M. Ross, consultant radiotherapist, for his advice on the preparation of this article.

References

- Frazell, E. L. (1954). *Cancer*, 7, 637.

Newcastle University Hospitals, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP

M. J. M. BLACK, M.B., F.R.C.S., Registrar in Plastic Surgery