more occasions has a 9 in 10 chance of being delivered of a severely affected infant. At the same time it should be noted that about one-third of severe cases will be missed, so that HPL levels cannot be relied on as the sole diagnostic criterion. Nevertheless, when the simplicity and low cost of this test is taken into account there can be little doubt that serial levels of HPL are a valuable adjunct to other methods of diagnosis in rhesus isoimmunization.

The assays were supported by grants from the Medical Research Council and the Board of Governors of St. Bartholomew's Hospital and were performed by Miss Linda Howard. We are grateful to the medical and nursing staff of University College Hospital and St. Bartholomew's Hospital for their co-operation in this study, and to Professor D. V. I. Fairweather for his encouragement and advice.

References

Fairweather, D. V. I., Billewicz, W., Loraine, J. A., and Bell, E. T. (1972). *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 79, 97.

Genazzani, A. R., Aubert, M. L., Casoli, M., Fioretti, P., and Felber, J. P. (1969). *Lancet*, 2, 1385.
 Genazzani, A. R., et al. (1971). *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 78, 577.
 Haour, F., Cohen, M., and Bertrand, J. (1971). *Revue Européenne d'Études Cliniques et Biologiques*, 16, 124.
 Josimovich, J. B., Kosor, B., and Mintz, D. H. (1969). *Foetal Autonomy*, ed. G. E. W. Wolstenholme and M. O'Connor. p. 117. London, Churchill.
 Josimovich, J. B., Kosor, B., Boccia, L., Mintz, D. H., and Hutchinson, D. L. (1970). *Obstetrics and Gynecology*, 36, 244.
 Letchworth, A. T., Boardman, R. J., Bristow, C., Landon, J., and Chard, T. (1971). *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 78, 542.
 Pridmore, B. R., Robertson, E. G., and Walker, W. (1972). *British Medical Journal*, 3, 136.
 Samaan, N. A., Bradbury, J. T., and Goplerud, C. P. (1969). *American Journal of Obstetrics and Gynecology*, 104, 781.
 Singer, W., Desjardins, P., and Friesen, H. G. (1970). *Obstetrics and Gynecology*, 36, 222.
 Spellacy, W. N., Teoh, E. S., and Bui, W. C. (1970). *Obstetrics and Gynecology*, 35, 685.
 Spellacy, W. N., Teoh, E. S., Bui, W. C., Birk, S. A., and McReady, S. A. (1971). *American Journal of Obstetrics and Gynaecology*, 109, 588.
 Ursell, W., Brudenell, M., and Chard, T. (1973). *British Medical Journal*, 2, 80.
 Varma, K., Driscoll, G. S., Emerson, K., and Selenkow, H. A. (1971). *American Journal of Obstetrics and Gynecology*, 38, 487.

Plasma Testosterone and Testosterone Binding Affinities in Men with Impotence, Oligospermia, Azoospermia, and Hypogonadism

DAPHNE M. LAWRENCE, G. I. M. SWYER

British Medical Journal, 1974, 1, 349-351

Summary

Mean plasma testosterone levels ($\pm$ S.D.), using Sephadex LH-20 and competitive protein binding, were 629 ± 160 ng/100 ml for a group of 27 normal adult men, 650 ± 205 ng/100 ml for 27 impotent men with normal secondary sex characteristics, 644 ± 178 ng/100 ml for 20 men with oligospermia, and 563 ± 125 ng/100 ml for 16 azoospermic men. None of these values differ significantly. For 21 men with clinical evidence of hypogonadism the mean plasma testosterone ($\pm$ S.D.), at 177 ± 122 ng/100 ml, differed significantly ($P < 0.001$) from that of the normal men.

The mean testosterone binding affinities (as measured by the reciprocal of the quantity of plasma needed to bind 50% of ^3H -testosterone tracer) were similar for normal, impotent, and oligospermic men. Though lower for azoospermic men the difference was not significant ($P > 0.1$). For 12 of the 16 hypogonadal males the testosterone binding affinity was normal, but raised binding affinities, similar to those found in normal adult females or prepubertal boys (about twice normal adult male levels), were found in four cases of delayed puberty. These findings help to explain why androgen therapy is usually useless in the treatment of impotence.

Introduction

It is common clinical practice to treat impotent men with androgen preparations either orally, by injection, or by implan-

tation but usually without noticeable clinical improvement. There appear to be few published data on plasma testosterone levels in impotent males, though there are reports that these men have a lower urinary testosterone excretion. Thus Ismail *et al.* (1970) compared 28 impotent men with 14 normal sexually active males and found the overall mean urinary testosterone excretion of the former to be significantly lower than that of the normal group ($P < 0.001$). Cooper *et al.* (1970) classified their impotent patients into predominantly psychogenic or predominantly constitutional groups and found the mean urinary testosterone of the latter group to be significantly lower ($P < 0.005$). Rosen and Weintraub (1971) reported on serum testosterone and luteinizing hormone concentration in idiopathic oligospermia and found a mean serum testosterone value of 560 ng/100 ml for 17 idiopathic oligospermic or azoospermic men aged 18 to 36. This was lower than the mean for normal men, 620 ng/100 ml, but the difference was not statistically significant ($P > 0.1$). The luteinizing hormone concentration was similar in both groups.

We have investigated the plasma testosterone concentration in normal and impotent men and in men with oligospermia and azoospermia, as well as in a group of men with clinical evidence of hypogonadism. We have also evaluated testosterone binding affinities in most of the subjects.

Subjects

The 27 normal men were hospital personnel aged 20 to 63 (mean 37 years), sexually active, and normospermic. Details of the 27 men with impotence aged 21 to 62 (mean 43 years) are given in the table.

The 20 men with oligospermia (sperm counts persistently less than 20 million/ml) aged 22 to 44 (mean 30 years) and the 18 men with azoospermia aged 25 to 34 (mean 29 years) had normal external genitalia and secondary sex characteristics on clinical examination. They were attending the fertility or endocrine clinics at University College Hospital. The group

Department of Obstetrics and Gynaecology, University College Hospital Medical School, London WC1E 6HX
 DAPHNE M. LAWRENCE, B.Sc., PH.D., Senior Biochemist
 G. I. M. SWYER, D.M., F.R.C.P., Consultant Endocrinologist

Clinical Data on Impotent Men

Case No.	Age	History of Impotence		Seminology	Children	Other Factors	Androgen Therapy
		Length in Years	Coitus before Onset of Impotence				
1	21	1½	Normal, unmarried	Normal	0	—	—
2	25	3	Inadequate erection	Normal	0	—	—
3	31	7	Normal	Normal	2	Wife in Pakistan and only 3 visits in 8 years Testicles rather high in scrotum	Methyl testosterone; no effect
4	31	8	Normal	Normal	5		
5	32	9	Unmarried. Unable to have coitus but can masturbate	Azoospermia	0	Atrophic right testicle after ? inflammation ? torsion	Fluoxymesterone; no effect
6	33	9	Normal, gradual onset	Oligospermia	0		
7	33	4	Normal. Can masturbate but reduced or absent orgasm	Normal	0	Diabetic for 11 years	Testosterone injections; no benefit
8	36	2	Normal	Normal	2		
9	36	10	Impotence from start of marriage		0	Divorced after 7 years. Remarried but impotence with second wife	Testosterone injections; no benefit
10	37	7	Erections only with manual stimulation; then increasing difficulty	Normal	2		
11	38	4	Satisfactory; gradual loss	Normal	0	Car accident 4 years ago. Ankylosing spondylitis	
12	40	13	No coitus. Tried for 4-5 years of marriage	Normal	0		
13	42	4	Normal	Normal	2	Divorced	
14	44	20	No erection before or after marriage	Normal	0		
15	48	5	Normal. Impotence since divorce 5 years ago		2	Wife died; wants to remarry	
16	49	1½	Normal	Normal	0		
17	49	1	Normal	Oligospermic	0	Pain in left groin Rather obese Alcoholic	
18	50	1	Normal	Normal	4		
19	52	1	Normal	Normal	2	Testicles retracted at onset of impotence. Complained of flushes and sweating.	
20	53	3	Normal	Oligospermic	5		
21	53	4	Normal	Oligospermic	1	Panic states Scrotal hernia	
22	55	2	Normal	Normal	0		
23	55	8	Normal	Normal	3	Wife died 8 years ago. Right testicle excised to remove cyst. Prostatectomy 4 years ago.	Fluoxymesterone; no effect
24	56	1	Normal	Oligospermia	0		
25	56	1	Normal	Normal	0		
26	61	20	Normal for 20 years	Normal	2		
27	62	8	Normal		2		

with clinical evidence of hypogonadism were aged 14 to 45 (mean 29 years), details of whom have been given previously (Lawrence and Swyer, 1973).

Materials and Methods

Blood was drawn into heparinized tubes between 09.30 and 12.00 and between 13.30 and 16.00 hours from both the patients and the normal controls. The significance of a circadian rhythm in testosterone levels with higher values expected in the morning and lower levels in the late afternoon was appreciated but it is believed that the scatter of times of venesection was similar for all the groups of men studied.

The plasma testosterone was extracted after separation on short columns (13 by 0.7 cm) of Sephadex LH-20 and measured by competitive protein binding using ammonium sulphate precipitation of the bound testosterone. The method used was that of Lawrence and Swyer (1973) as modified from those of Murphy (1970) and Mayes and Nugent (1968).

Testosterone binding affinity was measured by the method of August *et al.* (1969). In this method testosterone affinity is expressed as 1/S, the reciprocal of the quantity of plasma needed

to bind 50% of 0.5 ng of 1,2 ³H-testosterone (specific activity 40-45 Ci/mmol). Florisil was used to remove the unbound testosterone.

Results

The plasma testosterone levels in the 27 normal males ranged from 408 to 1,022 ng/100 ml (mean ($\pm$ S.D.) 629 $\pm$ 160). The range for the 27 impotent men was 352 to 1,147 ng/100 ml (mean 650 $\pm$ 205). The 20 men with oligospermia had values of 368 to 1,060 ng/100 ml (mean 644 $\pm$ 178), and the 18 men with azoospermia 365 to 766 ng/100 ml (mean 563 $\pm$ 125). The range for the group of 21 men with hypogonadism was 30 to 378 ng/100 ml (mean 177 $\pm$ 122).

The individual results are plotted in fig. 1. Though the scatter was greater in the impotent group the difference between the mean values of this and the normal group was not significant ($P > 0.7$). Similarly, the mean value for the men with oligospermia was not significantly different from that of the normal group though the mean value for the azoospermic group was lower, but again the difference was not statistically significant ($P > 0.1$). On the other hand, for the hypogonadal group the

FIG. 1—Plasma testosterone values in normal men and those with either impotence, oligospermia, azospermia, or hypogonadism.

FIG. 2—Binding affinities for testosterone (expressed as $1/S$, reciprocal of quantity of plasma needed to bind 50% of 3H -testosterone tracer) in groups of normal, impotent, oligospermic, azospermic, and hypogonadal subjects. O = Delayed puberty.

mean plasma testosterone value was highly significantly lower than that of the normal group ($P < 0.001$).

The mean testosterone binding affinity of the normal men did not differ significantly from that of the oligospermic men ($P > 0.6$), the impotent men ($P > 0.3$), or the azospermic men ($P > 0.1$). The hypogonadal group had a higher mean binding affinity in comparison with that of the normal men but this was due to four of the 16 subjects aged 15 to 19 with delayed puberty whose values were similar to those of our normal female subjects. These have nearly double the binding capacity of adult males and the same has been found in prepubertal male children by August *et al.* (1969). The individual results are plotted in fig. 2 together with 14 values for normal female subjects.

Conclusion

The finding that there was no deficiency of plasma testosterone in a group of impotent but otherwise normal men helps to explain why androgen therapy is usually unsuccessful in the treatment of impotence. Three of the present cases had previously been treated with testosterone oenanthate by injection or fluoxymesterone by mouth and had not shown improvement. Intramuscular injections of testosterone oenanthate 250 mg can markedly raise plasma testosterone—for example, up to 2,000 ng/100 ml testosterone seven days after injection—yet in spite of these high levels impotent males generally obtain little or no benefit—in striking contrast to hypogonadal males.

We have not attempted to explain the discrepancy between our findings of normal plasma testosterone and the low urinary testosterone reported by other workers. Ansari *et al.* (1973) reported reduced urinary testosterone excretion in impotent patients but, like us, they found normal plasma testosterone levels. They also reported a raised luteinizing hormone concentration and a reduced oestrogen excretion. One of our patients (case 27) aged 62 has been treated with oestrogen and showed some response. Teter (1972) claimed that mixed oestrogen and androgen therapy gives better results in selected cases of the so-called male climacteric with impotence than testosterone alone.

Our data on plasma testosterone in oligospermia and azospermia confirm the findings of Rosen and Weintraub (1971). We also find that the testosterone binding affinities in these groups are normal.

It is noteworthy that the older but not the younger hypogonadal males have normal adult male testosterone binding affinities in spite of their low plasma testosterone levels. Why adult men, whether normal or hypogonadal, have lower testosterone binding affinities than women and prepubertal boys remains obscure.

Thanks are due to Mr. V. Asta for preparing the figures. The Wellcome Trust are thanked for a generous grant for the purchase of the Packard Tri-Carb model 3375 liquid scintillation spectrometer.

References

- Ansari, M. A., Glover, T. D., Racey, P. A., and Rowe, P. H. (1974). *Journal of Endocrinology*, In press.
- August, G. P., Tkachuk, M., and Grumbach, M. M. (1969). *Journal of Clinical Endocrinology and Metabolism*, 29, 891.
- Cooper, A. J., Ismail, A. A. A., Smith, C. G., and Loraine, J. A. (1970). *British Medical Journal*, 3, 17.
- Ismail, A. A. A., *et al.* (1970). In *Reproductive Endocrinology*, ed. W. J. Irvine, p. 138. Edinburgh, Livingstone.
- Lawrence, D. M., and Swyer, G. I. (1973). *Annals of Clinical Biochemistry*, 10, 115.
- Mayes, D., and Nugent, C. A. (1968). *Journal of Clinical Endocrinology and Metabolism*, 28, 1169.
- Murphy, B. E. P. (1970). In *Steroid Assay by Protein Binding*, ed. E. Dzafalusky, p. 37. Copenhagen, Bogtrykkeriet.
- Rosen, S. W., and Weintraub, B. D. (1971). *Journal of Clinical Endocrinology and Metabolism*, 32, 410.
- Teter, J. (1972). *British Medical Journal*, 4, 114.