CORRESPONDENCE

Correspondents are asked to be brief

Outmoded Barbiturates

G. F. B. Birdwood, M.B., and Mollie Craven...148

Polio Vaccination for Travellers

J. E. Banatvala, M.D., and G. T. Spencer,
F.F.A.R.C.S.148

Dangers of High-solute Infant Feeding

B. W. Lewis, M.R.C.P.148

The Disc Sensitivity Test

J. C. Sherris, M.D., and others149

Quality Evaluation in Histopathology

D. A. Owen, M.B., and J. R. Tighe, F.R.C.P.149

Disclosure of Medical Records

J. Patterson, M.B., and others150

Retinitis Pigmentosa

A. C. Bird, F.R.C.S., and B. S. Jay, F.R.C.S.150

Diagnostic Test for Multiple Sclerosis

M. A. Crawford, Ph.D., and A. G. Hassam,
B.Sc.150

Duration of Action of Practolol

V. D. U. Hunt, M.D., and others151

Drug Combinations for Anaesthesia

D. W. Bethune, F.F.A.R.C.S., and others;
A. D. Forman, L.D.S.R.C.S.151

Epanutin and Isoniazid Interaction

J. Johnson, F.R.C.P.PSYCH.152

Oats and Coeliac Disease

G. D. N. Milne, M.R.C.G.P.152

Cardiovascular Disease and Peptic Ulcer

J. J. Segall, M.R.C.P.152

Thrombolytic Therapy in

Haemolytic-uraemic Syndrome

J. Stuart, F.R.C.P.ED., and R. H. R. White,
F.R.C.P.152

Strike Action in Hospitals

E. H. J. Smyth, F.R.C.S.ED.153

Interim Pay Review

M. S. Swani, M.B.; J. R. H. Heap, M.B.;
A. F. Crick, M.B.; A. Madwar, L.R.C.P.ED.,
and others; R. W. Billington, M.B., and others 153

Chairmanship of B.M.A. Committees

G. Scott, M.D.154

Increase in N.I. Payments for

Self-employed

J. H. Austin, D.O.M.S.154

Consultant Contract

J. Barron, M.R.C.PSYCH., and others; R. S.
Francis, F.R.C.P., and others; C. D. Lacey,
M.R.C.P.; A. W. B. Edmunds, F.R.C.P.;
C. S. Ogg, M.D.; S. R. Brennan, M.R.C.P.;
R. Brown, M.D.; J. S. Comaish, F.R.C.P.;
D. Adler, M.R.C.S.; R. W. Griffiths, F.R.C.S.;
J. H. M. Walker, F.R.C.S.; P. J. O'Sullivan,
M.B.; M. Ruben, F.R.C.S.154

Outmoded Barbiturates

SIR,—Your leading article (7 December, p. 552) was timely and helpful. Our aim is to prevent the opportunities for drug abuse as well as to supply health education in this field, so we are concerned that all doctors in general practice should fully appreciate the points you made. Firstly, the work in Ipswich referred to by Dr. Frank Wells, showing that the prescribing of barbiturates can be reduced by 65%, could be a prototype for all the medical profession. But only a few doctors prescribing less carefully can undo a great deal of good work. When the limitation of prescribing for amphetamines was accepted by most general practitioners it did not take drug-abusing groups of young people long to find doctors willing to prescribe methylphenidate, and now Drinamyl is around in fair quantities because some doctors have been "conned" into prescribing it.

Almost any drug with psychotropic qualities can be abused. Hence if an applicant unknown to the doctor applies—usually with elaborate excuses—for a supply of drugs which could possibly be used for the alteration of mood or as a drug of abuse, caution is essential. One certain way of limiting abuse is for the doctor to prescribe a maximum of five tablets, carefully writing in full in block letters to avoid a skilful alteration. This would aid a genuine patient. In the more likely case of the prescription being destined for the black market it would be almost useless and might never reach the chemist's counter. That particular doctor would not be troubled again by the wide circle of the applicant's drug-abusing friends. The consulting-room of a doctor who prescribes drugs on request is often filled in only a matter of hours by a queue of young hopefuls, and there have been cases of elderly doctors being intimidated by aggressive and threatening "temporary patients" demanding prescriptions. Where

prescribing can safely be limited we appeal to doctors to find the time and enthusiasm to give the patient confidence in a weaning process and never to prescribe more than minimal quantities of any drugs which can be abused. Many youngsters start their drug experiments from that over-stocked family medicine cupboard.—I am, etc.,

GEORGE BIRDWOOD

Chairman,

MOLLIE CRAVEN

Honorary Secretary,

Association for the Prevention of Addiction

London W.C.2

Polio Vaccination for Travellers

SIR,—In his paper on imported infections Dr. L. Roodyn (14 December, p. 648) states that it is still prudent to give a reinforcing dose of oral polio vaccine to children before departing to countries in which poliovirus is still endemic. However, we feel it particularly important to ensure that oral polio vaccine is offered not only to children, but also to adults if they have not previously had a full course of vaccine, if details of their polio vaccination status are in any doubt, or if they received a course of inactivated vaccine many years ago, for inactivated vaccines may not always induce persistent immunity to all three serotypes of poliovirus (personal communication from the Public Health Laboratory Service).

During the past four years we have seen five adults with poliomyelitis, one of whom died, while the others experienced severe and disabling disease resulting in permanent confinement to a wheel-chair. All contracted infection while abroad on holiday or business and none had ever been immunized against poliomyelitis. In addition, general practitioners must remember that an increasing

number of persons now travel on "package" holidays to non-tropical countries where poliovirus may still be prevalent—for example, parts of North Africa, Turkey, and Spain. We also feel that attention should be drawn to the need for immunization against poliomyelitis in brochures issued by travel agents, which at present is generally not done. Should there be insufficient time to provide a full course of polio vaccine, we recommend that travellers be given at least a single dose of oral vaccine before departure which, though not intended to provide adequate immunity, might prevent colonization of the gut by virulent poliovirus. The full course can then be completed either abroad if the visit is prolonged or on return.—We are, etc.,

J. E. BANATVALA

G. T. SPENCER

Department of Virology,

Department of Anaesthetics,
St. Thomas's Hospital,
London S.E.1

Dangers of High-solute Infant Feeding

SIR,—The Department of Health and Social Security has recently published a very welcome report¹ which lends support to our continued efforts to encourage breast-feeding of the newborn. It was further suggested that where breast-feeding was not possible a "low-solute" artificial feed should be used. Some excellent breast-milk substitutes are available but are relatively expensive—up to 70p for an average of five days' feeding. One might further hope that such expense would help to persuade the impecunious, at least, to breast-feed. This would be too naive a supposition however.

I have recently cared for an infant who has suffered brain damage as a direct consequence of the high cost of artificial feeding. An eight-week-old infant was admitted with severe hypernatraemic dehydration and convulsions. The precipitating cause was

pyrexia for 24 hours with reluctance to feed and the passage of one loose stool. During the previous three weeks he had been fed undiluted, unmodified cows' milk ("milkman's milk") with an added protein-rich cereal. He was desperately ill but recovered, only to be left with evidence of severe brain damage. It seems likely that this high-solute feeding regimen was the major factor in the production of his hypernatraemia. His mother's explanation for such an unusual feeding regimen in a young infant was that she could not afford the prepared dried milk powders.

I believe that a major campaign is required to alert mothers to dangerous feeding practices and this could be part of a national drive to encourage breast-feeding. There may also be a case for further subsidies on low-solute milk powders when breast-feeding is impracticable.—I am, etc.,

BARRY LEWIS

Whipps Cross Hospital,
London E.11

¹ Department of Health and Social Security, *Present-day Practice in Infant Feeding*. London, H.M.S.O., 1974.

The Disc Sensitivity Test

SIR,—Your leading article on this subject (13 July, p. 74) has come to our attention. It contained a number of misapprehensions about the Food and Drug Administration (F.D.A.) and International Collaborative Study (I.C.S.) recommendations for diffusion susceptibility testing. The most important is the statement that these procedures are "uncontrolled." Specific recommendations for the use of two recognized control strains and the statistical limits of acceptable results with them have been published in several places.¹⁻⁴ The question of quality control has also been broadly reviewed.^{1,6} Medium control at the manufacturer level is feasible to resolve finally the problem with *Pseudomonas aeruginosa* and the gentamicin group of antibiotics and hopefully will soon come about.⁷ With this one exception, inter-manufacturer and batch-to-batch reproducibility of Mueller-Hinton medium produced by the two major U.S. manufacturers is now very good.

The basic question raised by your article, however, is whether the aim should be towards the performance of susceptibility tests by standardized methods and with standardized reagents designed to give reproducibility in absolute terms or whether we should be satisfied with the results relative to those of control strains without requiring standardization of media or test conditions. We favour the first approach for several reasons. It permits relating diffusion to quantitative dilution test methods and allows the collection of cumulative statistical information as to population distributions. These can then be used in determining breakpoints for qualitative reports with both old and new antimicrobics. A statistical approach to control can be used which takes account of inherent technical variation in the results of individual tests with control strains but which will also detect variation in control strains because all other components have been standardized.

In the method recommended in your article the experimental error for individual

tests may be the sum of the errors with both the test and control strains and will be enhanced if objective methods of inoculum standardization are not used. The procedure uses preset criteria of "sensitivity" which have not been described in quantitative terms and which are the same for most antimicrobics irrespective of their diffusion rates. Results with all organisms and antimicrobics appear to be related to single strains of *Staphylococcus aureus*, *Escherichia coli*, and *P. aeruginosa* and variation or contamination could thus go undetected and lead to serious errors. The procedure in its manual form is also restrictive of the number of agents that can be tested on one plate.

Three other points in your article merit brief comment. A 100- μ g carbenicillin disc has now been approved in the United States as a result of studies such as that which was cited in the article. Neither cefoxitin nor cefamandole has so far been released for use in the United States and thus the issue of separate discs for these agents has not yet arisen here. As to a separate minocycline disc, evidence to date is insufficient to recommend this agent for the treatment of tetracycline-resistant staphylococcal infections, and diffusion tests are inapplicable to slow-growing organisms such as *Nocardia asteroides*. However, we know of no fixed positions as to restriction of discs and new congeners which have significantly different spectra of activity and would expect their licensure by the F.D.A. as and when the need arises and is supported by the necessary scientific data and the recommendations of their expert advisory committees.

We agree with you on the desirability of further studies to determine current inter-laboratory reproducibility with these techniques, and several involving the F.D.A. diffusion test are under way or are in the process of being reported. It is, however, equally important to be concerned about what can be achieved and we believe that results of both dilution and diffusion tests can be significantly improved only by the acceptance and use of standardized methodology at the very least for reference procedures.—We are, etc.,

JOHN SHERRIS

Department of Microbiology,
University of Washington,
Seattle, Washington

CLYDE THORNSBERRY

Antimicrobics Investigation Section,
Clinical Bacteriology Branch,
Bacteriology Division,
Center for Disease Control,
Atlanta, Georgia

ARTHUR L. BARRY

Chairman,
Subcommittee on Antimicrobial Susceptibility Tests,
National Committee for Clinical Laboratory Standards
Microbiology Laboratories,
U.C. Davis-Sacramento Medical Center,
Sacramento, California

- 1 Federal Register, 1972. Rules and regulations. Title 37: 20525-20529.
- 2 Federal Register, 1973. Rules and regulations. Title 38: 2576.
- 3 *Current Technique for Antibiotic Susceptibility Testing*, ed. A. Balows. Springfield, Thomas, 1974.
- 4 *Manual of Clinical Microbiology*, ed. E. H. Lennette, E. H. Spaulding, and J. P. Truant, 2nd edn. Washington, American Society for Microbiology, 1974.
- 5 Schoenkecht, F. D., and Sherris, J. C., in *Recent Advances in Clinical Pathology*, ed. S. C. Dyke, ser. VI, p. 275. Edinburgh, Churchill Livingstone, 1973.
- 6 Blazevic, D. J., Koepcke, M. H., and Matsen, J. M., *American Journal of Clinical Pathology*, 1972, 57, 592.
- 7 Reller, L. B., et al., *Journal of Infectious Diseases*. In press.

** Our correspondents are themselves under at least one misapprehension. Naturally if the type of test advocated in our leading article were to be commended by any authority for general use the test conditions (including inoculum standardization, now possible by a simple method not involving turbidimetry¹), and indeed its every feature except the composition of the culture medium would require definition. If a small fraction of the effort which has gone into the evolution of the F.D.A. and particularly the I.C.S. method were devoted to this task it should not present difficulty. Indeed, a description of one such method adequate for its performance without risk of any serious error already exists.² It will be little consolation to workers in Britain and many other countries if two manufacturers in the United States at last manage to control the composition of Mueller-Hinton medium sufficiently. Such control is most unlikely to operate world-wide.—ED., B.M.J.

¹ Felmingham, D., and Stokes, E. J., *Medical Laboratory Technology*, 1972, 29, 198.

² Stokes, E. J., and Waterworth, P. M., *Association of Clinical Pathologists, Broadsheet 55*. London A.C.P., 1966 (Rev. 1972).

Quality Evaluation in Histopathology

SIR,—Opinions have recently been expressed about the desirability of a quality control system in histopathology. Many factors affect the quality of histopathological opinion, not the least of these being the quality of the surgical specimen, the technical processing of the tissue, and clinicopathological consultation. However, we are concerned particularly with quality evaluation of the pathologist's opinion and consistency of diagnosis by a group of pathologists.

An ideal system of quality evaluation of histopathological opinion should reproduce the normal laboratory situation. The design of a system is complicated by (1) the difficulty of obtaining an acknowledged correct diagnosis based on opinion, (2) the need for clinical and other pathological data in reaching a diagnosis, and (3) variation in terminology. The Royal College of Pathologists of Australia base their system on sets of sections illustrating uncommon conditions, the diagnosis being "established" by acknowledged experts. Specialist histopathologists obtain an average of only 75% "correct" answers.¹ The histopathology proficiency testing programme of the American College of Pathologists uses randomly selected sections from five types of tissue. Consensus among the participating pathologists is reached when 50% or more of the participants make the same diagnosis. Consensus was reached in only 77% of cases.²

Both of these surveys suggest a disturbing disparity in histopathological opinion. For this reason a quality evaluation system has been instituted in this department. Each week 10 random sections from 1970 are used. The participating pathologist is informed of the age and sex of the patient and the site of origin (where known) of the tissue. Diagnoses are collected before group discussion, at which time full clinical information, including follow-up details, are available. Consensus is reached when all the participating pathologists agree on a final diagnosis after review of the clinical and pathological information. A final consensus diagnosis has