people. Thus for them the "open door" often becomes a "revolving door."^{2,3} On the other hand, they sometimes adjust surprisingly well to a therapeutic community approach in a prison unit.^{1,4} Under certain conditions, as Dr. Scott points out, prisons have therapeutic possibilities and prison officers can find such rehabilitative work stimulating and emotionally rewarding.⁴ Clearly, mentally abnormal offenders require treatment and not punishment but account must be taken of their ability to co-operate when deciding where they should be treated.

Obviously a range of community-based alternatives to prison are urgently needed, but it is unrealistic to expect that every abnormal offender will be suitable for treatment in community-based or in "open" hospital units or that within the foreseeable future enough community-based facilities will be available. Thus at present many such offenders fall between two stools. They are frequently deemed unsuitable for, or unacceptable to, "open" permissive hospital units and no adequate provision for their treatment is provided in prison. The establishment of therapeutic communities in prison should not be allowed to lead to the need for community-based rehabilitation facilities being ignored or to suitable offenders being refused admission to hospital units. The services run by the Home Office and the Department of Health and Social Security should be closely integrated—as suggested by Dr. Scott—with interested doctors and nursing staff working in both hospitals and prison. Thus, a co-ordinated long-term rehabilitation effort could often begin in a prison and be continued via a hospital unit to community services. Neither hospital nor prison treatment have a chance of lasting success without prolonged support in the community after discharge. But it does not help every offender if the slogan "these people need help not imprisonment" prevents the potential of prison therapeutic communities being used. A few psychopathic patients may be "carried" by a non-psychopathic majority, but to admit all of them into, for example, alcoholic units would completely alter the therapeutic atmosphere and the (voluntary) "group-code" and would greatly affect the morale of other patients.—I am, etc.,

M. M. GLATT

Alcoholism and Drug Dependence Unit,
St. Bernard's Hospital,
Southall, Middlesex

- 1 Glatt, M. M., *Drugs, Society and Man. A Guide to Addiction and its Treatment*. Lancaster, M.T.P., 1974.
- 2 Rollin, H. R., *The Mentally Abnormal Offender and the Law*. Oxford, Pergamon, 1969.
- 3 Shepherd, E. W., *British Medical Journal*, 1974, 2, 716.
- 4 Glatt, M. M., *British Medical Journal*, 1974, 1, 56.

Variations in Carboxyhaemoglobin Levels in Smokers

SIR,—It would be valuable to be able to identify those individuals who are at high risk of developing diseases associated with smoking. Present evidence suggests that a random measurement of the blood carboxyhaemoglobin (COHb) level may be helpful in this respect,¹ but since an individual's COHb level can fluctuate widely over 24 hours the use of a random COHb measure-

TABLE II—COHb Levels in Four Cigarette Smokers on Several Days

Subject	No. of Cigarettes Smoked/Day	No. of Test Days	Mean COHb% (range)		
			10.00 hr	15.00 hr	22.00 hr
7	25	6	4.8 (3.8-6.0)	6.8 (5.0-9.1)	8.9 (7.8-10.0)
8	22	5	3.9 (2.9-5.6)	3.9 (3.7-6.9)	6.0 (3.8-7.0)
9	26	5	5.9 (5.3-6.9)	6.8 (4.6-9.2)	8.6 (5.9-13.9)
10	20	6	3.0 (2.3-4.0)	5.3 (4.8-6.0)	6.5 (4.7-8.3)

ment must be examined critically. In a recent article (28 December, p. 736) Drs. C. M. Castleden and P. V. Cole concluded that "a random COHb estimation gives a good indication of the mean COHb of an individual," but since they do not define what is meant by "the mean COHb" and do not explain their grounds for judging how well an estimation indicated the mean, their conclusion cannot be evaluated. They claim that "in the individual smoker the COHb level does not increase gradually during the day but is kept within relatively narrow limits." However, their data do not support this. Some of the 10 subjects studied had COHb levels which did tend to rise during the day (for example, subjects 1, 3, 7, and 9) and in only one was there a tendency to fall.

We have collected similar data as part of a larger study, and in our experience COHb levels rise during the day, usually reaching a maximum in the evening. Table I shows

TABLE I—Mean Morning and Afternoon COHb Levels in Six Cigarette Smokers

Subject	No. of Test Days		Mean COHb%		Ratio of p.m. to a.m. COHb Levels
	a.m.	p.m.	a.m.	p.m.	
1	3	5	6.8	14.7	2.2
2	3	6	5.1	8.6	1.7
3	3	5	2.2	5.2	2.4
4	3	4	2.9	6.9	2.4
5	3	8	2.2	4.4	2.0
6	3	4	3.9	5.0	1.3

the results of one series of tests in which blood samples were taken from six smokers in the morning (about 10.00) and in the afternoon (about 16.00) on several occasions. In each subject the mean afternoon COHb level was higher than the mean morning level—twice as high on average—and the differences were statistically significant ($P < 0.05$). Table II shows the results of a second series in which four subjects were each tested at 10.00, 15.00, and 22.00 on at least five days, and in each subject the average of the COHb readings increased during each interval ($P < 0.001$). In neither of these experiments were the subjects asked to alter their smoking habits in any way, and there were no restrictions on the timing of the tests in relation to recent cigarette smoking. It therefore seems likely that a random COHb level corrected for the time of day provides a better indication of the mean daily level than an uncorrected random value. Indeed, a correction for recent cigarette consumption might give an even better estimate of the mean level and we have recently described a method of using a random COHb level in this way.² Though such a mean COHb level is likely to be a more powerful predictor of smoking-related diseases than an uncorrected random value, this can be confirmed only in a prospective study.—We are etc.,

N. WALD
SUSANNAH HOWARD

D.H.S.S. Cancer Epidemiology and Clinical Trials Unit,
Department of the Regius Professor of Medicine,
Radcliffe Infirmary,
Oxford

- 1 Wald, N., et al., *British Medical Journal*, 1973, 1, 761.
- 2 Wald, N., et al., *Thorax*. In press.

Discouraging Smoking

SIR,—Your leading article on health education (1 February, p. 233) emphasizes the failure to educate the public about cigarette-induced diseases, and the lack of response over the past 25 years.

Successive governments have raised the tobacco tax for the past 60 years, but this tax is now smaller owing to the present inflation. Also the cost of treating these diseases has increased out of proportion, with very little success. It is time that a sensible and realistic tax be imposed so as to reduce the amount of cigarettes bought and consumed. This revenue could help and pay for the increasing cost of the enlarging N.H.S. reorganization.—I am, etc.,

A. G. I. STOCKLEY

Thornton Heath, Surrey

SIR,—Your leading article (1 February, p. 233) on "Health Education in the Re-organized N.H.S." was most timely and, as it happens, coincided with the publication of our booklet *So Now You Know About Smoking* by Dr. Alfred Yarrow. Dr. Yarrow is in fact a senior medical officer concerned with the public health at the Department of Health and Social Security, so we regard it as something of a triumph to have persuaded him to write such an excellent booklet for us on this subject. Perhaps where we lead others may follow.—I am, etc.,

EVELYN BROWN

Executive Editor,
Family Doctor Publications

London N.W.1

Women Doctors in the N.H.S.

SIR,—According to your leading article, "State of Health" (28 December, p. 732), women form 32% of entrants to medical schools in England and Wales yet they constitute only 11.8% of principals in general practice and 7.9% of consultants.

I work six sessions as a consultant child psychiatrist and two as a clinical assistant in a teaching hospital. My specialty is very short-staffed. I do no private practice. I spend 53 hours a week in professional activities (compared with the full-time average of 60 hours). I do some of the activities conventionally left to mothers—household shopping, cooking, etc., and bringing up children. I do these willingly, but I make the point that male colleagues generally

do not do them. Some women would also be expected to entertain their husband's business colleagues.

My annual expenditure on costs incurred only because of my professional work amounts to £1980, made up as follows: car (£500 less mileage allowance), £300; domestic help (five hours daily at £1 an hour), £1250; extra help in school holidays (student help at £8 a week), £80; extra clothes (jeans and jumpers are unsuitable, even today, for consultants), £150; subscriptions to professional associations and journals and cost of books and conferences, £200.

Except for the B.M.A., most professional associations do not have earnings-related subscription rates at consultant level. Conference fees are paid pro rata to sessions worked, leaving a substantial sum to come from my pocket. My list of items of expenditure takes no account of hidden extras—for example, more easily prepared and therefore more expensive food; inability to shop round for bargains; no time to make clothes; hire of car or taxi when car out of action; and so on. My total annual take-home pay for 53 hours professional work a week and eventually a pension is £3600, leaving a net income of £1620.

If one-third or so of British-trained women doctors have to face these expenses in order to pursue their chosen career is it surprising that they drop out? This problem has not been adequately considered publicly. Of all hospital registrars 56% were born overseas. As you say, Sir, "international trends may cut off the supply." Can a solution be found to the problems of women doctors that would enable a large work force, eager to use their skills provided they are adequately paid, to return to work?—I am, etc.,

DORA BLACK

Fainting and Anaesthesia

SIR,—When a simple uncomplicated local dental anaesthetic is followed by "fainting" hypoxaemia may be the cause. This is due to the nervous and apprehensive patient involuntarily overbreathing in the interval before treatment and this being increased because of the sights and sensations related to the actual injection. When this type of faint occurs the lowering of the ionic calcium fraction consequent upon the hypoxaemia causes a tetanic contraction, particularly of the hands, followed by generalized clonic contractions. On recovery the patient complains of muscle pains, again particularly in the arms.

The use of a general anaesthetic for such a dental patient may be indicated because of the nervousness and apprehension. Thus even in the absence of an overt faint the respiratory drive of carbon dioxide may be greatly reduced and to this is added the respiratory depression of the particular drug used.

An effective premedication, as is given to the general hospital population prior to surgery, prevents overbreathing and its omission may be the reason why a general anaesthetic, as currently given for dental procedures, poses unique difficulties with, rarely, tragic results.—I am, etc.,

BARRY HEMPHILL

London W.1

Unexplained Fever

SIR,—In his article on unexplained fever (16 November, p. 397) Dr. A. M. Geddes states: "Benign tertian malaria can usually be treated successfully by a standard course of chloroquine. . . ." However, it should be noted that most patients with a *Plasmodium vivax* infection will relapse if treated with chloroquine alone, owing to the maturation of secondary exoerythrocytic schizonts in the liver and the subsequent invasion of erythrocytes. The 4-aminoquinoline group of drugs such as chloroquine acts only on the blood stages of *P. vivax*, having a potent schizonticidal effect on the asexual parasites and a lesser effect on gametocytes.

Eradication of tissue schizonts and the prevention of relapse are best achieved by a course of an 8-aminoquinoline drug such as primaquine. Usually, the standard course of primaquine (7.5 mg base twice daily for 14 days in an adult) will deal effectively with most strains of *P. vivax* imported into Britain, but people infected in some south-east Asian and western Pacific countries may require 7.5 mg primaquine base three or four times a day to clear all tissue schizonts. Since April 1973 11 such patients out of 367 with fully documented histories of treatment have come to the attention of the Malaria Reference Laboratory. These remarks on the need for primaquine treatment of *P. vivax* tissue stages apply equally to infections with *P. malariae* and *P. ovale*.

Dr. Geddes advises: "If there is any possibility of chloroquine resistance the patients must be treated with quinine." This needs qualification. Certain south-east Asian and central American strains of chloroquine-resistant *P. falciparum* have been shown to exhibit grade R-1 resistance to quinine, and we feel that such patients should be treated in hospital by a physician experienced in tropical diseases who has the facilities available for regular monitoring of parasitaemia together with experience of the other agents available for the treatment of multidrug-resistant *P. falciparum* infections. Further, there have been several proved cases of transplacental transmission of malaria in the U.K., and febrile illnesses in young children who have never left Britain may require careful history-taking of the mother's movements as well as blood-film examinations on mother and child.

It is also necessary to emphasize that, contrary to the implication of the last paragraph on malaria in Dr. L. Roodyn's article on immunization and other prophylactic measures against imported infections (14 December, p. 648), chloroquine resistance is not yet known from Africa.—We are, etc.,

B. A. SOUTHGATE
W. H. R. LUMSDEN
D. J. BRADLEY
D. M. MACKAY

Ross Institute of Tropical Hygiene and
Department of Medical Protozoology,
London School of Hygiene and Tropical Medicine,
London W.C.1

Erythema Infectiosum

SIR,—The B.M.J. of 23 November contained a leading article called "Fourth, Fifth, and Sixth" (p. 429) and also a letter from Dr. C. A. H. Watts (p. 466) reporting two cases of fifth disease, or erythema in-

fectiosum, occurring in close friends. I was interested at the time since, because I qualified fairly recently (1965) and despite having worked for three years in a fever hospital, I had never heard of the condition at all.

I am writing now because it may be of interest that over the past few weeks I have observed the same clinical picture, exactly as described in your leading article, in my own three children, aged 9, 6, and 2 years. They have had only minor constitutional upset but a brilliant red rash confined to their cheeks and a duller rash, variable and pleomorphic, on their limbs and trunks, lasting on and off for just over one week. Moreover, since being alerted to the condition I have so far seen at least three other children in High Wycombe with an identical illness, so presumably we are in the middle of a small local epidemic of this supposedly rare disease. I have so far been unable to find whether any definite viral cause for the illness has been established, or whether an increased incidence is being noted elsewhere in the country.—I am, etc.,

A. W. J. MEDHURST

High Wycombe, Bucks

Alpha-adrenoceptor-blocking Drugs in Asthma

SIR,—We have read with interest the paper by Professor S. Bianco and others on the prevention of exercise-induced asthma by indoramin (5 October, p. 18) and the letter from Dr. K. N. V. Palmer and others (16 November, p. 409) on the increased effect of salbutamol after intravenous administration of indoramin and thymoxamine.

We have drawn similar conclusions about the role of alpha-receptors from our studies with guinea-pig trachea and human bronchus in vitro. The tracheal chains of normal and egg-protein-sensitized guinea-pigs or spirals of human bronchus obtained during pulmonary operations were used in Krebs' bathing fluid perfused with carbogen. Several combinations of alpha-blockers and beta-mimetics were studied. The alpha-blockers dihydroergocristine, Hydergine (a combination of dihydroergotoxines), phenoxylbenzamine, phentolamine, and thymoxamine and the beta-mimetics salbutamol, terbutaline, isoprenaline, and ephedrine were used.

Small concentrations (10 $\mu\text{mol/l}$) of each of the alpha-blockers potentiated (to about double) the effects of beta-mimetics, though these doses of alpha-blockers alone did not show any effect. Concentrations over 10 $\mu\text{mol/l}$ were less effective. Egg-protein sensitization and the use of egg protein in the bathing fluid did not significantly change the potentiation. The degree of potentiation in human bronchus preparations was of the same order as in guinea-pig tracheal preparations. On the other hand, when alpha-blockers were combined with the same beta-mimetics in preparations of rat atria no potentiation of the beta-mimetics was observed.

Though the tracheobronchial alpha-receptors seem to be almost "silent," they seem to decrease the therapeutic effects of beta-mimetics to a considerable degree. The fact that the same is not true of the heart is additionally favourable for the combination of an alpha-blocker with a beta-