(4) That while we may regard health education as wholly desirable if it reduces true poisoning in children, this education in its present forms of commercial advertising and television programmes may also have the effect of generating poisoning scares. Parents are naturally anxious about the health of their children and these health education techniques may have heightened this anxiety. The consequence of this may be that some children have to suffer unnecessary clinical procedures in the hospital. Health education methods must therefore be examined in terms of their costs and benefits.

We are all agreed that suspected poisoning is not a random event but occurs under certain special social circumstances. We believe it to be useful to examine these circumstances and to recognize that those which produce a poisoning scare may be different from the circumstances which lead to a case of true poisoning. We expect then to be in a better position to reduce the latter without inflating the former. Policy until now has meant that parents have had to take full responsibility for the prevention of suspected poisoning in children and have had to suffer the consequences. However, evidence in the above correspondence suggests that in practice parents may have limited control over the events which lead to suspected poisoning. The responsibility for prevention must be shared and parents should be provided with preventive measures which have been shown to be effective through experiment and field studies.—We are, etc.,

M. W. CALNAN

National Children's Bureau,
London E.C.1

J. W. DALE

S.E. Thames Regional Health Authority,
Croydon, Surrey

¹ Calnan, M. W., *Community Health*, 1974, 6, 91.

SIR,—Two points raised by Drs. A. W. Craft and R. H. Jackson in their letter (11 January, p. 95) about the research by Calnan¹ call for comment. I set up the research project.

By considering all the evidence it was possible to distinguish clearly between those children who developed symptoms and/or ill effects and those who did not. Those who had a quantity of a toxic substance retrieved from the stomach were classified as poisoned, even though no symptoms developed. It was found that all children admitted, irrespective of symptoms or ingestion, were recorded by the hospital as "poisoned" whereas children not admitted never reach the national statistics. Therefore nationally there appeared to be a gross overestimate of those actually poisoned and a considerable underestimate of poisoning scares. I do not think that distinguishing between true poisonings and poisoning scares will lead to an increase in the number of children dying; it can only clarify the situation and help more rational management.

The second point I wish to make is that no one, however experienced or otherwise, could hope to predict the onset of symptoms unless he first seeks evidence of possible ingestion. The following recent personal clinical experience illustrates my point. It involved an 18-month-old child found by the mother with an open bottle of

aspirin and a few fragments in the mouth. Since the number of missing tablets was unknown, the simplest decision would have been to wash its stomach out and admit it, to be eventually classified as "poisoned." However, on close questioning of the mother it became quite evident that she had not left the child alone in the room with the bottle for more than one minute, which could have given the child enough time to open the bottle and put one tablet at the most in its mouth. I then reassured the mother, who had herself by then realized the possibilities with obvious relief.

The research demonstrated that the appropriate questions were not being asked and therefore admission appeared arbitrary, possibly reflecting clinical judgement without facts. Assessment of risks in management is accepted clinical practice. Only appropriate questions can determine the facts on which to decide whether a child with a head injury needs admission. Many children with such injury are sent home because intracranial complications are thought highly unlikely.

I find that even very anxious mothers can be adequately questioned when first seen and provide very reliable information. The demonstration of this to young doctors is very easy and they quickly grasp what it is all about. But it requires experienced people directly involved in the primary-care front line to lay down the guidelines.—I am, etc.,

C. P. DE FONSEKA

Accident and Emergency Department,
Royal United Hospital,
Bath

¹ Calnan, M. W., *Community Health*, 1974, 6, 91.

Small Bowel Tumours

SIR,—Your leading article (18 January, p. 115) contains an ambiguous statement which is confusing. In referring to complications of adult coeliac disease you say: "Both lymphoma and carcinoma occur, but the incidence of carcinoma is reduced if the patient is treated with a gluten-free diet; this is as yet unproved for lymphomas. These findings justify the continued use of gluten-free diet for the whole of the patient's life."

Such a statement in an article on small bowel tumours implies that the lymphoma and carcinoma are situated in the small bowel. Though it is now well established that lymphomas of the small bowel may complicate adult coeliac disease,¹ the true relation of small intestinal carcinoma to adult coeliac disease is not known. Indeed, the same series of adult coeliac disease which established its association with abdominal lymphoma¹ contained no carcinomas of the small bowel. The carcinomas referred to were situated in the oesophagus and elsewhere in the gastrointestinal tract. You support your premise by reference to a paper by Holmes *et al.*,² which is from the same group and which presumably contains substantially the same cases. This paper refers vaguely to "all cancers" and to "carcinoma of the oesophagus and reticulosarcoma." No specific mention of small bowel carcinoma is made.

Some 16 cases of jejunal carcinoma associated with either steatorrhoea or histological villous abnormality have been described in

the literature.³⁻⁶ In all but one of these the mucosal histology was studied at the time of, or within months of, the carcinoma being resected. This makes them subject to the hypothesis that the mucosal abnormality is a result of the carcinoma. In the other case³ the diagnosis of adult coeliac disease was established seven years before resection of the carcinoma. I studied a similar (unpublished) case in a woman who had suffered from steatorrhoea in early childhood and was diagnosed as having adult coeliac disease with jejunal biopsy evidence of subtotal villous atrophy at the age of 56 years. She went on to develop a jejunal carcinoma five years later.

I think it is likely that jejunal carcinoma may be a complication of adult coeliac disease, but as yet the case is supported by circumstantial evidence only and not proved as your leading article implies.—I am, etc.,

N. R. PADLEY

Department of Pathology,
Poole General Hospital,
Poole, Dorset

¹ Harris, O. D., *et al.*, *American Journal of Medicine*, 1967, 42, 899.

² Holmes, G. K. T., *et al.*, *Gut*, 1974, 15, 339.

³ Kenwright, S., *Postgraduate Medical Journal*, 1972, 48, 673.

⁴ Asch, T., and Seaman, W. B., *Radiology*, 1971, 100, 271.

⁵ Blackwell, J. B., *Gut*, 1961, 2, 377.

⁶ Joske, R. A., *Gastroenterology*, 1960, 38, 810.

Metoclopramide and Breast-feeding

SIR,—Paediatricians in developing countries are deeply involved in the battle against the decline of breast-feeding. It has recently been reported¹ that metoclopramide is a potent stimulator of prolactin release. This could be important in the prevention of the premature weaning that has serious consequences in preindustrialized areas. I have therefore tried the effect of giving 10 mg of metoclopramide three times daily to five lactating mothers who had suffered a severe decrease in milk volume for more than 72 hours. In every case the results were most encouraging. Both mother and child were put under close observation during the 7-10 days of treatment because metoclopramide might produce side effects in the mother² and I do not know how much of the drug is excreted in breast milk. In these five cases no side effects were observed.

I am at present investigating the clinical use of the drug in larger trials and trying to determine the amount of metoclopramide excreted in breast milk.—I am, etc.,

PAULO L. R. SOUSA

Department of Paediatrics,
Universidade Católica de Pelotas,
Pelotas, Brazil

¹ McNeilly, A. S., *et al.*, *British Medical Journal*, 1974, 2, 729.

² Shaklay, M., Pinkhas, J., and de Vries, A., *British Medical Journal*, 1974, 2, 385.

Removal of Potassium from Fruit Juices by Ion Exchange

SIR,—Many attractive drinks are denied to patients with renal failure because of their high potassium content. The addition of resins to juices has been described,^{1,2} potassium being exchanged for sodium or hydrogen. But these have the disadvantage of increasing the sodium intake or lowering

TABLE I—Mean ($\pm$ S.E.) Concentrations of Potassium and Sodium in Fruit Juices before Passage through Column of Zerolith and Concentrations after Passage

	Potassium (mmol/l)		Sodium (mmol/l)	
	Apple Juice	Grape Juice	Apple Juice	Grape Juice
Before passage	28.2 $\pm$ 1.5	22.7 $\pm$ 3.0	1.0 $\pm$ 0.1	6.0 $\pm$ 0.5
After passage of:				
1 bottle (0.75 l)	0.0	0.0	0.8	0.5
2 bottles (1.5 l)	0.0	0.0	0.0	0.7
3 bottles (2.25 l)	0.8	0.8	0.5	1.0
4 bottles (3.0 l)	1.5	1.2	0.5	1.5
4.7 bottles (3.5 l)	3.0	2.9	0.6	1.5
5 bottles (3.75 l)	4.1	4.0	0.8	1.5

Conversion: SI to Traditional Units—Potassium: 1 mmol/l = 1 mEq/l. Sodium: 1 mmol/l = 1 mEq/l.

TABLE II—Concentrations of Potassium and Sodium in 5 ml of Fruit Juices before and after Hand-mixing with Zerolith

Zerolith in Mix (g)	Potassium (mmol/l)		Sodium (mmol/l)	
	Apple Juice	Grape Juice	Apple Juice	Grape Juice
0	25.1	16.8	1.0	10.0
1	7.2	4.1	0.5	5.0
2	3.5	2.4	0.0	3.0
3	2.7	2.0	0.0	3.0

pH. We have removed potassium from commercially available fruit juices by ion exchange chromatography in a column packed with a calcium exchanging resin.

Dry Zerolith 225 Ca-form resin 150 g was swollen in distilled water for one hour and then, after thorough washing, poured into a 60-cm $\times$ 3-cm column. The column was washed with 300 ml distilled water. Commercially available fruit juices were then passed through the column and the eluate collected. The potassium and sodium contents of the juices were determined by flame spectrophotometry before and after chromatography. The results were compared with those obtained by hand-mixing 1 g 2 g, or 3 g dry Zerolith with 5 ml of the same juices over a period of five minutes. With the Zerolith column the potassium content of up to 3 l (4 bottles) of apple or grape juice could be lowered to acceptable levels (table I). The column allowed a flow rate of 1 l per hour. After 3.5 l of juice had passed through the column the potassium concentration of the eluate reached an unacceptable level of 3.0 μ mol/l and the column had to be discarded or regenerated. Mixing the juices by hand with the resin gave less satisfactory results (table II).

The column procedure is unsuitable for turbid juices like orange or tomato juice, which tend to block the column. The calcium exchange resin does not lower pH or increase the sodium concentration in the juice, and this is preferable to other resins like Dowex 50W,¹ which exchanges potassium for hydrogen, and Resonium A, which exchanges potassium for sodium. The drinks treated by our method were judged highly palatable by the patients. However, we think it would be too risky to allow patients to make their own potassium-free fruit juices, as proposed by Levy *et al.*¹—We are, etc.,

G. J. VAN KAMP
W. BAKKER
J. G. M. C. ROSIER

University of Nijmegen,
Department of Medicine,
Division of Nephrology,
Sint Radboud Ziekenhuis,
Nijmegen, The Netherlands

¹ Levy, N., Boxer, J., and Carter, A., *New England Journal of Medicine*, 1973, 289, 753.

² Cacioppo, Ph.L., and Hollander, L., *New England Journal of Medicine*, 1972, 287, 361.

Blow-out Fracture

SIR,—Your excellent leading article (11 January, p. 50) drew attention to an important and easily overlooked bony injury. However, it is not an infrequent event to detect all the features of this injury, including inferior rectus trapping, the hanging-drop x-ray sign, and operative evidence of an orbital floor defect, without a history of blunt trauma to the orbital contents. In fact, many of these cases in our experience arise from an impact at the inferior orbital margin or to the body of the malar bone.

This discrepancy also stimulated Fujino¹ to test the conventional blow-out hypothesis using a weighted striker on the inferior orbital margin of dried skulls. He succeeded in producing by this method a characteristic "punched-out" fracture of the orbital floor and concluded that a direct buckling force transmitted from the orbital margin across the orbital floor in the absence of increased hydraulic pressure can produce a "blow-out" injury. This agrees with our own clinical experience.—We are, etc.,

MALCOLM HARRIS
GEOFFREY FORMAN
J. H. SOWRAY

Department of Oral and Maxillo-Facial Surgery,
King's College Hospital and Dental School,
London SE5

¹ Fujino, T. J., *Plastic and Reconstructive Surgery*, 1974, 54, 81.

SIR,—In these troubled times a leading article on "black eyes" (11 January, p. 50) seems appropriate. Unfortunately your clinical description of the condition is inaccurate and the management is confused.

Confining my remarks to pure blow-out fractures, with an intact orbital rim, then the presence of enophthalmos with retraction of the eye on attempted upward gaze is clinically diagnostic, whatever the x-ray findings may be. A more subtle test is the rise in intraocular pressure on trying to look upwards. These signs mean unequivocally that the orbital floor has been fractured and that herniated orbital contents have become trapped. Orbital emphysema following blunt trauma is diagnostic of a

blow-out fracture and usually implies ethmoidal involvement, in isolation or in association with a floor fracture. Of our 103 cases, 21% had ethmoidal fractures.¹

A patient presenting with a black eye and diplopia should have orthoptic assessment of the ocular movements which includes plotting the field of binocular fixation and the Hess chart. If the diplopia improves during the first few days repeated orthoptic testing will quantify this. Provided that the x-rays show only a small fracture and hernia and the enophthalmos is 2 mm or less, no orbital exploration is necessary since the prognosis for the disappearance of diplopia and the final cosmetic appearance is good.

The indications for surgical intervention¹ have been established as follows: (1) incarceration of orbital contents in the fracture with consequent globe retraction and increased intraocular pressure on attempted upgaze; (2) a fracture with large herniation of tissue into the antrum; (3) enophthalmos of 3 mm or more; and (4) diplopia not resolving significantly within the first 10 days. No decision about operation need be made before 10 days have elapsed. This allows time for bruising and oedema to settle and reveal the true situation concerning enophthalmos and diplopia. Surgical treatment of a pure blow-out fracture is not an emergency but should be confined to those cases fulfilling one or more of the criteria listed. Once the condition of blow-out fracture had been described in 1956² the fashion was for early surgical treatment, preferably within 24 hours. When many patients regained normal ocular movements after early surgery most surgeons were vain enough to believe that the improvement was because of the operation. A moment's reflection on the outcome for most patients with "black eyes" before 1956 will confirm the wisdom of the current approach.

It is clear that the ophthalmologist is the best person to deal with blow-out fractures not associated with generalized injuries, since the correct management of the diploma is paramount.—I am, etc.,

PETER FELS

Moorfields Eye Hospital,
London E.C.1

¹ Dulley, B., and Fells, P., *British Orthoptic Journal*, 1974, 31, 47.

² Smith, B., and Regan, W. F., Jun., *American Journal of Ophthalmology*, 1957, 44, 733.

S.L.E.-like Syndrome Associated with Mitochondrial Antibodies

SIR,—We were interested in the report of Dr. J. Guardia and others (15 February, p. 370) on four cases of a systemic lupus erythematosus-like syndrome associated with mitochondrial antibodies (A.M.A.) instead of the more usual antinuclear (A.N.A.) reactions found in the collagenoses. This syndrome was first identified by German authors, who described 27 cases in 1973.^{1,2} The illness is almost confined to the female sex and is characterized by recurrent attacks of fever, arthralgia, myalgia, pleural effusion with or without pericarditis, a high E.S.R., relative lymphopenia, normal liver function tests, and A.M.A. of high titre, mainly of IgG class. L.E. cells and A.N.A. are usually absent. No evidence of specific infections can be obtained and response to antibiotics is disappointing. Some patients require