possibility of updating it is also an added bonus. All forms of the media presentations (films, slides, and exhibitions) are expensive if produced commercially, but the N.H.S. already employs professional photographers and illustrators who, with encouragement and very modest expansion of their facilities, could provide most of its needs.—I am, etc.,

L. BOWCOCK

Department of Medical Illustration,
Suffordshire Area Health Authority,
North Staffs District

Inquiry among Rheumatism Unit Outpatients

SIR,—As our outpatients sat patiently waiting in the rheumatism unit a few months ago at Westminster Hospital we asked them to fill in two questionnaires asking them (1) for details of their daily commuting to work and the effect it had on their arthritis, and (2) what forms of unofficial therapy they had had for this condition while attending our clinic or their general practitioners. This action was taken as we felt that, in the London area, if a patient was fit to get there she was probably fit for her work, and that patients with arthropathies, where cure is rare and treatment largely symptomatic, would in most cases try some unorthodox form of therapy. The results of our inquiry might be of interest.

Firstly, of 65 patients who completed the form as regards daily commuting to work 38 had arthritis, 12 ankylosing spondylitis, six gout, and nine osteoarthritis. They gave very different stories. For the 16 men and 22 women with arthritis travelling time averaged 67 minutes a day. Much the most popular form of transport was by private automobile, the least popular were walking and travelling by bus. A common complaint was of being jostled and pushed aside by hurrying commuters when walking. This often caused as much emotional as physical upset. For the spondylitics the least popular forms of transport were essentially those which entailed prolonged immobility. With them, unlike the patients with arthritis, travel on foot or by car were equally popular. For osteoarthrotics the most unpleasant feature was standing waiting at bus stops. In the whole group travelling time varied from six to 140 minutes daily, averaging 76 minutes.

The question about forms of unofficial therapy was answered by 184 patients. Rather to our surprise, 100 had not gone outside the treatment that their general practitioners and we had given them. Among the various unofficial forms of therapy were the wearing of copper bangles, faith healing, acupuncture, and manipulation. Though in individual cases benefit was seen on occasion, in all but one group therapeutic results were considered by the patients as unsatisfactory. The exception was manipulation. Of the 23 patients who had received it outside hospital and general practice, 11 had rheumatoid arthritis, six osteoarthritis, three ankylosing spondylitis, and three soft tissue lesions. Nine had lasting relief—five with osteoarthritis, three with rheumatoid arthritis, and one with soft tissue rheumatism. None of the three spondylitics improved, one being considerably aggravated. Eleven patients, seven with rheumatoid arthritis, had tried acupuncture. In none

was it considered helpful by the patient and in none did it produce remission or lasting relief. Faith healing was practised by 11 patients, six with rheumatoid arthritis. In two there was temporary and in one lasting improvement, the remainder reporting no benefit. Only two of 32 wearers of copper bangles thought that any benefit resulted.—We are, etc.,

F. DUDLEY HART

J. A. WOJTELEWSKI
P. M. G. REYNOLDS

Westminster Hospital,
London S.W.1

Hyperglucagonaemia in the Surgical Patient

SIR,—The article by Mr. R. C. G. Russell and others (4 January, p. 10) prompts us to record our own experience of this phenomenon in a two-year study in which changes in the plasma glucose level have been related to the plasma levels of immunoreactive glucagon (I.R.G.), insulin (I.R.I.), growth hormone, and cortisol, and the urinary excretion of adrenaline and noradrenaline in more than 50 patients.

Plasma glucose (hexokinase method) rose significantly above basal levels by the fifteenth minute of operation, from 5.08 ± 0.12 to 6.38 ± 0.20 mmol/l (91.6 ± 2.1 to 115 ± 3.6 mg/100 ml) (mean $\pm$ S.E.M., $P < 0.005$) and reached its peak of 7.71 ± 0.38 mmol/l (139 ± 6.8 mg/100 ml) by 30 min. I.R.G. values showed no change, being respectively 165 ± 25 , 164 ± 28 and 152 ± 34 ng/l, and no rise was seen during the remaining operation time. Values of I.R.I. fell during operation. On the first postoperative morning the plasma glucose had already declined to 6.44 ± 0.14 mmol/l (116 ± 2.5 mg/100 ml) and yet the I.R.G. (286 ± 36 ng/l, $P < 0.001$) and I.R.I. values were significantly greater than at any previous observation. During the next six days the plasma glucose fell steadily and there was a parallel fall in I.R.I., yet I.R.G. values remained elevated. We would thus agree that hyperglucagonaemia is a feature of the post-operative situation but have not found any rise during operation itself.

In the conduct of such studies a number of factors may introduce bias and we have therefore studied a stereotyped response in otherwise healthy adult males whose diet, activity, drugs, anaesthetic, operation, and convalescence could be precisely controlled. We have already shown that in this model situation changes in plasma glucose correlate precisely with the surgical injury and recovery¹ and that the relationship of insulin to glucagon is influenced by more than operation alone.² The use of heterogeneous groups of patients, some of whom have serious complications, detracts from the precision of an already difficult investigation. In addition, we are uncertain why Mr. Russell and his colleagues have used the mean of two basal observations as a comparison for a single highest value obtained during any one period of the operation. Their data do, however, show that overall the greatest increases in I.R.G. occurred not during operation itself but on the first post-operative day.

While supporting the evidence that hyperglucagonaemia occurs some 16 or more hours after operation, we do not think that glucagon can yet be implicated in the immediate surgical response. In our experience the

onset of hyperglucagonaemia occurs with that of hyperinsulinaemia and both are preceded by increases in plasma glucose.—We are, etc.,

ANTHONY GIDDINGS

Royal Devon and Exeter Hospital (Wonford),
Exeter

K. J. O'CONNOR

Diabetes Research Unit,
Wellcome Foundation,
Dartford, Kent

B. J. ROWLANDS

D. MANGNALL

RONALD G. CLARK

University Surgical Unit,
Northern General Hospital,
Sheffield

¹ Giddings, A. E. B., et al., *British Journal of Surgery*, 1974, 61, 920.

² Giddings, A. E. B., *British Journal of Surgery*, 1974, 61, 787.

Glucagon and Pseudogout

SIR,—I was interested to read the paper by Mr. R. C. G. Russell and his colleagues (4 January, p. 10) on hyperglucagonaemia in the surgical patient. Glucagon has been used in the treatment of Paget's disease of bone, one of the suggested mechanisms of action being that it plays a role in pyrophosphate metabolism.¹ Calcium pyrophosphate crystal deposition disease (pseudogout, articular chondrocalcinosis) has a clear-cut but little understood association with diabetes mellitus.²

It would be interesting to know if diabetic patients who develop pseudogout or chondrocalcinosis fall into the hyperglucagonaemic group and whether those patients who have postoperative attacks of pseudogout have very high levels of plasma glucagon.—I am, etc.,

A. J. RICHARDS

McMaster University Medical Centre,
Hamilton, Ontario

¹ Condon, J. R., *British Medical Journal*, 1971, 4, 719.

² McCarty, D. J., jun., in *Arthritis and Allied Conditions*, ed. J. L. Hollander and D. J. McCarty, jun., 8th edn., p. 1151. Philadelphia, Lea and Febiger, 1972.

Genitourinary Medicine

SIR,—The council of the British Association of Urological Surgeons has discussed the proposed change of name of the specialty of venereology to genitourinary medicine and has registered its disapproval of the change. Venereologists from various centres in Great Britain have also expressed concern at this proposed change.

In your leading article on this subject (11 January, p. 51) you state that the Royal College of Physicians has approved the change of name. This is not wholly accurate because at the time this was discussed by the council of the Royal College of Physicians it was noted that the proposed change of title was not acceptable to the Scottish Health and Home Department. Again, the problem of nomenclature was at this time still being considered by the Medical Society for the Study of Venereal Diseases, and no decision had then been reached. The council of the Royal College of Physicians felt that a compromise could be attained by allowing hospitals and clinics to use the title of genitourinary medicine instead of venereology if they so wished or any other designation of their choice.

In your article you stress that genitourinary medicine should not be confused with urology, and would not encroach on the realm of urology. In spite of this you immediately state that the work of genitourinary physicians could "expand to include infertility and other diseases of the genitourinary tract when appropriate." This could only be seen as an encroachment on the realm of the urologist. May I stress again the deep concern which the council and members of the British Association of Urological Surgeons feel regarding the change of title. In a recent issue of the *B.M.J.* a junior post in genitourinary medicine was advertised under "Venereology." Surely this is an anachronism.

It would appear that the time is now opportune for the whole subject to be discussed logically not only by the Royal College of Physicians but in association with the Medical Society for the Study of Venereal Diseases, and more particularly with the British Association of Urological Surgeons.—I am, etc.,

R. A. MOGG

President,
British Association of Urological Surgeons

London W.C.2

Television Programme on Induction of Labour

SIR,—The main theme of the B.B.C.2 "Horizon" programme of 27 January was induction of labour. On this controversial subject I am not qualified to comment. However, epidural analgesia in obstetrics was also referred to and several misleading or frankly incorrect statements were made.

The information leading to these statements did not come from discussions between the "Horizon" production team and the anaesthetists working at the John Radcliffe Maternity Hospital, where much of the programme was filmed, as none took place. In particular, one patient's recollections of her care after an epidural analgesic were quite at variance with established theory and practice at this hospital. She was also allowed to state that the epidural was the direct cause of her emergency caesarean section for fetal distress. In fact, at that operation the reason for the fetal distress was found and it was in no way related to the epidural.—I am, etc.,

L. E. S. CARRIE

John Radcliffe Hospital,
Oxford

Leeds Infirmary Blues

SIR,—Your correspondents Mr. R. A. Elson and Dr. R. H. Townshend (22 February, p. 455) raise two points in regard to your leading article (8 February, p. 297) on the threatened disruption of the development of the Leeds medical school and General Infirmary. Mr. Elson questions why the Leeds situation is unique. The answer is simple. About £12m. has been committed already in relation to the project and the Hospital for Women at Leeds has been demolished as the first part of the "development." Tolerating failure to improve hospital facilities is one thing, but to live with their destruction through administrative incoordination quite another. Dr. Townshend suggests that the local population might find the necessary

capital and this raises an issue which deserves the closest consideration, but any policy decision could not be applied to Leeds in isolation.

We are very conscious of the parlous financial state of the nation, but it is in this light that the problems which have arisen are particularly serious. The detailed planning of an integrated hospital-medical school complex began as you say in 1962 with the enthusiastic support and encouragement of the Department of Health and the University Grants Committee—the bodies ultimately financially responsible. The first serious setback arose in the late 1960s, when the U.G.C. reneged on its side of the integrated plan. Subsequently a fresh plan was devised involving separate, but closely related hospital and medical school buildings. At all stages the closest liaison was maintained with the two administrative bodies concerned. Now, after more than 12 years of work, when the U.G.C.'s side of the joint plan is well on the way to completion, it seems possible that the D.H.S.S. might fail to produce its part of the scheme. It is the great financial waste which would occur in this eventuality—attributable to lack of co-ordination between two Government bodies—that is so serious from a national point of view.

Locally the implications are rather different. The initiation of the scheme occurred with demolition of the Hospital for Women, which was sited adjacent to the General Infirmary and relied upon its wide range of ancillary resources in emergency situations. Its "temporary" replacement is miles away from these resources and this involves a deterioration in patient safety standards. Apart from the patient welfare aspect, perpetuation of this on a semi-permanent basis would serve along with the £5m. energy-generating complex as memorials to the massive waste of public funds which has achieved nothing but deterioration in the local medical facilities.

It will be no surprise that, faced with such a situation, medical staff morale is on the point of collapse; problems of contracts and the like are of little significance by comparison with having to work in the shadow of this chaos. Those like ourselves who have not made a major contribution to the planning are aghast that the dedicated work of our colleagues—some part-time, some whole-time N.H.S., and some university—might at this stage be squandered. These consultants have given this effort without stint at the prime of their lives and at the cost of developing their own practices and research. To maintain dedicated doctors of the best quality in this country it must be shown to them that, whatever the financial state at any time, millions of pounds are not wasted through lack of co-ordination of expenditure from the left and right Government pockets and that their talents and enthusiasm will not be exploited.—We are, etc.,

J. C. GOLIGHER
J. S. SCOTT

General Infirmary at Leeds,
Leeds

SIR,—May I congratulate you on putting the unfortunate position of the Leeds General Infirmary complex into such clear perspective in your leading article (8 February, p. 297). As a past student I have

a deep interest in its fortunes. There are perhaps two points which could be made with the benefit of local knowledge.

Both the General Infirmary at Leeds and the medical school are buildings of unusual, and in the former case distinguished architecture. This quality is one which endears itself to those using the buildings and cannot be replaced when building in current styles. Both buildings are basically sound and, though attention would be required to bring them up to present needs functionally this would cost little compared with the task of rebuilding. The site which you mention is behind the infirmary and is available to enlarge the hospital backwards as far as is needed. Current plans are to retain only the facade of the present building and to demolish everything behind this. The medical school is to go in its entirety. Is it possible that the enforced pause in reconstruction might give time for second thoughts?

With regard to the increase in the intake of students, 80 per year ideally suited the personal teaching methods used at Leeds. With the further expansion of teaching at St. James's Hospital this number has been increased modestly without impairing the standards in any way. Should medical schools, however, go on increasing their intake to make up for losses in medical manpower which are the direct result of the policy of present and past governments?—I am, etc.,

D. J. RODGERS

Sheffield

Fibrinolysis and Venous Thrombosis

SIR,—Your leading article (16 November, p. 365) and subsequent correspondence (25 January, p. 208) relating to fibrinolysis and deep vein thrombosis suggest that drugs which increase blood fibrinolytic activity may be valuable in the prevention of deep vein thrombosis. An assumption basic to the arguments put forward in your article and by your correspondents is that a considerable and prolonged reduction in blood fibrinolytic activity usually follows surgical trauma. We believe that this is open to serious doubt.

Accurate measurement of blood fibrinolytic activity after operation is complicated by the marked rise in fibrinogen concentration that invariably occurs. Fibrinogen forms the substrate for the lysis time methods of measuring fibrinolytic activity and it has been shown that an increase in fibrinogen concentration alters the lysis times of plasma and whole blood clots both *in vitro*^{1,2} and *in vivo*.³ We applied an isotopic method⁴ for the measurement of fibrinolytic activity in 50 patients after operation. This method was used because it has been shown to be independent of alterations in the level of plasma fibrinogen.³

As we reported last year,⁵ the results showed, in contrast to previous studies using lysis time methods, that no consistent sustained decrease in fibrinolytic activity or "fibrinolytic shutdown"⁶ occurred. Indeed, by the sixth postoperative day activity was actually significantly greater than mean pre-operative values. Postoperative fibrinolytic activity was reduced only on the first post-operative day and this reduction reached statistical significance only in patients who developed venous thrombosis.