(aged 24-50) were then visited without prior warning and their height and weight measured. Their reported heights and weights were on average 0.5 in (12.7 mm) above and 6.1 lb (3.13 kg) below their actual measurements. As in Ashwell and Etchell's paper, overweight was assessed using the Metropolitan Life Insurance Company tables.² For 90% of the women the reported measurements indicated a lower degree of overweight than the actual measurements, the difference being roughly proportional to the degree of overweight.

As well as broadly classifying body size as either "overweight," "suitable," or "underweight" Ashwell and Etchell define categories of over- and underweight. Individuals 30%, 20%, and 10% above ideal weight were classed as severely, moderately, and mildly overweight respectively, with corresponding groupings for underweight. Using these categories 15 of the 40 women in our study were classified in a lower category on reported than on actual measurements. Four of these changes were from the broad classification "overweight" to that of "suitable weight."

Our data indicate that reported height and weight should be treated with great caution, as there appears to be systematic underestimation of body size. It may well be that this underestimation of body size as assessed from reported data reflects awareness of overweight status. We would suggest that any further studies on perception of overweight should obtain information on reported as well as actual height and weight and also perceived weight status.—We are, etc.,

DOUGLAS ALTMAN
LESLIE IRWIG

Department of Community Medicine,
St. Thomas's Hospital Medical School,
London S.E.1

¹ Ashwell, M., and Etchell, L., *British Journal of Preventive and Social Medicine*, 1974, 28, 127.
² Metropolitan Life Insurance Company, *Statistical Bulletin*, 1959, 40, Nov.-Dec.

SI Units

SIR,—If and when SI units are introduced into clinical practice I foresee very real difficulties in the management of patients. Apart from the expense involved in converting to SI, the management of common diseases is surely going to be very much more difficult. When one is informed that a patient in diabetic ketoacidosis has a blood sugar of 780 mg/100 ml one has an instant picture of what that patient's clinical state is likely to be. The SI unit normal range of blood sugar is 3.0-5.0 mmol/l and we are told that the approximate range found in hospital practice could vary from 0.6 to 60 mmol/l. It will not be easy to decide on insulin dosage in such patients. I would think that the same difficulties would arise in uraemic patients when the blood urea is measured in SI units.

We are told that prescribing will not be changed to SI; neither will blood pressure recordings. These have presumably been left alone because of expediency. I think many practising clinicians would feel that much the same argument could be applied to many of the parameters routinely measured in blood chemistry and that to introduce a completely new set of values at this time, when the Health Service is in dire trouble,

will not help in the management of patients.

There is still doubt that the Americans will go over to SI units and this must mean that the American and British medical literature will be in two completely different sets of units. Medicine is not a science; is empiricism now a dirty word?—I am, etc.,

B. H. BASS

Department of Medicine,
Good Hope General Hospital,
Sutton Coldfield, Warwicks

Tafracher

SIR,—I was most interested in Dr. F. I. D. Konotey-Ahulu's Personal View of the Ghanaian word *tafracher* (8 February, p. 329). Though English has no direct equivalent the sentiment has not, at least in the past, been foreign to us, as is witnessed by the expression of an Irish woman describing the death of her daughter's child to my father some years ago: "... and then, saving your presence, his little bowels gave way."—I am, etc.,

D. S. THOMPSON

London N.19

** We have received several letters drawing attention to similar expressions in use in a variety of languages and cultures.—ED., *B.M.J.*

Comparison between Free Thyroxine Index and Effective Thyroxine Ratio

SIR,—We read with great interest the paper by Dr. E. G. M. D'Haene and others (21 September, p. 708) on the comparison of the Mallinckrodt effective thyroxine ratio (E.T.R.) with the serum free thyroxine index (F.T.I.). Using Spearman's rank correlation analysis, the authors found a highly significant relationship between the E.T.R. and the F.T.I., with a correlation coefficient of 0.68. The correlation coefficient was considerably less than that found by us—namely, 0.95—when we used the Pearson product-moment correlation coefficient in a study of three groups of patients classified as hypothyroid, euthyroid, and hyperthyroid.¹ A similar correlation coefficient (0.93) was found between the E.T.R. and the free thyroxine concentration (F.T.C.). We found it necessary to select patients and subjects in order to yield diagnostic groups of approximately equal numbers. This leads to a much more sensitive regression analysis because of the wider spread of values.

Dr. D'Haene and his colleagues, however, consider that if a given variable has a normal distribution in the population it is necessary, in assessing the variable, to select test subjects (or specimens) by random sampling and hence are critical of our statistical handling of our data. However, we submit that their approach, which studies a randomly selected group of subjects, has no application in the regression model used by us. The essential point in the design is the random distribution of assay errors and not of the population from which the specimens were collected.

It seems that these authors have confused regression with correlation analysis. As pointed out by Hays,² each is appropriate to

a particular kind of research enterprise. Correlation analysis applies especially to problems of prediction. Regression analysis on the other hand determines the relationship between a set of variables—for example, does the E.T.R. increase as the F.T.I. increases and, if so, is there a mathematical rule which describes the relationship? Dr. D'Haene and his colleagues have found a poorer correlation between E.T.R. and F.T.I. because 70-75% of their specimens had normal values and because the E.T.R. has a very low coefficient of variation (C.V.) (3.7% in our hands) in euthyroid subjects. Furthermore, they would have encountered a substantial number of multiple tied values, which would also detract from the degree of correlation found, and there is no indication in their paper that they have corrected for ties.

Further studies from our laboratory³ have indicated that the E.T.R. is a good screening technique, as in euthyroid patients with low activity of thyroxine-binding proteins the diagnostic accuracy of the E.T.R. (80.5%) is superior to that of the F.T.I. (62.5%) and F.T.C. (67%). In euthyroid patients with elevated thyroxine-binding proteins the diagnostic accuracy of E.T.R. is 93.5% compared with 83% for F.T.I. and 91% for F.T.C.

Though the conclusion of Dr. D'Haene and his colleagues that the E.T.R. is not a satisfactory substitute for the F.T.I. is probably incorrect in the light of our studies, we do agree that there are some limitations in the E.T.R., as continuing studies in our laboratory have demonstrated. It is not a good parameter to use in patients undergoing triiodothyronine (T-3) suppression, nor is it a good assay in either thyroid-stimulating hormone (T.S.H.) or thyroxine-releasing hormone stimulation tests. Furthermore, the very low C.V. of the E.T.R. may indicate relative insensitivity rather than a high degree of precision. This leads to a marked overlap in values between clinically hypothyroid patients and normal subjects, particularly where treatment has been given for thyroid disorders. In clinical situations, such as that following radioiodine treatment, it is mandatory to use serum assays such as T.S.H. and T-3 concentration to supplement the E.T.R. test.—We are, etc.,

M. L. WELLBY
M. W. O'HALLORAN
JANET MARSHALL

Department of Clinical Chemistry,
The Queen Elizabeth Hospital,
Woodville, South Australia

¹ Wellby, M. L., O'Halloran, M. W., and Marshall, J., *Clinica Chimica Acta*, 1973, 45, 255.

² Hays, W. L., *Statistics for the Social Sciences*, p. 536. New York, Holt, Rinehart and Winston, 1972.

³ Wellby, M. L., O'Halloran, M. W., and Marshall, J., *Clinical Endocrinology*, 1974, 3, 63.

Venous Gangrene in Multiple Myeloma

SIR,—The incidence of thromboembolism in myelomatosis based on a preliminary estimate of 376 patients admitted to the Medical Research Council's myelomatosis trial up to March 1970 was reported to be about 3%,¹ but the incidence of gangrene of the feet as a presenting feature is uncommon. Catovsky *et al.*¹ reported 14 cases of myelomatosis associated with major