The estimated date of a recurrence is not always recorded by a registry, although it gives an indication of the *quality* of life after treatment as opposed to total survival time. Histology is often not verified, although this varies with region.

4. DATA ANALYSIS

Techniques used to calculate T-year survival rates vary from the direct method to the life table or actuarial method^{3 4} and the prediction model method^{5 6} and may or may not be adjusted for the age and sex distribution of the particular group of cancer patients.⁷ This should always be made clear to enable comparisons to be made.

5. DATA PRESENTATION

Presentation of data is largely a matter of common sense, but the publications by Reichmann⁸ and Mould⁴ may prove helpful.

DHSS SH3 returns

Other figures are produced yearly for the DHSS on hospital return form SH3. These returns are subdivided by medical specialty and include data on "new" patients, attendances, and treatments during the year separated into inpatients and outpatients. Instructions for the completion of the form are ambiguous in many places. Since statistics on cancer treatment work load must in part determine future policy such as purchase of equipment, the consequences of inaccurate returns are serious. The definition of the term "new patient" varies. For the purposes of the SH3 a new patient for the registration year is defined as one who starts a new course of treatment in that year, regardless of whether previous treatment courses have been given. If hospitals do not realise that a single patient may be counted more than once for SH3 forms the radiotherapy work load will be underestimated. The table gives some 1973 SH3 statistics for eight different hospitals, and there seem to be inconsistencies. These are probably in part explained by (a) some of the patients originally seen at outpatient clinics

receiving treatment as inpatients (number of new inpatients treated in 1973: hospital A 756, B 516, C 325, D 156); (b) outpatients seen but not treated not being recorded (hospitals E and F)—these will probably form a minority and include those of advanced age and with advanced disease at initial presentation—and (c) patients referred for radiotherapy by other consultants not having also been included under the radiotherapy section.

Figures derived from parts 2 and 3 of hospital return form SH3 completed by eight hospitals (A-H) for the year 1973 showing apparent inconsistencies in numbers of new outpatients

Hospital	New outpatients		Implications!!!
	Part 2, SH3: consultation statistics, line 18	Part 3, SH3: treatment statistics, line E(c)	
A	3248	1841	1407 new patients seen but not treated
B	1868	1335	533 new patients seen but not treated
C	1271	927	344 new patients seen but not treated
D	944	825	119 new patients seen but not treated
E	1560	1560	All patients seen were treated
F	241	241	All patients seen were treated
G	838	1248	410 patients were treated but not seen
H	796	892	96 patients were treated but not seen

References

- ¹ *British Medical Journal*, 1975, 4, 245.
- ² Office of Population Censuses and Surveys, *Report of the Advisory Committee on Cancer Registration*, p 10. London, HMSO, 1970.
- ³ Ministry of Health, *Reports on Public Health and Medical Subjects*, No 33. London, HMSO, 1926.
- ⁴ Mould, R F, *Clinical Radiology*, 1976, 27, 33.
- ⁵ Boag, J W, *Journal of the Royal Statistical Society* (series B), 1949, 11, 15.
- ⁶ Mould, R F, and Boag, J W, *British Journal of Cancer*, 1975, 32, 529.
- ⁷ Berkson, J, and Gage, R P, *Proceedings of the Staff Meetings of the Mayo Clinic*, 1950, 25, 270.
- ⁸ Reichmann, W J, *Use and Abuse of Statistics*. Harmondsworth, Penguin, 1973.

Every emergency department should have one— an interview room

D H WILSON

British Medical Journal, 1976, 1, 87-88

The accident and emergency department is at the interface between the hospital and the outside world, and it is the focal point for many acute social catastrophes. Parents may have to be told that their children have been killed on the road, or a wife may learn that her husband has just had a fatal coronary thrombosis. A room should be set aside for such devastating interviews.

Design of the room

In 1971 I was given the opportunity of helping to design a new accident and emergency department for the General Infirmary at

Leeds. In the first draft of the operational policy I asked that a pleasant quiet room should be designed close to the reception area for personal interviews with patients' relatives. I had already experienced the seemingly callous way in which it was sometimes necessary to inform relatives of a sudden brutal death and then expose their grief to public gaze. The secretary's office, the sister's office, or even the staff sitting room are used in some departments as a haven for distressed relatives, but this can embarrass both the relatives and the staff and disrupt the work of the department. I had come firmly to the opinion that a room specifically designed for this purpose was essential.

Eventually, as the plans for our new department developed, we located the interview room next to the new patients' waiting area and close to the nurses' station. This has proved to be an ideal arrangement. The door opens discreetly off the waiting area and only the nurse can see directly in. The room is carpeted, has soft lighting, and is furnished with easy chairs grouped around a coffee table. Behind the door is a hand basin and mirror and on the table a telephone to enable people to contact other relatives or friends.

In the six months since these new premises came into use we have kept a record of the various purposes for which the interview room has been used. Normally the room is kept locked but the key is in the nurses' station. When relatives leave the room is locked again, the key is returned to its hook, and details of the interview are entered in a

Accident and Emergency Department, General Infirmary, Leeds
LS1 3EX

D H WILSON, FRCS, consultant surgeon

register. This register now has page after page recording catastrophic personal tragedies. It makes very sombre reading, but the staff are all agreed that this new facility has greatly improved the way in which we are able to care for the relatives as well as the patient.

Using the room

We have had to break the news of the death of a son or daughter from road accidents, drowning, or acute illnesses. Other parents have learnt that their child has suffered severe brain damage or physical mutilation from a road accident, that they are still alive, but that they have little chance of making a full recovery.

Among adult patients myocardial infarction is by far the commonest cause of unexpected sudden death. Attempts at resuscitation may persist for an hour or more and we have found it invaluable to have this room where relatives can wait quietly and privately until we have some definite news for them. If the resuscitation fails they may wait until it is convenient for them to view the body or collect any valuables. During this time they can use the telephone to contact other friends or relatives who can share their anxiety or grief with them.

Apart from "informing the relatives" the room has also been used to interview eye witnesses, workmates, or relatives about the circumstances of an accident or, in one instance, to talk with the remorse-filled 16-year-old father of a battered baby. Staff from other departments of the hospital have also occasionally used the room to talk with relatives who have arrived when it is inconvenient to see them on the ward.

The onerous task of passing on sad news is certainly made easier if it is shared. I have often greatly appreciated the help of a senior nurse. I start the interview by making sure of the relationship between the injured or deceased patient and the people with whom I am speaking to avoid the possible anguish of mistaken identity. Then, having told

them what has happened, I find that a female nurse in uniform is a more readily acceptable source of consolation than a male doctor in a white coat. For most people the offer of a cup of tea is much appreciated. Our hospital chaplain recently pointed out to me that it is impossible to cry and drink tea simultaneously. For people who want to stop crying but cannot find the strength to do so a cup of tea may enable them to regain control of their emotions. They then may want to ask questions about necropsies, inquests, and funerals or just to sit and talk about the person who has died. A few minutes spent listening sympathetically is probably of more value than some stereotyped words of consolation, and physical contact by an arm around the shoulder or holding a hand is often more eloquent than words.

The hospital chaplains, the medical social workers, and the police have all been helpful at times in consoling and counselling patients' relatives. The police also often help in taking relatives home, especially at night.

Future use

Although the interview room was originally designed as a place for passing on catastrophic news, the consultant staff of the department can see a further use for it in the future as a room for counselling patients about the circumstances which led up to their visit to the accident department or seeking to help them to face up to the social environment which they are returning to after their treatment. Our brief experience has already convinced us that every emergency department should have an interview room where members of the staff can talk with patients or relatives with quietness and dignity and so seek to help them with the problems and tragedies of life and death.

A patient in her 50s suffers from separation of some of her fingernails from the nail bed almost as far as the lunule. There is no obvious disease and her general health is good. What reasons are there for this and is there any treatment?

Onycholysis (separation of the nail from its bed) is a common complaint. Diseases which may be responsible include psoriasis (it is rarely the only sign), fungus infections, dermatitis of the finger tips, thyroid disorders, poor peripheral circulation, and hyperhidrosis. It has been noted in people taking photosensitising drugs like demethyl-chlortetracycline. In many patients there seems to be no obvious cause. Idiopathic onycholysis is usually confined to women. Repeated minor trauma may be an important cause, particularly if the nails are kept too long. The gap between the nail and the bed often becomes infected with bacteria (especially *Pseudomonas aeruginosa*) and yeasts, causing discoloration of the nail. Any associated disease should be treated on its own merit. There is no specific treatment for idiopathic onycholysis. The nails should be kept short and Samman¹ states that the application of 15% sulphacetamide in spirit or 22% thymol in chloroform is useful. These help to combat secondary infection, which is probably the main reason for the nail failing to reattach itself to the bed.

¹ Samman, P D, *The Nails in Disease*, 2nd edn. London, Heinemann, 1972.

What is the best advice to an elderly patient who has difficulty opening his bowels? Investigations have proved negative but he has taken laxatives for 50 years.

A patient who has taken laxatives for 50 years is unlikely to be helped by changing from one purgative to another. It is important to review other medication. Many drugs cause constipation. Codeine is well known, as are all drugs with an anticholinergic action. These include antispasmodics, anti-Parkinsonian remedies (but not levodopa or amantidine), phenothiazine tranquillisers, many hypotensives, and tricyclic antidepressants. Iron preparations, antacids—particularly calcium carbonate and aluminium hydroxide—barbiturates, benzodiazepines, and monoamine oxidase inhibitors may cause constipation. Diuretics remove fluid from the stool and predispose to constipation.

This may also lead to potassium loss and muscular weakness may be an additional factor. He should take extra fluid, particularly beer, and increase the fibre content of the diet. Bran is particularly effective, but start with one or two teaspoonfuls of natural bran with a breakfast cereal at first, and if possible work up to two heaped dessertspoonfuls. Other bulking agents, Isogel, Normacol, and Celevac, may be tried as alternatives to bran. The risk of bolus formation with these agents is reduced if they are taken with plenty of fluid and more than once a day.

Is there any value in transcendental meditation?

"Value for whom?" must be the counter-question. There can be no doubt that in certain cultures religious beliefs of a predominantly mystical sort, including transcendental meditation, are of value to their practitioners. In our European culture, however, there can be very few genuine adherents of such religions, and certainly it would be unacceptable for, say, a local authority health committee to make provision for them. Those who feel a need for such religious experience seem capable of finding their own kind either at home or, more frequently, in the Orient. It is worth mentioning that some schizophrenics find satisfaction in contemplating their religio-philosophical navels.

How great is the risk of lead-poisoning from the glaze of the old-fashioned earthenware gallon/two gallon jars, used in homemade beer and wine making?

The Pottery (Health) Special Regulations 1947 prohibit the use of earthenware glazes other than those described as "leadless" (a glaze not containing more than 1% dry weight of a lead compound calculated as lead-monoxide) or "low solubility" (a glaze that does not yield to dilute hydrochloric acid more than 5% of its dry weight of a soluble lead compound calculated as lead monoxide). Hence the use of such vessels for home brewing or winemaking should not present a lead-poisoning hazard. Vessels of unknown origin, however, may present a hazard.