frequently be avoided if the subjects show the fixity of purpose described here; their determination to minimise the wind-chill factor (by digging a substantial snow cave) and to overcome the urge to sleep undoubtedly prevented a serious outcome for this party, which carried no specific equipment to spend a night out near the summit of Everest.

CHARLES CLARKE
Medical Officer,
British Everest Expedition, 1975

Middlesex Hospital Medical School,
London W1

¹ Ward, M., *Mountain Medicine*. London, Crosby Lockwood Staples, 1975.

² Smythe, F. S., in *Everest*, 1933, ed H. Rutledge. London, Hodder and Stoughton, 1934.

Heart block and autoimmunity

SIR,—In reply to Dr F I Lee (29 November, p 524) the four patients quoted in our paper who had chronic heart block and hypothyroidism did not develop these disorders simultaneously. In three patients heart block occurred between four and eight years after they had become clinically hypothyroid, while they were on replacement therapy. The fourth patient, whose history was quoted, was clinically euthyroid when heart block developed, but a raised thyroid stimulating hormone level at this time, together with evidence of multiple autoimmune disorders, led to the institution of thyroid replacement therapy, which has not, after 12 months, reversed his conduction disturbance. None of these patients had evidence of ischaemic heart disease but all had thyroid autoantibodies present. As in Dr Lee's case, the irreversibility of heart block by administered thyroid hormone suggests that a direct metabolic effect alone is insufficient to explain these data.

Permanent damage to the sinoatrial node has been reported in thyrotoxicosis,¹ and recently "lone" atrial fibrillation has been associated both with "silent" thyrotoxicosis and, surprisingly, with early hypothyroidism.² An autoimmune aetiology seems to be a possible unifying factor to explain these observations, and it is possible that circulating immune complexes, which have been proposed as a cause of the skeletal muscle damage in thyrotoxic ocular myositis,³ could be one of the mechanisms by which the heart is damaged in these autoimmune disorders.

A J FAIRFAX
A LEATHAM

Department of Cardiology,
St George's Hospital,
London SW1

¹ Wan, S. H., Lee, G. S., and Toh, C. C. S., *British Heart Journal*, 1972, **34**, 942.

² Myers, A. et al., *Abstracts of the British Cardiac Society Autumn Meeting*, 1975, p 20.

³ Konishi, J., Herman, M. M., and Kriss, J. P., *Endocrinology*, 1974, **95**, 434.

First-aid treatment of poisoning

SIR,—Your issue of the 29 November contained two items on this topic, a letter on "Salt overdose" (p 517) and the leading article on "Childhood poisoning: prevention and first-aid management" (p 483). On the first item we informed you in 1974 (9 November, p 342) that the joint manual of the voluntary aid societies would be modified

to omit the use by first-aiders of the induction of emesis in the treatment of swallowed poisons; this has now been done.

At the time we decided to make this modification we considered recommending the use of ipecacuanha in the first-aid treatment of childhood poisoning. We concluded that to try to define for first-aiders the indications for this procedure would lead to confusion and that the occasions on which first-aiders would be required to administer ipecacuanha were so few that it would be wasteful of resources and contrary to the aim of simplicity in first-aid equipment to provide it.

ANDREW RAFFLE
Chief Medical Officer,
St John Ambulance Association

JOHN GRAY
CMO, British Red Cross Society

HECTOR R F MACDONALD
CMO, St Andrews Ambulance Association

Chronic peritoneal dialysis for diabetic nephropathy

SIR,—Your leading article (16 August, p 397) entitled "Renal transplantation in diabetes," concludes that renal transplantation is probably the most promising treatment for patients with end-stage diabetic kidney disease. Renal transplantation may be the preferred treatment in younger diabetic patients. The mean age of those transplanted in Minneapolis was 34 years, the oldest patient being 49.¹ Success is more likely in those lacking severe cardiovascular disease and other complications of diabetes and in circumstances where a living donor is available. However, such patients make up only a small fraction of the total number of patients with end-stage uraemia due to diabetic nephropathy.

In the United States, with the recent enactment of new social security Medicare laws, renal replacement therapy is becoming feasible for nearly all patients with end-stage renal disease. The magnitude of the problem is great. It was conservatively estimated that in the US approximately 15 diabetic patients develop end-stage nephropathy per million population per year (3200/year in the US).² Many of these patients are in the older age group and have multiple complications of diabetes. For this group of patients transplantation is not feasible. The difficult medical and moral problem has been whether these patients are best treated by dialysis or by "conservative measures," which implies that they will soon die as the consequence of the disease. The results of dialysis therapy in this group often prove less satisfactory than in non-diabetic uraemic patients, but an increasing number of dialysis centres are accepting these patients for care. Although the results of haemodialysis in this population have for the most part not been good, maintenance peritoneal dialysis may offer a promising alternative in the patient with diabetes and end-stage uraemia.^{3,4} The mean age of the eight diabetic patients we have treated with maintenance peritoneal dialysis at this centre is 54 years. This mode of therapy may be better suited for the patient with severe cardiovascular disease who is prone to haemodynamic instability and arrhythmias. Since rapid fluctuations of blood pressure

are less frequent and systemic anticoagulation is not employed eye disease may not progress at as rapid a rate with haemodialysis, while problems of vascular access necessary for haemodialysis are obviated.

None of the current methods available for managing end-stage uraemia in diabetic patients provide the chances for rehabilitation seen in the uraemic patient without diabetes. In the older diabetic patient with severe cardiovascular disease it is likely that the physician, the patient, and the patient's family must accept more limited goals. Under such circumstances peritoneal dialysis may offer certain advantages over haemodialysis: it can be safely carried out at home even without the presence of another person, and a patient can be trained for self-care at home in less time than is required for haemodialysis. Thus a well-motivated patient may achieve satisfactory rehabilitation while peritoneal dialysis is performed during sleep three to five nights per week.

M J BLUMENKRANTZ
J E BAILEY
M YARON
J W COBURN

Medical Service,
VA Wadsworth Hospital Center, and
Department of Medicine,
UCLA School of Medicine,
Los Angeles, California

¹ Kjellstrand, C. M., et al., *Kidney International*, 1974, **6**, suppl. p 15.

² Knowles, H. C., *Kidney International*, 1974, **6**, suppl. p 2.

³ Jones, K. M., and Ivanovich, P., *American Society of Extracorporeal Technology Proceedings*, 1974, **2**, 97.

⁴ Blumenkrantz, M. J., et al., *Kidney International*, 1974, **6**, suppl. p 108.

Immunisation against whooping cough

SIR,—During the past year you have published several reports and letters about whooping cough, including notably the report of the Joint Committee on Vaccination and Immunisation (20 September, p 687). None of these publications offered data for epidemiological analysis, so readers were dependent for guidance upon the recommendations of the committee and of your leading article (25 October, p 186). In this article you referred to "evidence" that current vaccines give such a high degree of protection that experts are "almost unanimous" in recommending its use routinely, thus endorsing the view of the committee that the decline in the incidence of whooping cough since 1958 is attributable to immunisation.

Immediately after reading this leading article I sent to you a short paper, based largely upon data available to the committee, in which I questioned the assumption that routine immunisation was responsible for the decline in the incidence of whooping cough. I drew attention also to the continuing importance of overcrowding and other socioeconomic factors which appeared, in the UK generally and in Glasgow especially, to be at least as important as lack of immunisation in determining the distribution of recent outbreaks. So that the "evidence" could be assessed independently I stated that the raw data, national and local, on which my conclusions were based would be made available to anyone who wanted it through the Communicable Diseases (Scotland) Unit at Ruchill Hospital, Glasgow G20.