low perinatal mortality rate, whereas in fact it had been shown that "those parts of Holland with the highest incidence of hospital confinement have the lowest perinatal mortality rates."

Secondly, they appear to suggest that the physical safety of mother and baby measured in terms of mortality and morbidity should not be the overriding preoccupation. What more important considerations are conceivable? It is surely a truism to state that it is better that the mother and child should be delivered in an efficient maternity unit where unpredictable emergencies can be rapidly treated (although they occur with comparative infrequency) rather than in an inevitably improvised fashion in the patient's own home. I believe that a great majority of doctors think hospitals could be made more friendly and less impersonal with obvious benefit to patients and indeed to themselves. Nevertheless, the prime objective of safety to mother and infant should not be obscured by more nebulous factors such as those mentioned in the study group's letter.

MARK REID

Craigavon Area Hospital,
Craigavon,
Co Armagh

SIR,—I read with interest the correspondence (31 January, p 279) on the question of domiciliary or hospital confinement. This is a matter which is rightly generating concern in both public and professional quarters since it is increasingly clear that the research which has led to major advances in clinical obstetric practice has not been adequately matched by research into childbirth as a social experience. Undoubtedly, the technical quality of modern obstetric care is a great source of pride to its practitioners, but it is also apparent that it leaves many patients cold and is responsible for the sort of nostalgia about which Professor A G M Campbell is rightly sceptical. It is a matter for regret that the policy of 100% hospital confinement has been pushed through without adequate public, as opposed to professional, debate, but I think it is unrealistic to suppose this can be easily reversed or would be clinically desirable. What should be a matter of concern, though, is the social climate of our maternity hospitals which allows so many women to feel so unhappy about their experiences in them.

Both Professor Campbell and the National Childbirth Trust group agree on the need to involve patients and their families more closely in clinical decision-making. I would not wish to decry this, but I would suspect that the changes which are needed must be both deeper and more extensive. Such a question, however, could be settled only by a thorough review of the social aspects of maternity care, examining the social interactions which take place between the various parties who are involved with the service—obstetricians, midwives, social workers, health visitors, ancillary staff, patients and their friends and kinfolk. The maternity hospital is a social organisation through which people come to make sense of a biological event and to respond to it on the basis of that interpretation.

In the past there has been a shortage of interested medical sociologists with the relevant theoretical and methodological training to carry out ethnographic studies of maternity services grounded in precise and detailed observations of their social life. There has, too,

been a certain reluctance to commit research funds and for obstetricians to extend the sort of access which such investigations require. This situation is changing slowly, but such work as is being done is somewhat partial and fragmentary, although useful both intrinsically and in persuading maternity staff that they have no grounds for feeling threatened by the presence of adequately trained observers in their workplaces. It is perhaps now time to consider whether a more systematic and comprehensive programme of studies should not be set up. This should be able to compare practices in both high- and low-technology settings, in teaching, non-teaching, and GP units, and in domiciliary practice. It could provide some solid evidence for a debate which is so far founded largely on assertion. In this it represents a challenge to both the "back to nature" and the "forward with technology" lobbies. If each is genuinely worried about the activities of the other, then they should welcome such inquiries into their own practice.

ROBERT DINGWALL

Institute of Medical Sociology,
Aberdeen

Immunisation against whooping cough

SIR,—Professor G T Stewart (10 January, p 93) regrets the sharp decline in childhood immunisations which has followed the 1974 adverse publicity and which results from "continuing doubts . . . about the safety and efficacy" of pertussis vaccine. Regarding safety, why was there no reply to last year's letter from Dr A H Griffith (8 November, p 347)? What is the evidence, if any, that pertussis vaccine may be too dangerous to use? Regarding efficacy, there was no reply to a letter published last May¹ which showed that current vaccine was highly effective in the 1970-1 and 1974-5 outbreaks and which called for "evidence that it does not protect."

Professor Stewart (31 January, p 283) provides no such evidence, either in your correspondence columns or in the raw data that he has kindly sent me, but makes misleading deductions from articles by Dr Christine L Miller and Mr W B Fletcher (17 January, p 117) and Dr N D Noah (p 128). He asks, "What kind of epidemiology is this which advocates immunisation by excluding . . . factors other than immunisation?" He may be right that "immunisation schedules are often better maintained . . . where socioeconomic conditions are favourable"; but this does not justify his deduction that "association between protection and immunisation could be an expression of better social conditions and child care." In both the 1970-1 and 1974-5 outbreaks,^{2,3} pre-1967 vaccine was associated with poor protection and current vaccine with good protection. What evidence is there that social conditions suddenly improved at precisely the time when the vaccine was modified?

Moreover, we must ask, what kind of epidemiology is *this* which equates actual cases of pertussis infection (bacteriologically confirmed) with mere notifications, based on clinical diagnosis, of varied accuracy? At least, Drs Miller and Fletcher were aware of the difference: "The survey was limited . . . by its dependence on notified cases"; and so was Dr Noah: "Some of the cases notified are probably due to organisms other than *Bordetella pertussis*, against which whooping-cough vaccine would provide no protection."

Again, Professor Stewart says, "There is no

evidence . . . that immunisation of older children protects younger ones." It "leaves those . . . below 6 months of age unprotected." Is this true? There is, at least, a "widely held belief"—to use his own phrase—that whooping cough is taken home by a child of school age; and several family infections in the 1974-5 outbreak illustrated such transmission. Is there any evidence that immunity of older children does *not* protect their younger siblings?

To conclude, Professor Stewart's misleading letters could undermine the confidence in current vaccine which you so rightly encouraged last year (25 October, p 186). Hence the urgent need to provide this antidote, pending completion of my analysis of the 1974-5 outbreak. Suffice it to say at present, for the immediate guidance of those still unsure, that the trend, which was reported³ after a study of only 37 children, has now been amply confirmed by data from nearly 400 bacteriologically confirmed cases. Current pertussis vaccine does give good protection.

NOEL W PRESTON

Department of Bacteriology and Virology,
University of Manchester

¹ Preston, N W, and Stanbridge, T N, *Lancet*, 1975, **1**, 1089.

² Preston, N W, and Stanbridge, T N, *British Medical Journal*, 1972, **3**, 448.

³ Preston, N W, *Lancet*, 1974, **2**, 1138.

SIR,—The report "Severity of notified whooping cough" by Dr Christine L Miller and Mr W B Fletcher (17 January, p 117) was, as its title implies, concerned solely with the severity of *notified* cases of whooping cough and under these circumstances severity was found to be related to both age and previous immunisation. Because of the current schedule there were no fully immunised children under six months, but from the age of one year the proportion of vaccinated children admitted to hospital and severely ill at home was less at all ages than the proportion unvaccinated. The conclusion that previous vaccination reduced the severity of the disease seems perfectly reasonable and Professor Stewart's criticisms do nothing to invalidate it.

This report was, as it clearly stated, not concerned with *prevention* of the disease by vaccination and Professor Stewart is mistaken in inferring that the relative proportion of vaccinated and unvaccinated cases which he quotes from it can provide any such evidence. An assessment of the efficacy of the vaccine requires a comparison of the attack rates in vaccinated and unvaccinated children and Dr N D Noah's report (17 January, p 128) was concerned with this aspect. His findings were indeed combined in a single table—for the sake of brevity—but as clearly stated in the text "for children born after 1969 the whooping-cough notification rates were invariably greater in unimmunised children in every period and every age group."

T M POLLOCK

Epidemiological Research Laboratory,
Central Public Health Laboratory,
London NW9

It's a baby!

SIR,—I recently had the opportunity of seeing Henry Kemp's untitled film depicting several mothers' reactions to the birth of their babies. As seen on the film, and from my own experience, it is common practice to declare the

sex of the baby at an early stage, generally before the mother has handled or often even seen the baby. By chance alone approximately half the mothers are going to be mildly to markedly disappointed in the sex of their offspring. In such cases there is a risk that the set of the mother towards her child may be detrimentally affected.

In the terminology of ethology the binding of the mother to her child is dependent on certain releaser stimuli. The clinical circumstances in which the first contact takes place between the mother and her newborn child can hardly be conducive to what is sometimes a precarious binding process.¹ However, it would be possible to avoid declaring the sex of a newborn baby and allowing the mother to discover this for herself. Satisfaction of her curiosity will necessitate both looking at and handling the baby to discover the sex. This exploration may provide important releaser stimuli²⁻⁶ uncontaminated by maternal disappointment. To avoid unnecessary anxiety the practice of not declaring the sex at birth could be explained beforehand.

M WELLER

Royal Victoria Infirmary,
Newcastle upon Tyne

- ¹ Kennell, J. H., *et al*, *Developmental Medicine and Child Neurology*, 1974, **16**, 172.
- ² Collias, N. E., *Ecology*, 1956, **37**, 228.
- ³ Hersher, L., Moore, A. U., and Richmond, J. B., *Science*, 1958, **128**, 1342.
- ⁴ Klaus, M. H., *et al*, *New England Journal of Medicine*, 1972, **286**, 460.
- ⁵ Barnett, C. R., *et al*, *Pediatrics*, 1970, **45**, 197.
- ⁶ Klaus, M. H., *et al*, *Pediatrics*, 1970, **46**, 187.

Laparoscopy explosion hazards with nitrous oxide

SIR,—Readers of your correspondence columns may be confused as to the difference between hysteroscopy, the subject of our letter (1 March 1975, p 511) and laparoscopy (Professor J. S. Robinson and others, 27 December, p 760). During hysteroscopy the uterine cavity is insufflated with gas or liquid to allow telescopic inspection, whereas with laparoscopy it is the peritoneal cavity which is insufflated with gas. The changes we observed in PaCO₂ during hysteroscopy have indeed been further investigated, and our original thoughts (1 November, p 288) seem to have been confirmed.

Turning back to laparoscopy we find it hard to comprehend the statement by Professor Robinson and his colleagues concerning "initial opening of the bowel by diathermy" as this is not usually a part of laparoscopic sterilisation, although it may occur as a complication.¹ However, diathermy of the bowel may be part of an elective surgical procedure, and N₂O is almost universally used during anaesthesia. Explosions such as those described by the Birmingham workers (27 September, p 764 and 27 December, p 760), although unfortunate, do not seem to be very frequent. Nevertheless, an explosion in a closed cavity may be more hazardous than one in an open cavity.

Leaving the above points aside, we are in agreement with Professor Robinson and his colleagues that more work needs to be done in this area. Nevertheless, for those performing tubal diathermy the choice lies between using CO₂, with some risk of cardiovascular collapse (on occasion fatal), and N₂O, with a theoretical risk of causing an intraperitoneal explosion. Regrettably the data currently available do

not permit, in our opinion, quantification of these two risks.

I M CORALL
J A ELIAS
LEO STRUNIN

King's College Hospital,
London SE5

- ¹ Rawlings, E. E., and Balgobin, B., *British Medical Journal*, 1975, **1**, 727.

"Whites only" at the University of Natal

SIR,—On 17 December the cabinet of the South African government sent instructions to the Medical Faculty of the University of Natal to stop admitting first-year African medical students. This was done without consulting and with the subsequent opposition of the Faculty.

The Medical Faculty of the University of Natal was established in 1951 to educate non-White medical students. It has developed over the past 25 years into a leading medical school and is the main place of medical education for non-Whites in South Africa. Limited numbers of non-White students are educated at Cape Town and Witwatersrand universities, while the White students have a choice of five medical schools.

Now that the Natal Medical Faculty is established as a leading centre the South African government has decided unilaterally to take it over for White students. First, African students are to be excluded and will have to go to an African medical school north of Pretoria in the intellectual wilderness of the "Homelands." This is not yet built and will not produce doctors before 1982. The Indian medical students will then be excluded and will also have to start up a new faculty in Durban in a run-down tuberculosis hospital—King George V Hospital.

Thus once again the South African government is taking the best for the White community and the disfranchised non-Whites become the victims of a political ideology that must be condemned by all people of the democratic Western world.

SOUTH AFRICAN DOCTOR

Durban

Steatorrhoea complicating therapy with mefenamic acid

SIR,—We were interested to read the report by Drs J. S. Marks and M. H. Gleeson (22 November, p 442) describing mefenamate-induced steatorrhoea.

We have seen two such cases in women aged 44 and 69 years respectively, who had profuse steatorrhoea, abdominal distension, and weight loss during mefenamic acid therapy. One became asymptomatic when ibuprofen was temporarily substituted for mefenamic acid, but the diarrhoea recurred on restarting mefenamic acid. Faecal fat output was increased (14.2 and 12.5 g/24 h respectively), with normal serum vitamin B₁₂ and folate levels. However, jejunal biopsy appearances were abnormal. The villi were irregular and many leaf-forms were present. There was plasma-cell, macrophage, and neutrophil-cell infiltration. Goblet cells were increased in numbers, both in the crypts, which were hyperplastic, and in the surface epithelium. These appearances reverted to normal on stopping therapy.

Experiments in animals have shown that mefenamic acid and related compounds cause small-bowel mucosal damage and ulceration.¹ This is also seen in indomethacin therapy. It is worthy of note that both drugs are recirculated enterohepatically.^{2,3} Our results suggest a direct toxic action on the small-bowel mucosa.

R G CHADWICK
A HOSSENBOSCH

General Hospital,
Southampton

D G COLIN-JONES

Queen Alexandra Hospital,
Cosham, Portsmouth

- ¹ Kaump, D. H., *Annals of Physical Medicine*, 1967-8, **9**, suppl, p 16.
- ² Brodie, D. A., *et al*, *Toxicology and Applied Pharmacology*, 1970, **17**, 615.
- ³ Glazko, A. J., *Annals of Physical Medicine*, 1967-8, **9**, suppl, p 23.

Laboratory proficiency

SIR,—Your leading article (3 January, p 5) refers to the objective assessment of histological opinions. Semantics apart, a consultant histopathologist is appointed after a long training and must defend his opinions daily with clinicians who together quickly establish the measure of his professional competence. Finally, difficult and questionable necropsy findings are commonly debated at clinicopathological conferences, which in our view provide the best form of medical audit that we can devise. Medicine, including pathology, is still a long way from being an exact science.

RICHARD GOODBODY
JOHN GUTHRIE

Department of Histopathology,
Southampton General Hospital,
Southampton

Hazards of cephalosporins in penicillin-allergic patients with meningitis

SIR,—In discussing the treatment of meningitis and encephalitis Dr C. C. Smith (8 November, p 335) states a number of times that when the patient is allergic to penicillin cephaloridine should be used intravenously and perhaps intrathecally.

We suggest that this is not the best alternative therapy. Firstly, if cephaloridine is to be used then intrathecal therapy is mandatory. None of the cephalosporins efficiently penetrate the blood-brain barrier.¹ A recent review of 106 cases of bacterial meningitis treated with cephalosporins² indicated that response to intravenous therapy was poor unless accompanied by simultaneous intrathecal administration of the drug. Cephaloridine, the cephalosporin which crosses the inflamed meninges with the least inefficiency, did not achieve therapeutic levels in the cerebrospinal fluid despite large intravenous doses.

Secondly, there is evidence that patients who are allergic to penicillin have an increased risk of allergy to cephalosporins as well. In a recent review of nearly 10 000 patients receiving treatment with cephalosporins Petz³ showed that the overall incidence of allergic reactions was 1.2% in those with no history of allergy to penicillin. However, this was increased nearly five-fold to 8.7% in those who did have a positive history of penicillin allergy.

Chloramphenicol has neither of these disadvantages. It crosses the blood-brain barrier