Comparison of mean times ($\pm$ SD) taken to reach various labour indices

	Fixed dose	Incremental dose
Induction-IC (min)	27.5 $\pm$ 14.06	10.95 $\pm$ 10.65 (P<0.05)
IC-AC(h)	2.50 $\pm$ 1.39	0.96 $\pm$ 1.16 (P<0.005)
IC-5 cm (h)	3.76 $\pm$ 3.45	2.80 $\pm$ 1.43
IDI (h)	10.21 $\pm$ 6.27	5.25 $\pm$ 2.98 (0.5>P>0.01)

IC = Initial contractions (recorded tocographically).

AC = Adequate contractions (painful).

5 cm = End point of trial.

IDI = Induction-to-delivery interval.

Comment

The buccal route does not have any disadvantages when compared to the oral route of prostaglandin administration. The incidence of vomiting (25%) described for the oral route by other workers is similar.^{2,3} The buccal route also has the advantages of oral drug administration. These are staff convenience and patient comfort, the latter being without the discomfort of an intravenous infusion and the attendant danger of circulatory overload. It may, however, have the added advantage in that the tablet may be removed if uterine hyper-tonus, or any other side effect, maternal or fetal, should occur. Administering prostaglandin E₂ by the buccal route is a satisfactory method for inducing labour at term. An incremental mode of administration is advantageous.

The prostaglandin E₂ tablets were kindly supplied by Dr B S de Wet, Upjohn, South Africa. Dr June Juritz kindly did the statistical tests.

¹ Scher, J, *et al*, *South African Medical Journal*, 1972, **46**, 2009.² Thiery, M, *et al*, *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1974, **81**, 303.³ Haeri, A, *et al*, *Journal of Obstetrics and Gynaecology of the British Commonwealth*, 1975, in press.

Department of Obstetrics and Gynaecology, University of Cape Town, Cape Town

J SCHER, MD, MRCP, consultant

P DRYDEN, MB, BS, registrar

D A DAVEY, PHD, FRCOG, professor

Systemic lupus erythematosus and sickle-cell anaemia

In view of the relationship between immunity deficiency states and immune complex diseases such as systemic lupus erythematosus (SLE) a possible association between SLE and sickle-cell disease, in which there is evidence of immunity deficiency, is of interest.

Case reports

Case 1—A 50-year-old Black woman was first seen in 1956 with a six-month history of pleuritic chest pain and arthritis of the fingers, wrists, and elbows. Homozygous sickle-cell disease was diagnosed from alkaline haemoglobin electrophoresis, a positive sickling test, and a characteristic blood film. Levels of Hb A₂ were not measured but were not visibly raised on paper electrophoresis. Haemoglobin was 6.5 g/dl, mean corpuscular haemoglobin concentration 27 g/dl, reticulocytes 13%, and fetal haemoglobin 13%. During the next 10 years she continued to have exacerbations of the arthritis. In 1967 she was readmitted in congestive cardiac failure. Her blood pressure was 150/90 mm Hg, she had spindling of the proximal interphalangeal joints, active synovitis of the wrists, and finger clubbing. Her temperature varied between 37.2°C and 37.9°C. Haemoglobin was 8.8 g/dl, reticulocytes 3.5%, and white cell count 5.6 $\times$ 10⁹/l. The antistreptolysin O titre was normal, serum total protein 68 g/l and albumin 32 g/l, blood urea 7.8 mmol/l (47 mg/100 ml) rising to 17 mmol/l (103 mg/100 ml), ESR 143 mm in 1 h, venereal disease research laboratory test negative. Nine blood cultures were sterile. LE preparations were strongly positive on five occasions. There were many pus cells, occasional red blood cells, and granular and hyaline

casts in the urine. Chest x-ray examination showed cardiac enlargement with diffuse pulmonary infiltration in the lower zones compatible with fibrosis or infiltration. Her fever and arthralgia failed to respond to salicylates. Prednisone 40 mg daily was started, and was increased to 80 mg daily when she became psychotic and had continued fever. It had little effect, and she died suddenly after 10 days. Necropsy showed no evidence of valvular disease or of infectious or Libman-Sachs endocarditis.

Case 2—The patient was a 16-year-old Black girl who had been admitted to hospital 13 times previously, nine of them for pneumonia. When she was 4 months old homozygous sickle-cell disease was diagnosed on haemoglobin electrophoresis at alkaline and acid pH, positive sickle test, and the presence of the sickle-cell trait in both parents. On admission she had a two-week history of jaundice, pleuritic chest pain, and swelling of both knees and ankles. She was jaundiced, had clubbing of fingers and toes, and 18-cm hepatic enlargement. The spleen was palpable. Both knees were warm and contained small effusions. Investigations showed haemoglobin 6.1 g/dl; white cell count 9.1 $\times$ 10⁹/l, 68% neutrophils; prothrombin time normal; electrolytes normal; plasma creatinine 53 μ mol/l (0.6 mg/100 ml), total bilirubin 410 μ mol/l (24 mg/100 ml) (50% conjugated), alkaline phosphatase 53 KAU. Blood and urine cultures were persistently sterile and the serum was Australia antigen negative. Creatinine clearance was 0.9 mmol (100 mg)/min, and urine protein 700 mg/24 h. Antinuclear factor (ANF) tests were consistently positive and DNA binding values (Farr assay) were 50% and 60% (normal 0-30%), C3 was 0.72 g/l (normal 1.2-1.6 g/l). Three days after admission she developed a persistent fever. Treatment with prednisone 60 mg daily was started after 22 days and the temperature fell to normal immediately. She improved clinically and her liver function tests reverted to normal. Six months later she was taking prednisone 10 mg daily, had returned to school, and was clinically well.

Discussion

In each case homozygous sickle-cell disease was diagnosed on adequate evidence. In case 1 five of the American Rheumatism Association criteria for the diagnosis of SLE were met by the presence of pleurisy, arthritis, LE cells, psychosis, and casts in the urine. In case 2 SLE was diagnosed on the arthritis, proteinuria, strongly positive ANF, high DNA binding, and hypocomplementaemia.¹ Although joint symptoms are common in sickle-cell disease these usually arise in the juxta-articular bone and are referred to the joint. The finger clubbing and hepatomegaly in both cases remain unexplained, although an association of gross hepatomegaly and clubbing in sickle-cell disease has been reported.²

The occurrence of SLE in sickle-cell anaemia is of theoretical interest because of the immunological aspects of the two diseases. Johnston *et al*³ have reported a selective abnormality of the alternative pathway of complement activation in patients with sickle-cell disease in whom the C3 and total haemolytic complement (CH₅₀) levels were normal. In studies still in progress we have confirmed this and shown that it is related to a partial deficiency in factor B (glycine-rich B protein), which is a component not only of the alternative pathway but also of the C3_b feedback amplification system.⁴ Such a defect might be partly responsible for the increased susceptibility of sickle-cell patients to infection. Immune complex diseases—particularly SLE—may be associated with genetic deficiencies of various components of complement, and complement deficiency may lead to such diseases through failure to eliminate foreign viral, bacterial, or other antigens. The defect in complement metabolism in sickle-cell anaemia may similarly lead not only to recurrent infection but possibly to immune complex disease.

¹ Hughes, G R V, *British Journal of Haematology*, 1973, **25**, 409.² Konotey-Ahulu, F I D, *Ghana Medical Journal*, 1969, **8**, 104.³ Johnston, R B, Newman, S L, and Struth, A G, *New England Journal of Medicine*, 1973, **288**, 803.⁴ Wilson, W A, Hughes, G R V, and Lachman, P J, *British Medical Journal*, 1976, **1**, 367.

Departments of Medicine and Pathology, University of the West Indies, Kingston, Jamaica

W A WILSON, MD, MRCP, consultant physician and lecturer in medicine
G D NICHOLSON, BM, MRCP, consultant physician and lecturer in medicine
G R V HUGHES, MD, MRCP, consultant physician and lecturer in medicine
(present address: Department of Rheumatology, Royal Postgraduate Medical School, London W12 0HS)

S AMIN, MB, MCPATH, consultant in pathology

GAO ALLEYNE, MD, FRCP, consultant physician and lecturer in medicine

Medical Research Council Epidemiology Unit, University of the West Indies, Kingston, Jamaica

G R SERJEANT, MD, MRCP, head of unit