CORRESPONDENCE

Sterilisation of children under 16 S D M Court, FRCP, and June K Lloyd, FRCP.....830	Exchange transfusion in prevention of iron overload J A McLean, FRACP.....834	Thrombotic complications in acute polyneuritis C P Bredin, MRCP, and others.....837
After stroke, what? B Isaacs, FRCPGLAS.....830	Carcinoma-in-situ of cervix in sisters S A Way, FRCOG.....834	Tubal pregnancy and surgery D F Hawkins, FRCOG.....837
Cross-sensitivity between practolol and other beta-blockers? Anna-Karin Furhoff, MD, and others.....831	Immunological findings in multiple sclerosis S Poser, MD, and others.....834	Rectal loss of vaginal vibrators W K Moores, FRCS; B J Mayou, FRCS; Ann G Morgan, MB.....837
Recoverable organic psychosis after hypopituitary coma S I Cohen, FRCP.....831	Failure of phenobarbitone to prevent febrile convulsions D M T Gairdner, FRCP.....834	Position of myocardial infarct and results of alprenolol treatment G Ahlmark, MD, and H Sætre.....837
Laparoscopy and laparotomy in staging of Hodgkin's disease V T DeVita, jun, MD.....831	Hazards of ergotamine tartrate J M S Pearce, FRCP.....834	Progression and regression of atherosclerosis D E FitzGerald, PHD, LRCP&SI; R Mulcahy, FRCP, and N J J Hickey, MD.....838
Medical and surgical emergencies in divers R I McCallum, FRCP, and R A F Cox, MB.....832	Ibuterol hydrochloride and terbutaline in asthma S S Poulsen, MD, and B N Petersen, MD.....835	Hazards of multilaxative mixtures E W Godding, FPS.....838
Radiology and endoscopy in acute upper gastrointestinal bleeding G M Fraser, FRCP, and D H Cummack, FFRCSI.....832	Neuroblastoma in infancy Jane V Bond, FRCP, and H H Nixon, FRCS.....835	Misdiagnosis of urinary tract infection in women M A Waugh, MB.....838
Smoking tables for carbon monoxide M A H Russell, MRCPsych.....832	Antibiotics for kicks Freda S Reed, MRCPsych.....835	Conservative Policy Group on Mental Health C E Parkinson, MP.....838
Laparoscopy explosion hazards with nitrous oxide P C Steptoe, FRCOG.....833	Alpha-chain disease cured with antibiotics O Wetter, MD.....835	Local structure of the BMA R A R Lawrence, MRGP.....839
Use of pressurised aerosols by asthmatic patients T J Coady, MB, and others.....833	C3 inactivator and hypocomplementaemia in a "healthy" woman R A Thompson, MRCPATH.....835	Lessons from the past J A Davis, FRCP.....839
Hypertriglyceridaemia and abdominal pain L A Donaldson, MB, and W McIntosh, MB.....833	Otitis media C Salkin, MB; J F Birrell, FRCSed.....836	Specialists (British style) M C T Morrison, FRCS.....839
Reticuloendothelial phagocytosis in nephritis E N Wardle, MD.....833	Detection of somatic muscle fasciculation on electrocardiograms R W Gilliatt, FRCP, and R G Willison, FRCPed.....836	Consultants and junior hospital staff contract D Kirk, FRCS.....840
Aplastic anaemia and hair dye S Hamilton, MB, and J G Sheridan, MRCPi.....834	Domiciliary oxygen in advanced chronic bronchitis P B Anderson, MRCP, and others.....836	Campaign for Independence in Medicine M S Jones, FDSRCS, and others.....840
	Carcinoembryonic antigen and mucus D L J Freed, MB, and G Taylor, MD.....836	Points from letters Psychiatric aspects of shoplifting (H B Minchom); Herpes zoster—a new diagnostic sign (S W V Davies); Vehicles for streptokinase (V F Standing); Treatment of rib fractures (S V Humphries); Specialists (British style) (G I B Da Costa).....840

Correspondents are urged to write briefly so that readers may be offered as wide a selection of letters as possible. So many are being received that the omission of some is inevitable. Letters should be signed personally by all their authors.

Sterilisation of children under 16

SIR,—The Department of Health and Social Security has recently prepared a discussion paper on sterilisation of children under 16 years of age. The document has had wide distribution and views have been sought from a large number of organisations, particularly the professions likely to be involved, together with a range of organisations concerned with the welfare of children. Because this is a subject of especial concern to paediatricians the British Paediatric Association would like to make its comments more widely known.

There are two distinct groups to be considered: the first in which sterilisation proves unavoidable in the course of treatment of a serious medical condition (therapeutic sterilisation); and the second in which sterilisation of children with severe mental handicap is undertaken in order to prevent the possibility of pregnancy (non-therapeutic sterilisation). In some of the latter cases the fetus may also be at considerable risk (for example, maternal phenylketonuria).

In the case of therapeutic sterilisation the decision must be made by the doctor in charge of the case. In an important issue of this kind it would be advisable to follow the normal practice of obtaining a second opinion but this may not always be possible—for instance,

when during the course of an operation for cancer involving the pelvic organs the surgeon finds that removal of the uterus and/or ovaries is necessary.

It was suggested in the Department's discussion paper that there may be certain genetic indications. After taking expert advice we are clear that the presence of a genetic condition alone cannot justify sterilisation under the age of 16 years.

A decision regarding non-therapeutic sterilisation should not be made by a single doctor; there are too many issues concerning the accuracy of diagnosis, prognosis, genetic implications, etc. Professional consultation should include further medical opinion, and for particular children the opinion of medical social workers, psychologists, and other health workers should be sought. It is understood that confidentiality would be observed by all involved in these consultations. Full discussion with the parents (and, where possible, explanation to the child) should, of course, precede the final decision. Although it will be necessary in all cases to consider the problem against the background of the family, including the risks to the baby who might be conceived, the interests and well-being of the child concerned (that is, the girl for whom sterilisation

is being considered) should be paramount. No parent should be able to secure a daughter's sterilisation because it is feared that the child may become promiscuous. Wherever possible decisions on sterilisation should be postponed until the age of consent. We realise, however, that in cases of severe mental handicap the individual may never be able to give "informed consent" and we would wish to see a procedure which safeguards the interests of all such persons irrespective of age.

We are not in favour of a central registry or any form of notification. Decisions should not be delegated to ethical committees. If after consultation consensus cannot be reached then a ward of court procedure should be followed.

S D M COURT
President

JUNE LLOYD
Secretary,
British Paediatric Association

London WC1

After stroke, what?

SIR,—The glow of pleasurable anticipation which was aroused when I saw this question posed in your leading article (6 March, p 541) gave way to bewilderment and dismay when I read your extraordinary conclusion that "common-sense measures [not defined] given in the early stages after the patient has completed his medical treatment (my italics) are likely to lead to as great a degree of functional

improvement as any more esoteric methods of management [not defined]."

My own non-esoteric common sense tells me that severely disabled stroke patients need and respond to intensive rehabilitation, started on the day of onset of the stroke and provided by an expert and committed team of doctor, nurse, physiotherapist, occupational therapist, speech therapist, and social worker.¹ Your reference to "physical treatment . . . unenthusiastically applied at infrequent intervals over a long period in a vain attempt to achieve unrealistic goals" certainly does not describe what happens in the all-too-few stroke rehabilitation units and in the many departments of geriatric medicine where stroke rehabilitation is taken seriously; nor do the staff of such units need to be reminded that "contractures, stiff and painful shoulders, bed sores, and loss of morale may be preventable." (And you might have added incontinence, loss of balance, muscle wasting, mental confusion, and loss of family support.)

The severely disabled stroke patient needs two or three months of intensive team work in hospital and prolonged follow-up in day hospital and community, with the active participation of his family, if he is to have a fair chance of regaining his full potential for recovery. If the acute medical units to which most such patients are admitted in the first instance are unable to provide this level of care—and it is understandable that most units should find this difficult—then it is time that collaborative arrangements were made with departments willing and able to do so.

Your article appeared one week before a heavily oversubscribed conference organised by the Chest and Heart Association on stroke clubs—an indication of how eager the community is to respond to the needs of stroke patients. You rightly quote the Royal College of Physicians' report² as having stressed that stroke causes more severe disability than any other medical condition. You might well have quoted their conclusion that "the resources available for . . . treatment and management should be better concentrated and co-ordinated in health assessment."

BERNARD ISAACS

University Department of Medicine,
Queen Elizabeth Hospital,
Birmingham

¹ Isaacs, B, et al, *The Two Stage Management of Stroke*. In press.

² The Royal College of Physicians of London, *Report of the Geriatric Committee Working Group on Strokes*. London, RCP, 1974.

Cross-sensitivity between practolol and other beta-blockers?

SIR,—During 1974 and 1975 the Swedish Adverse Drug Reaction Committee received 95 reports of adverse reactions involving the skin and eyes following treatment with practolol. The causal relationship between the condition described in the reports and practolol was classified as "probable" or "not excluded" by the committee. When the possibility of a cross-reaction between practolol and other beta-blockers started to be discussed¹ inquiries were sent to the reporting physicians asking if their patients had been treated with other beta-blocking agents. Eighty-six answers were sent in and indicated that 64 of the 86 patients had received another beta-blocker to replace practolol. All beta-

blocking agents on the Swedish market had been used, alprenolol in 26 cases, propranolol in 17, pindolol in 17, metoprolol in seven, and oxprenolol in three (six patients had received two drugs).

In six instances the symptoms thought to have been provoked by practolol either reappeared when another beta-blocker was administered (two patients) or failed to disappear in spite of the change of drug (four patients). Of the first two patients, one, a 70-year-old woman, got macular skin changes which had developed when she was taking practolol, disappeared when this drug was discontinued, but came back when alprenolol was administered one month later. The other, a man aged 56, had psoriasiform skin eruptions and complained of dry eyes. Both complaints vanished when practolol was discontinued. The eye symptoms reappeared when treatment with metoprolol was started some weeks later. Five months later a challenge test with metoprolol provoked the same eye symptoms. A test for antinuclear factor was negative. In the remaining four cases the symptoms remained in spite of a change from practolol to another beta-blocker (alprenolol, pindolol (two patients), or propranolol). Irritation of the eyes was described in three patients, one of whom also had cutaneous symptoms. The fourth patient had psoriasiform changes of the skin which did not improve when her medication was changed from practolol to pindolol.

The possibility of a cross-reaction between practolol and other beta-blockers is suggested by the first two cases, in which there was an interval with no symptoms between the periods of treatment. The chances of making even a tentative diagnosis are much less when there was a direct change from one drug to another. Four of these six patients had eye symptoms. Only one of them had undergone a specialist's examination, which had revealed no abnormalities. In three cases the causal relationship between drug(s) and suspected adverse reaction must be regarded as highly dubious, especially since Wright² has pointed out how common reduced tear flow with variable hyperaemia and vascular engorgement is in this age group.

About a year ago Felix *et al*³ reported on 32 patients with cutaneous reactions to practolol, all of whom tolerated treatment with other beta-blocking agents. Our inquiry confirms the impression that serious side effects of the type induced by practolol are uncommon with other beta-blockers, even in patients who do not tolerate practolol.

A-K FURHOF

Secretary,
Swedish Adverse Drug Reaction Committee

MARGIT NORDLANDER

CURT PETERSON

Section of Adverse Reactions,
Department of Drugs,
National Board of Health and Welfare,
Stockholm, Sweden

¹ Lyall, J R W, *British Medical Journal*, 1975, 2, 747.

² Wright, P, *British Medical Journal*, 1975, 4, 577.

³ Felix, R H, Ive, F A, and Dahl, M G C, *British Medical Journal*, 1975, 1, 626.

Recoverable organic psychosis after hypopituitary coma

SIR,—The case reported by Dr R R Parker and others (17 January, p 132) is very similar to that reported by myself and Dr Isaac Marks in 1961.¹ Our patient suffered from Addison's disease and was originally admitted to hospital in coma with a blood sugar of 1.1 mmol/l (20 mg/100 ml). She recovered from this but an organic confusional state persisted for two months and she was admitted to the Maudsley Hospital. She had been treated with steroids

by mouth and her blood sugar before admission had been 5.5 mmol/l (100 mg/100 ml) on two occasions. The steroids were continued after admission and we found when we made numerous random blood sugar estimates that it could drop as low as 2.9 mmol/l (52 mg/100 ml).

It was long ago pointed out² that symptoms of hypoglycaemia might develop at higher levels of blood sugar in patients with Addison's disease than in normals and it seems likely that the same might be true in hypopituitary patients. We thought that in our patient the blood sugar may not have been maintained at a consistently normal level for long enough for the brain to recover after a prolonged period of hypoglycaemia. We therefore gave her regular and frequent glucose drinks and by the end of a week she was quite normal and remained so during follow-up. It seems possible that because of the confusional state she had not fed herself properly and had probably not taken her medication regularly so that there was a vicious circle tending to perpetuate the hypoglycaemia and confusion. It is possible that during the confusional state she might not have been normally responsive to hunger and that once her mental state became normal she remained well because she responded normally. The recovery of the patient reported by Dr Parker and his colleagues seems to have been rather slow and I think it is possible that she might have recovered more quickly had more emphasis been placed on the correction of hypoglycaemia.

SAMUEL I COHEN

Psychiatric Department,
The London Hospital,
London E1

¹ Cohen, S I, and Marks, I M, *Journal of Neurology, Neurosurgery, and Psychiatry*, 1961, 24, 366.

² Thorn, G W, et al, *Journal of Clinical Investigation*, 1940, 19, 813.

Laparoscopy and laparotomy in staging of Hodgkin's disease

SIR,—I was pleased to see the article by Dr P Spinelli and others (6 December 1975, p 554) on this subject. Our previous work indicated that laparoscopy with multiple biopsies by needle was as likely to be useful in diagnosing Hodgkin's disease of the liver as biopsy is at time of surgery. However, I think it should be further emphasised to those who might attempt to stage patients using peritoneoscopy that splenic aspiration carries a considerable risk for the low yield. Dr Spinelli and his colleagues found that only 38% of patients ultimately found to have disease in the spleen at the time of laparotomy had identifiable disease at aspiration. This low yield is no surprise since spotty involvement of the spleen has been demonstrated in all laparotomy studies. Obviously if one relies on laparoscopy to sample the spleen a significant number of false-negative results will be achieved. There were two episodes of bleeding requiring surgical intervention which, in my view, is an unacceptably high complication rate.

The main value of laparoscopy in staging is the identification of disease in the liver. If disease is demonstrated this may change the entire approach to the therapy of the patient regardless of what is ultimately found in the spleen. If disease is not demonstrated in the liver and the patient is already known to have stage III disease the spleen, if it is of normal size, can be encompassed in a radiation field