that sugar raises the plasma concentration of cholesterol in man. The report refers to the "very low carbohydrate diets" commonly used for the treatment of obesity and says that they cause an increase in plasma cholesterol. The commonly used diet is not very low in carbohydrate—it contains something like 60 g a day—and it certainly does not increase plasma cholesterol.⁵

The main failing of the report is its dedication to the hypothesis that the concentration of blood cholesterol is the dominant feature of CHD. The fact is that there are many other features; among these are a raised concentration of triglyceride and uric acid and a diminished glucose tolerance. A search for the causes of CHD should seek factors that produce as many as possible of these features. Alternatively, it should seek hormonal changes that almost certainly underlie this range of abnormalities that characterise the disease; how else do we explain the considerable difference in disease prevalence between men and premenopausal women? So far an acceptable candidate is insulin; it may, however, well be that the primary disturbance is in some other hormone.^{6,7}

JOHN YUDKIN

London NW3

¹ Joint Working Party, *Journal of the Royal College of Physicians of London*, 1976, 10, 213.

² *The Times*, 1 May 1976.

³ Department of Health and Social Security, *Diet and Coronary Heart Disease*. Reports on Health and Social Subjects, No 7. London, HMSO, 1974.

⁴ *Mass Field Trials of the Diet-Heart Question*, American Heart Association Monograph No 28, 1969.

⁵ Evans, E, Stock, A L, and Yudkin, J, *Nutrition and Metabolism*, 1974, 17, 360.

⁶ Yudkin, J, *Acta Diabetologica Latina*, 1972, 9, suppl 1, p 395.

⁷ Yudkin, J, in *Controversy in Internal Medicine II*, ed F J Ingelfinger et al. Philadelphia, Saunders, 1974.

Myocardial infarction

SIR,—Dr H G Mather and others (17 April, p 925) have made inadequate conclusions from their own study. Although they quote Christiansen *et al*¹ as showing that the introduction of coronary care units has been associated with a fall in the hospital mortality rate their own figures demonstrate that the mortality rates for patients treated at home do not for the most part differ significantly from those for patients treated initially in coronary care units in hospital. Not only was it confirmed that some patients, especially those who are initially hypotensive or who have a history of angina or of myocardial infarction, have a poor prognosis wherever they are treated, but also it is shown that other patients, especially those over 60 years of age, may do better if treated at home. What is certainly not demonstrated is that there is a place at all for coronary care units. The recently published joint report on coronary care by the Royal College of Physicians and the British Cardiac Society² is more courageous in its conclusion in this respect.

It might be thought unethical to include in a controlled study a group of patients for whom the medical profession did nothing whatever in hospital or in general practice. None the less, I venture to suggest that one of the conclusions that may possibly be drawn from this paper is that the influence of the medical profession is questionable, and moreover the paper may even reinforce Illich's view that not only are coronary care units of doubtful advantage³ but

in some cases there may be a direct correlation between the amount of professional medical activity and the increased morbidity of the patient. This surely is the true conclusion to be drawn from the figures that demonstrate that no group of patients does specifically better in a coronary care unit and one group may indeed do significantly worse.

The opening statement of the paper, "Because of the increase in the number of deaths from coronary heart disease, improvement in the treatment of this condition is a matter of urgent concern," is fundamentally wrong.

The report of the Joint Working Party² summarising current knowledge and research work concluded that 40% of coronary deaths occur within the first month, the majority of those within the first hour, and the majority of those within the first few minutes. Two-thirds of all fatal attacks are medically unattended. The disease now kills 35 000 men and women of under retirement age every year in the UK. My belief is that there is no practical system of postcoronary care that could possibly influence these figures to any significant degree. There is no possible framework of clinical organisation that could cope with these numbers at the right time and in the right place even if the treatment was effective. Prevention, not treatment, is what should be the matter for urgent concern. To that event the authors' suggestions on the management of acute myocardial infarction are an irrelevance in the wider sphere of the management of ischaemic heart disease.

ROBERT LEFEVER

London SW7

¹ Christiansen, I, Iverson, K, and Skouby, A P, *Acta Medica Scandinavica*, 1971, 189, 285.

² Joint Working Party, *Journal of the Royal College of Physicians of London*, 1975, 10, 5.

³ Ivan Illich, *Medical Nemesis*, p 51. London, 1974.

Regression of atherosclerosis

SIR,—There has been much interest lately in the possibility of causing regression of atheromatous plaques by giving polyunsaturated fats by mouth. I should like to suggest that, of all sites in all arteries, it is the progress of atherosclerotic plaques in cerebral arteries that can be assessed most effectively because the reduction of cerebral blood flow caused thereby—and the increase of cerebral blood flow when plaques regress—can be monitored readily by the resulting effect on behaviour, which changes in certain recognisable ways that are observable and recordable.

Some 14 years ago I conducted an investigation along these lines,¹ using large doses of corn oil emulsion combined with a reduction of dietary animal fats, and found indeed that behaviour in atherosclerotic demented (though not in senile demented) improved, in particular in respect of incontinence. Since then I have been using corn oil emulsion in other cases of atherosclerotic dementia, enabling many patients to leave hospital and to lead relatively normal lives in the community again.

H DALE BECKETT

Cane Hill Hospital,
Coulston, Surrey

¹ Beckett, H D, *Journal of Mental Science*, 1962, 108, 453.

Sociological realities

SIR,—It is interesting to see from your leading article (3 April, p 790) that you are a disciple of Chairman Mao. Your conversion is probably not total, for I doubt whether you want to promote the notion of the "barefoot doctor"; but the barefoot social worker is obviously your ideal, and positively Mao-like is your scorn for the pedagogues who seek to keep the student social worker in the classroom rather than the rice-fields, or at any rate their urban equivalent.

All the internal evidence suggests that you know little at first hand about social work and nothing about social work education. Ignorance, however, is venial; arrogance is not. So powerful is the unexamined assumption which dominates the article, that social workers are useful pairs of hands with minimal educational needs, that the uninformed reader is bound to emerge with wholly misleading notions of the purpose and arguments of the document supposedly under review. The CCETSW paper is not faultless: in common with most global reviews of educational policy it has its pretentious moments. But its goal is a useful one and its concerns are essentially practical. Who would suppose, having read the *BMJ* account, that the main emphasis of the paper in question was precisely on the need to give priority to training for practice skills rather than to academic knowledge?

In fact, neither the authors of the report nor you need feel unduly alarmed at the prospect of social work students waddling out of their training courses like so many Strasbourg geese, stuffed to bursting point with arcane theories. The exigencies of time-tableing alone impose a more varied and less heady diet. My own department, which is typical of many, takes young (and some not so young) graduates and completes their professional training in either one or two years, according to the type of first degree course they have followed. At least half of that time must be spent in learning practical skills, under supervision, in local authority departments, hospitals, health centres, or voluntary organisations. A student whose first degree was in Russian or zoology has therefore only the equivalent of one year in which to acquire the necessary "knowledge base" for social work practice, the social sciences graduate only half a year. Uncertainty about the essential components of the "knowledge base" does not imply, as you imagine, doubts as to what a social worker's job is but rather a continuing attempt to define those specific areas of knowledge most relevant to the job. Inevitably this means stringent selection and the elimination of a very great deal which, though of considerable interest, is not of central relevance to the professional task.

For you, of course, there are few problems. You know what a straightforward task the social worker has and how useless it is to do very much more than throw him into the world of practice. Having spent more than a decade on the staff of a medical school, I can imagine the outcry that would (quite properly) go up if I were sufficiently philistine to argue in your columns that it is an absurd waste of time to fill the heads of medical students with anatomy, physiology, biochemistry, and pharmacology instead of concentrating on practicalities. I would rapidly be reminded that a professional man is distinguished from a technician, inter alia, by the fact that he understands the theoretical principles underlying the practical measures he takes. You would no doubt deny