While rarely heard of nowadays except in Dr Apley's metaphorical sense, a tooth-comb was in the past an essential weapon in the battle against head-lice. Perhaps its apparently superfluous prefix was suggested by the pain caused by its being dragged across the scalp and through tangled hair in the search for the elusive nit.—Ep, *BMJ*.

In defence of the under-40s

SIR,—Dr P Buisseret (1 May, p 1075) has many strictures for my recent Personal View article—so many that I wonder he bothered to read it “two or three times” and to tot up its grammatical errors, tautologies, and clichés (none of them specified but some patently intentional, by the way). Despite his assertion that I could not be taken seriously he seems to have taken me even more seriously than I take myself: I never count my clichés. They seem, however, to have hatched, for Dr Buisseret has clearly got much of the message I meant to convey.

Pace Dr Buisseret, I did not invent “jugglery”: it is, according to the *Shorter Oxford Dictionary*, good Middle English. If he really knows “acceptable and more subtle English counterparts” for “angst” and “manqué,” not to say for my neologism “medico-bureaucrat,” perhaps he will let us have them (though I'd prefer simple English synonyms, neither more nor less subtle than the originals). His statement that I do not understand the meanings of “apposite” and “sacerdotal” is, of course, impossible to refute, although if I do not the editors of the *Shorter OED* are in like case, for the meanings I have in my head for those words are, I discover, the meanings they provide. It is, however, certainly possible to contest his statement that my article contained “seven words or phrases which convey no meaning” for I know of no word that does not convey a meaning; and perhaps therefore he will tell us which words he had in mind.

May the kettle call the pot black? Dr Buisseret writes that “since you had evidently seen fit to publish such nonsense, you should receive an appropriate protest.” A plain “since you published such nonsense” would have avoided a double tautology. And does he often go in for inappropriate protests? He says too that I made “statements regarding the social and intellectual behaviour of baboons, subjects which have recently been thoroughly and lucidly documented and found to be of a remarkably advanced kind.” Are the subjects remarkably advanced, or is it the behaviour that is? And if the latter, may we have details of this documentation of the “intellectual behaviour” of baboons?—not, by the way, that I made any reference to their social or intellectual behaviour, with which I cannot claim acquaintance.

None of the doctors who did me the kindness of writing to me direct about the article had anything but praise for it; and they were split equally as to the age-40 barrier. My last and much my most important point is that I did in a sense, as Dr Buisseret suggests, write the article after gazing into a mirror. I wrote it after holding a mirror up to the whole profession, both the under-40s and the over-40s; and he seems not to have noticed that my criticisms—as distinct from statements of fact or of purported fact—were reserved almost entirely for the second group, among which, I

regret to say, I am certainly numbered. It was not the doctors of Dr Buisseret's generation I was accusing (“Who can blame them?” I wrote), but mostly those of my own, myself included, for having failed his.

JOHN S BRADSHAW

How Caple,
Hereford

“Automatic” sun-glasses: a warning

SIR,—Several newspapers are carrying advertisements for special sun-glasses, which are recommended also as suitable for driving. The novelty consists of photochromic glasses—referred to as “automatic light filters”—and, on the face of them, they appear to have homeostatic properties. The glasses darken on exposure to light and vice versa. In their present form they may constitute a driving hazard, and I was told at the recent Northern Optical Congress in Harrogate that one of the principal manufacturers has no plan at present to deal with this aspect. Published data—for example, for Reactolite glass—show that, irrespective of the tint, the time in which half the total darkening is completed is of the order of one minute. The time for the reverse process—namely, half the recovery of the maximum transmission—can be up to half an hour or more.

The advertisements contain no warning to the effect that such glasses can be a grave risk; in fact one of them states, “You can wear them all the time.” The glasses are, however, a hazard even in daytime, as anyone driving through the Kingsway or any other similar tunnel or underpass will discover with ease and, on Italian motorways, with alarm.

R A WEALE

Institute of Ophthalmology,
London WC1

Skin contact pressures

SIR,—The article by Professor P H Fentem and others (31 January, p 254) contained a timely discussion of pressure measurement techniques. Skin contact pressures are important in “tissue viability” research, the emphasis of which is on decubitus pressure problems, estimated to affect 25 000 patients and to cost Britain £60 million annually.¹ Skin contact pressures may be helpful in fitting surgical appliances, prostheses, splints, bandages, and pressure garments other than stockings. A review of the literature² shows that skin contact pressure research is a growing area, with increasing numbers of workers using their own methodologies. Discrepancies in results are becoming apparent, and may be attributed to factors such as sensor size and thickness as well as the topography and consistency of the site to which the sensor is applied.

Apart from a minority of methods which include the “extraordinary” and the “blow by” the commonly used are of two kinds: (1) Space-occupying fluid-filled “fixed volume” bladders often connected via a transducer to a system capable of continuous recording; and (2) the empty bladder kind usually of the “electro-pneumatic” type. Both kinds may have large or small and thick or thin sensors. Comparison of the 8 × 4 cm square Scimedics sensor developed for cushion prescription³ and

the 28 mm diameter sensor used by Isherwood *et al*⁴ for assessing stump bandaging showed that for nearly flat interfaces readings were identical. The 28 mm sensor was used on a series of hard cylinders of decreasing circumference. If the sensor was taped in apposition, and a baseline reading subtracted from the “test” reading, a correction for sensor distortion could be made. However, the limit of the sensor was reached on a 3 in (8 cm) circumference.

We believe that the field of skin pressure research has a requirement for three sizes of sensor—say, about 8, 3, and 1 cm in diameter, whether or not these are capable of “spot” measurements or continuous recording. We would like to see more work comparing different techniques so that perhaps agreed methods can be used in the future. Would a society such as the Biological Engineering Society take on the promotion of standardisation?

P A ISHERWOOD

A ROSSI

London

J E and J E DRUETT

J C ROBERTSON

Salisbury

¹ Fernie, G R, PhD thesis, University of Strathclyde, Glasgow, 1973.

² In preparation.

³ Reswick, J B, and Rogers, J, Strathclyde Symposium on Tissue Viability.

⁴ Isherwood, P A, Robertson, J C, and Rossi, A, *British Journal of Surgery*, 1975, **62**, 982.

Low-dosage treatment with tetrabenazine

SIR,—The drug tetrabenazine was introduced for the management of Huntington's chorea and allied disorders associated with choreiform movements in 1971. McLellan *et al*¹ recommended it as the drug of choice in such disorders, but two of their 10 patients developed Parkinsonian symptoms. Astin *et al*² used it in 26 such patients, but three had to stop taking it because of side effects. The following case report might help to extend the use of this drug.

A mentally retarded woman of 52 was seen as an outpatient and diagnosed as suffering from congenital choreoathetosis. She was referred back to her general practitioner with the advice to start her on tetrabenazine 25 mg daily and to increase the dose until an optimal response had been obtained. When she was seen six months later her choreiform movements had completely disappeared but she showed typical Parkinsonian symptoms. These disappeared on withdrawal of the drug, but the choreiform movements reappeared. Restarting treatment with the smallest tablet available, 25 mg, twice a day again precipitated Parkinsonism. She was eventually stabilised satisfactorily on a dose of approximately 6.25 mg twice to three times a day.

We should like to make a plea to the makers for a smaller tablet in order to make the management of such patients easier.

R SCHNEIDER

A P BUCHAN

T A COUSTON

St Margaret's Hospital,
Birmingham

¹ McLellan, D L, Chalmers, R J, and Johnson, R H, *Lancet*, 1974, **1**, 104.

² Astin, K J, and Jumpert, E W J, *Lancet*, 1974, **1**, 512.