was advised not to "cover" a "specialist midwife" who was proposing to insert IUCDs as part of the domiciliary service provided by the local branch of the Family Planning Association. I presume the same advice would still be given. I do not think that nurses should be allowed to insert IUCDs as the complications which inevitably arise might bring this valuable method of contraception into disrepute.

T G E WHITE

Croydon, Surrey

Antibiotics again

SIR,—Your leading article (8 May, p 1107) disapproves of the fact "that chloramphenicol continues to be prescribed for illnesses other than typhoid fever." If this remark discourages use of chloramphenicol in haemophilus meningitis and other life-threatening infections due to *Haemophilus influenzae* Type b it could have serious consequences.

In their 1967 leaflet about chloramphenicol the Committee on Safety of Drugs stated: "It is a highly effective drug in typhoid fever and in *H. influenzae* meningitis. In these conditions its advantages outweigh its dangers." Since then the case for using ampicillin (the only serious alternative to chloramphenicol) for treatment of haemophilus meningitis has been weakened by reports of its failure to eradicate the infection in some cases despite normal ampicillin sensitivity of the organism¹; of more rapid resolution of signs of infection when chloramphenicol is used²; and of cases of meningitis due to β -lactamase-producing (and therefore ampicillin-resistant) strains of *H. influenzae*.³

I hope to publish soon laboratory data supporting the suggestion that chloramphenicol is more effective than ampicillin against *H. influenzae* Type b in circumstances which may well be relevant to conditions in the meninges in haemophilus meningitis. Available information suggests that children with haemophilus meningitis and similar severe haemophilus infections who are given ampicillin rather than chloramphenicol may not improve as rapidly as they should, may relapse, or may (if they are unfortunate enough to have an ampicillin-resistant haemophilus) die from lack of effective treatment. In these circumstances I must sharply disagree with the reprimand implied in your leading article.

D C TURK

Bacteriology Department,
Gibson Laboratories, Radcliffe Infirmary,
Oxford

¹ Kandall, S R, Davis, T C, and Abramowicz, M, *Clinical Pediatrics*, 1972, **11**, 264.

² Shackelford, P G, et al, *New England Journal of Medicine*, 1972, **287**, 634.

³ Nelson, J D, *Journal of the American Medical Association*, 1974, **229**, 322.

Devolution

SIR,—As a prospective Plaid Cymru parliamentary candidate I found that the transcript of your conference on devolution (8 May, p 1127) brought out some very interesting points.

Relating specifically to medicine, there are two points I would like to emphasise. Firstly, the medical service in Wales, in common with many parts of England, is grossly over-

dependent on medical staff from other countries. Any suggestion that we could not plan our health service better than at present is quite laughable when one considers that our own doctors are frequently in a minority in our own hospitals. Such a situation would be quite unacceptable in any other country except possibly in the Third World, and then it would undoubtedly only be on a temporary basis. I have greater faith in the ability of the medical world in Wales to plan for this relatively small country in a more efficient manner and with greater foresight than the Elephant and Castle has done for the UK in the past and to develop such a service without constant political interference.

The other point I would wish to lay before the doubters of devolution is the important one of pay differentials arising in a devolved UK. My only comment here is that I should hope they would. Ironically, in *General Practitioner* the same week as the seminar it was shown that general practitioners in the UK are now the poorest in Europe. Even Ireland is apparently now in a position to offer medical staff a better deal than in the UK.

Things could hardly be worse in the Health Service than they are at present. There is a lack of confidence which I have not experienced previously and which I believe will be righted only as and when we have devolution and a system of government which is readily answerable to the people it serves. A self-governing Wales with its rightful priorities would indeed not only benefit Wales but help England to find its true role as a partner in Europe. A recent decision of the London Government, for example, to invest in 600 new multirole combat aircraft, despite our economic problems, at an average cost of £5m each smacks of imperialistic grandiosity. That age is gone if for no other reason than that the UK cannot afford such priorities, and I am convinced that this is the opinion of the vast majority of the English people too.

In the Western developed world many of our social problems arise through a lack of identity and rootlessness. In Wales particularly there is a very rich tradition of community. This can be enhanced only by relating this tradition to the ideal of full nationhood status for our country.

Of one thing we can be certain: the next five years will be very exciting ones for our respective countries.

CARL IWAN CLOWES

Llanaelhaearn,
Gwynedd

SIR,—Your discussion of devolution (8 May, p 1127) contained data on the under-financing and under-staffing of English hospitals which look even more remarkable on recalculation per caput.

	England	Wales	Scotland	Excess, Scotland v England
<i>£/head population 1973-4 and 1975-6</i>				
Central administration	0.45	0.41	0.44	-2%
General medical services	4.05	4.1	4.4	+9%
Personal social services	10.2	8.9	7.6	-25%
Hospital services	36.6	36.4	46.9	+28%
Total health services budget 1973-4 ..	65.3	56.4	65.6	+1%
Estimated health services budget 1975-6 ..	79	82	97	+23%
<i>Health Services Staff per million population</i>				
Hospital doctors	551	521	848	+54%
GPs (including assistants and trainees) ..	485	490	568	+17%
Community physicians	28	98	59	+110%
Hospital nursing and midwifery	6869	6385	8483	+23%

Per head of population Scotland in 1973-4 compared with England had 17% more family doctors, 23% more nurses and midwives, and 54% more hospital doctors, and this year each Scot is to be allowed 23% more money on total health services than each Englishman.

J H BARON

London NW8

Oestrogens as a cause of endometrial carcinoma

SIR,—We have read your leading article (3 April, p 791) and related correspondence with considerable interest, for, like many other gynaecologists who were initially conservative towards oestrogen replacement therapy, we have come to appreciate the benefits that can be afforded to patients with climacteric and post-menopausal disturbances and have become "moderates," supporting some symptomatic therapy. Like other clinicians we have also been perturbed by the reports^{1,2} from America suggesting that such therapy might be associated with an increased risk of endometrial carcinoma and have noted the criticisms of these studies by Mr J W W Studd (8 May, p 1144). His comments relating to the retrospective nature of these reports and the need for an independent review of the relevant histology seem particularly apposite. With regard to inadequate socioeconomic matching being a confounding factor, however, in the series of Ziel and Finkle² patients and controls were matched for areas of residence and one wonders therefore whether such socioeconomic groupings in this report were that unbalanced.

We note the reservations in your leading article on the value of administering progestational compounds with oestrogen therapy in hormone replacement regimens, but the results of treatment of persistent endometrial hyperplasia with progestogens by such authorities as Wentz,³ Kistner,^{4,5} and Taylor⁶ and their effectiveness in causing regression of an endometrial state that many authorities believe antedates uterine carcinoma make the concomitant use of progestogens a very pertinent consideration. The observations of Dr E Schleyer-Saunders (8 May, p 1145) give further support to the adoption of an oestrogen-progestogen regimen and a sequential form of replacement therapy has been commonly favoured by Mr Studd⁷ and other authorities.

The need for continued vigilance even with an oestrogen-progestogen mode of treatment nevertheless is emphasised by a small number of cases of endometrial carcinoma reported in patients taking oral contraceptives of both the sequential and combined forms.^{8,9} It has been postulated that such patients may have localised "islands" of endometrium refractory to