bonding, but I wonder how many of us could get to the cure in time.

These adhesives should be handled with a non-touch technique, but as the general public know little about this perhaps it would be safer were these modern adhesives restricted to use by technical staff until a general awareness evolves.

W G STRAWBRIDGE

Pontypridd,
Glamorgan

Alternatives to levodopa

SIR,—I am surprised that in your leading article (15 May, p 1169) you are luke-warm about combined therapy for Parkinsonism. You refer to a relatively enthusiastic article by Professor Marsden,¹ but conclude that "nausea and slow initial response can be overcome by combining levodopa with a peripheral decarboxylase inhibitor, but this has few other real advantages." In the elderly patients that I have studied (50 patients of average age 76 years) combined therapy has several advantages over levodopa. The speed with which dosage can be increased means that patients spend less time in hospital or can be treated as day patients. The relative lack of nausea means that more patients can achieve a therapeutic dosage (about 500 mg a day of levodopa with 50 mg of carbidopa in my group). In addition, the success rate measured by improvement in functional ability is higher. The relative lack of cardiac side effects is valuable in an age group in which multiple pathology is common. The limiting factors in the elderly appear to be the mental side effects of confusion, hallucinations, and sleep disturbance. Involuntary movements are not a problem on the low dosage recommended.

I agree with Professor Marsden's conclusion in a later article² that combined therapy is the present treatment of choice in Parkinsonism.

M HILDICK-SMITH

Department of Geriatric Medicine,
Nunnery Fields Hospital,
Canterbury, Kent

¹ Marsden, C D, *et al*, *Journal of Neurology, Neurosurgery and Psychiatry*, 1973, **36**, 10.

² Marsden, C D, *et al*, *Lancet*, 1973, **2**, 1459.

Future of community medicine

SIR,—National Health Service Reorganisation was viewed with considerable trepidation by many of the older medical officers of health and with qualified optimism by some of the younger members of the specialty of preventive medicine. It is now clear that the older members of the specialty were right. Preventive medicine, as practised by doctors, no longer exists in Britain because it cannot compete successfully with the curative services for a share of the single purse from which the hospital and community health services are now provided. In the past, of course, preventive medicine was funded separately, mainly by the local authorities.

The new specialty of community medicine is merely medical statistics with some administration and is increasingly concerned with planning services for patients who are the victims of the lack of preventive medicine. At district level we are medical statisticians without the statistical staff or the time to

enable us to produce the detailed and accurate information required by the district management teams and without the power to put into effect the few ideas we may have time to produce.

The new circular HC(76)18 on joint care planning weakens the district management team further by making strategic and service planning a function of the new joint care planning team at area level and leaving the health care planning teams (now to be reconstituted as district planning teams) to carry out operational planning within the policy laid down by the joint care planning team, which incidentally has a large local authority input.

I will not take up your time with all the other problems of community medicine which make many community physicians feel that the specialty may not survive after the retirement of its present members. Perhaps this was one of the undeclared aims of reorganisation.

L M MAYER-JONES

Durham Health District,
Dryburn Hospital,
Durham

Operation baby lift

SIR,—In a leading article (17 January, p 112) you suggested that the British Paediatric Association might take the lead in planning for the reception of large numbers of infants transported by air from a disaster area. The problem was therefore carefully considered by the Joint Accident Committee of the British Paediatric Association and British Association of Paediatric Surgeons at a recent meeting.

In the article it was suggested that there should be, among other things, a store of paediatric supplies and lists of accommodation and staff (both professional and volunteer) for the reception of up to 1000 babies, and that all large cities with a major airport should make plans along these lines. However, much time and money would be needed to establish the organisation necessary for this aid and the lists would quickly get out of date without continuous review. The chances of any one major city having to receive a consignment of babies without reasonable warning must be so remote that the committee felt that such a major exercise, with the inevitable expense, would be unjustified.

R H JACKSON

Children's Department,
Royal Victoria Infirmary,
Newcastle upon Tyne

Convener,
BPA/BAPS Accident Committee

Quality of life

SIR,—I read with interest your leading article concerning the quality of life for the elderly in long-stay wards and old people's homes (15 May, p 1168). Two points are raised—one is the quality of life that residents in these institutions may have and the second is the great need for appropriate allocation of scarce resources.

Many people concerned with the care of the elderly feel the need for urgent reappraisal of the type of accommodation we provide. While there may be differences in the amount of

nursing and other care that patients in different forms of institutions may require, fundamentally all such persons need a home-like environment and it is with this in mind that social services departments and hospital authorities should consider future joint planning. At the same time, of course, we must not revert to the deficiencies of the old "Poor Law" institutions.

At a time of extreme financial stringency appropriate use of scarce resources must be a firm aim. Early referral to a social worker and geriatrician is clearly an essential part of assessment of an elderly person for an old people's home, as it may be that alternative action is more appropriate. It will also avoid persons in need of hospital care being placed wrongly in welfare homes with the consequences that your article describes. However, on the other side of the fence one of the greatest problems at present is to utilise sheltered housing for the needs for which it was originally intended. There is certainly one London borough where it is difficult to house people in sheltered housing who require walking aids, as the housing department finds it impossible to recruit wardens for these developments if the residents are disabled. It is urgently necessary to reconsider the qualifications that are needed to fill a warden's post, and it may be that a period of training, with experience of minor nursing procedures, may be valuable. At present many elderly people who could benefit from sheltered housing are being maintained in totally unsatisfactory housing conditions for this reason.

More geriatric beds are clearly needed in some areas, but, as the recent consultative document¹ shows, there will not be a dramatic increase. Geriatricians have to distribute their scarce resources to as many as possible. Inter-mittent and holiday admissions go a long way to solving this, allowing the person to feel wanted by the family while at the same time relieving them of some of the burden of caring for an elderly relative. Early referral is essential and is best undertaken at an outpatient clinic or at a clinic for the elderly in a general practitioner health centre.

A C D CAYLEY

Geriatric Department,
Central Middlesex Hospital,
London NW10

¹ Department of Health and Social Security, *Priorities for Health and Personal Social Services in England: A Consultative Document*. London, HMSO, 1976.

Neuropathy or myelopathy?

SIR,—In their interesting paper on "folate-responsive neuropathy" Drs M Manzoor and J Runcie (15 May, p 1176), by not using conventional nomenclature, may have added confusion to what they admit is a controversial subject.

Neuropathy is commonly taken to mean peripheral neuropathy, and bilateral extensor plantar responses would not be "characteristic." The authors appear to indicate that their patients had spinal cord disease, and "folate-responsive myelopathy" would have been more accurate and less misleading. If their patients also had a peripheral neuropathy (thus emphasising a similarity to vitamin B₁₂ deficiency) the use of well-established neurophysiological techniques might have given valuable information about peripheral nerve function in these patients before and after treatment. An opportunity has been missed to show that folic acid may have a role in