of the baby are impossible to carry out. Nevertheless, all the indications are that the practice is nearly always, in hot countries or cold, unnecessary and undesirable. Newborn babies often need frequent body contact and always require abundant opportunities to suckle. Their mothers, no less, need support and encouragement, as is recommended by the authors of the study.

MICHAEL GURNEY

World Health Organisation,
Regional Office for the
Eastern Mediterranean,
Alexandria, Egypt

¹ Laupus, W E, in *Nelson Textbook of Pediatrics*, ed V C Vaughan and R J McKay. 10th edn, p 164. Philadelphia, Saunders, 1975.

² Newton, M, *American Journal of Clinical Nutrition*, 1971, 24, 987.

³ Applebaum, R M, *Pediatric Clinics of North America*, 1970, 17, 203.

Encouraging breast-feeding

SIR,—The recent papers by Dr E Eastham and others (7 February, p 305) and Drs C J Bacon and J M Wylie (p 308) have focused attention on the problem of breast-feeding, and attempts have been made to determine what influences a mother's decision to breast-feed her baby.

At Fazakerley Maternity Hospital in Liverpool (approximately 3500 deliveries a year), where many of the patients are from the lower socio-economic groups, the proportion of mothers breast-feeding had fallen to 10% in October 1975. Efforts to remedy this included a short questionnaire, designed to be answered in a "yes/no" manner, given randomly to women who had just been delivered. Only 65 usable replies were received, making detailed analysis impossible; nevertheless, a broad area of ignorance was highlighted.

The results obtained mirror those from more exhaustive surveys. When they came into hospital only 13 of the 65 mothers hoped to breast-feed their babies. Thirty-three would have liked more advice and plainly this had not been available to them from medical sources in an understandable form, if at all. Only one patient had received any instruction from her general practitioner, other sources being the antenatal clinic (6) parentcraft class (7), relaxation class (10), the patient's mother or another relative (26), or a women's magazine (10). Of the 13 patients hoping to breast-feed, 12 wanted more advice; 21 others intending to feed their babies artificially were potential "converts" had they received more advice. The need to return to work, often assumed to be a significant factor, seemed to be irrelevant, as only three mothers stated that they would be returning to a full-time paid job within three months.

Patently a great deal more could be done in the antenatal period. The opportunity to attend parentcraft classes is open to all patients, but the minority who do so seem to represent disproportionately the higher socioeconomic groups, many of whom already intend to breast-feed. At Fazakerley Hospital, finances permitting it is intended to have a continuous running audiovisual projection unit in the clinic waiting area providing breast-feeding propaganda, but at best this can only give limited information, though it is hoped that it will stimulate interest.

In a utopian obstetric service infant feeding would be discussed individually with every patient early in the pregnancy and advice would always be readily available; but in a busy clinic in reality there is not the time available. Perhaps there is a role for the paediatrician in the antenatal period at least to interview every primipara.

Fundamentally, however, the problem is one of changing irrational social attitudes to breast-feeding. Should not those concerned with infant welfare be going to the secondary schools and explaining sensible feeding practices to the potential parents (mothers and fathers) of a few years hence?

J G DAVIES

Alder Hey Children's Hospital,
Liverpool

Bottle-feeding

SIR,—Drs B A Wharton and H M Berger (29 May, p 1326) begin their article on bottle-feeding: "Whenever possible mothers should be encouraged to breast-feed their babies—even as little as two weeks is valuable." That pathetic "two weeks" aside, when is it not possible to encourage breast-feeding? Did they not mean "When there is no contra-indication"? And how often is there a contra-indication, as distinct from a culturally determined unwillingness to breast-feed, a conditioning doctors and nurses at worst take part in and at best commonly ignore? There is no mention in the article of the effect of the practice of some maternity hospitals of supplying mothers with preformulated feeds. And neither in this article nor in the preceding article by Dr D P Addy (22 May, p 1268) was there any detailed discussion of how to induce mothers to breast-feed, and this despite the many major advantages, some of them lifelong, detailed in Dr Addy's article.

The bottle-feeding article was obviously well-intentioned and most accurate; but it would need another article to comment on it adequately. I will make only four points. Firstly, sterilisation of bottles and teats by boiling: the description of this is daunting indeed. No wonder diarrhoea in bottle-fed babies is common, and in underdeveloped countries often fatal.

Secondly, making up the feed: although Drs Wharton and Berger do commend the 1974 Report of the DHSS Working Party,¹ they do not mention the fact, which seven expert members of the working party acknowledged² some months after the report appeared, that they themselves had used for the analyses quoted in their tables in the report a 1 in 8 dilution of National Dried Milk as against the 1 in 10 to 1 in 12 recommended on the NDM pack. Their report stated, as regards the making-up of feeds, "it is not only mothers who make mistakes. . . . Nurses and midwives as well as mothers were inaccurate, and all erred usually on the side of over-concentration." They could by their own admission have added "and the country's leading experts on baby feeding." In this context the working party's indication that NDM had for years contained undesirable excess of sodium and phosphate was almost an irrelevant detail.

Thirdly, Drs Wharton and Berger state that "infant formulae based on unmodified cows' milk with added vitamins and iron were used for 30 years or more before the dangers of too much vitamin D or too much sodium were fully realised." With that background and the defects of NDM as a *donnée*, might it not be advisable to proceed most cautiously with any further modification of artificial feeds? And yet we read a little later, in connection with the missing anti-infective properties of such feeds, that "possibly by genetic selection a cows' milk containing larger amounts of lactoferrin might be developed" and "experiments are already being made to produce cows' milk rich in immunoglobulins by immunising cows with appropriate organisms." As we currently have some 600 000 births a year and so could have available a plethora of human milk with all its highly beneficial qualities, why bother immunising the cows? And are they not to be given a feed with a formaldehyde-casein coating to increase the

polyunsaturated fat content of the milk, and so bring it even more into line with human milk?³

Fourthly, I found no mention anywhere in the article of the practice in underdeveloped countries, encouraged by the West, of giving up breast-feeding in favour of bottle-feeding; and yet the *BMJ* is read widely in such countries. A recent official estimate⁴ for the number of children in the world suffering from malnutrition due to the decline in breast-feeding was 200m (*sic*). I think of bottle sterilisation and scoops and of an illiterate African woman with no running water and an open fire paying one-third of her husband's wage for the milk powder, which she therefore over-dilutes with the obvious consequences; and of a Western baby, obese from his over-concentrated feeds.

The article by Drs Wharton and Berger, and indeed our whole attitude to breast- and bottle-feeding, remind me of what was said very recently about obstetric delivery today: "Experience has shown that labour is far from safe if left to the capricious whims of Nature."⁵ Substitute "infant feeding" for "labour" and the sentence seems to express exactly the sentiments of much of the Western medical establishment towards breast-feeding. If the human body cannot teach us the dangers of interfering overmuch with the capricious whims of Nature, surely what we have done in the last 30 years to the ecosphere might. But then we doctors often wear blinkers. It was not by chance that it was a layman⁶ who uttered recently what has for some time been a truism: that "The medical establishment has become a major threat to health."

JOHN S BRADSHAW

How Caple,
Hereford

¹ Department of Health and Social Security, *Present Day Practice in Infant Feeding*, Reports on Health and Social Subjects No 9. London, HMSO, 1974.

² Arneil, G C, et al, *Lancet*, 1975, 1, 450.

³ *Lancet*, 1975, 2, 960.

⁴ Food and Agriculture Organization of the United Nations: *The State of Food and Agriculture 1974*. Rome, FAO, 1975.

⁵ Beard, R W, and Chamberlain, G, *Lancet*, 1976, 1, 904.

⁶ Illich, I, *Limits to Medicine*, p. 3. London, Marion Boyars, 1976.

Alcohol and the brain

SIR,—In your leading article on this subject (15 May, p 1168) you were kind enough to quote the air encephalographic, psychometric, and electroencephalographic study of alcoholic cerebral atrophy written by Dr Lance Perrett and myself.¹ I would like to add a few comments in the light of recent developments in the diagnosis of brain disorders.

Our own study relied on relatively simple psychometric tests such as the Wechsler Adult Intelligence Scale and the Benton Visual Retention Test. Even so there was a significant correlation ($P < 0.01$) between these tests and the degree of cortical atrophy. More accurate and informative psychometric tests are available in the shape of the Halstead-Reitan test battery,² and the advent of the EMI-Scanner means that information about the state of the cortex and ventricles can now be obtained painlessly. The delineation of the cortex by the EMI-Scanner is possibly a little less precise than that provided by air encephalography, but may well be adequate for most purposes.³ Most studies, including ours, have found the EEG to be singularly unhelpful in the diagnosis of alcoholic brain damage.

I support your view that a combination of

brain damage and continued excessive drinking represents a particularly serious threat to the lives of other road users, and if we are to take this threat seriously the assessment of convicted drunken drivers with the Halstead-Reitan test battery and the EMI-Scanner deserves our careful consideration not only as a possible index of current disability but also as a possible prognostic guide.

I hope shortly to produce a six-year follow-up of our series of 33 alcoholics which may tell us whether the likelihood of relapse is related to objective indices of the extent of brain damage.

COLIN BREWER

University Department of Psychiatry,
Queen Elizabeth Hospital,
Birmingham

¹ Brewer, C. and Perrett, L., *British Journal of Addiction*, 1971, **66**, 170.

² Filskov, S. B. and Goldstein, S. G., *Journal of Consulting and Clinical Psychology*, 1974, **42**, 382.

³ Huckman, M. S., Fox, J., and Topel, J., *Radiology*, 1975, **116**, 85.

Drinking drivers and the law

SIR,—I am prompted to write to you by reading again your leading articles (8 May, p 1103) concerning drinking drivers and the law and alcohol and the brain (15 May, p 1168).

If we are honest with ourselves, is it not true to say that alcohol is really a drug far more dangerous even than heroin? The heroin addict, whatever risks he runs for himself, is very little danger to others. If we are truly concerned for the damage done by alcohol, should we not demand that any public advertising of alcoholic drinks should be forbidden by law? The experience of the United States of America and, I believe, of India proved that prohibition is counterproductive essentially because it creates a situation advantageous to Mafia-type pusher groups. For this reason would we not be wiser to make hard drugs fairly readily available for those foolish enough to use them but punish any advertising of them severely? I visualise a situation in which drugs and alcohol could be purchased at moderate costs discreetly, rather like hard-core pornography. They would be in containers labelled "Drugs for Dopes."

Needless to say, penalties for driving under the influence of such drugs would be draconian. Because of the importance of our example drinking by doctors is even more disgraceful than is smoking.

KEITH NORCROSS

Department of Accident Surgery and
Orthopaedics,
Dudley Road Hospital,
Birmingham

Childhood accidents

SIR,—It is welcome to see in the review by Dr R. H. Jackson and Professor A. W. Wilkinson on the prevention of childhood accidents (22 May, p 1258) reference to the particular implications for the eye and vision in these. It is germane to this matter that during a visit of members of the Faculty of Ophthalmology to Russia in 1963 one of the matters which excited considerable favourable comment was the presence in every school that we visited (and those who have participated in such tours will know that the Russians are fond of showing off their schools) of a large rotating illuminated picture-postcard type of photograph holder. This demonstrated the

wide variety of methods and instruments with which a child could injure its eyes, including, for example, scissors in the hand, eating fork, bows and arrows, and so forth. We were told that considerable attention was paid to seeing that the children had these dangers brought well to their attention.

Ophthalmic surgeons would agree that the majority of childhood accidents to the eye are in a degree avoidable. If educational authorities could be persuaded to follow this example we might at least be making a beginning in the reduction, if not the elimination, of such accidents.

There is no doubt that ophthalmology, through the Ocular Safety Committee of the faculty and other appropriate bodies, would very much wish to play its part in determined efforts in this field of accident prevention. It is perhaps also appropriate to remember that this is the World Health Organisation's Ocular Injury Year.

M. J. GILKES

Sussex Eye Hospital,
Brighton

SIR,—At the BMA Clinical Meeting at Exeter recently we were shown a slide of children on holiday (presumably) riding on a farm tractor and trailer, which the speaker on the subject of farm accidents pointed out was illegal and the cause of many accidents to children. When it was asked why the local official children's playgrounds provided old farm tractors for children to climb on, and so breed the contempt of familiarity, the only reply given was that the playground tractors were stationary. When it was asked if the authorities concerned with safety would act in the matter the reply was that this was the responsibility of "another department."

A similar observation was forthcoming from officials on an exhibit on "Safety" at the Devon County Show... "they wouldn't say 'thank you' if we poked our noses into the affairs of another local authority department, and anyway we don't know under which department council-provided playgrounds are equipped."

At least let us have sufficient sense not to encourage children to play with potential dangers.

"Never, never, let your gun pointed be at anyone; That it may unloaded be, matters not the least to me."

JOHN B TRACEY

Pinhoe,
Exeter

What do community physicians do?

SIR,—Having read your recent leading article on "The community physician of the future" (24 April, p 976) I was intrigued to notice that there was no mention of what such a doctor would actually do. It is clear that his training is immaculate, that he is highly qualified, and that he has a considerable knowledge of epidemiology and statistics together with "social policy and social sciences" (he probably would be able to define what sociology actually means). To cap it all he is also fully versed in the principles of administration and management. Not only all this but he has also passed exams and it clearly would be helpful to him in his career if he had higher medical qualifications.

Such a highly qualified person is obviously of extreme potential value to "the community." But what does he do and what is "the community"? If a drain is seen to be seeping foul-smelling green liquid is the community physician notified, summoned, and questioned, or does he delegate one of his staff to look into the matter, or is it nothing whatsoever to do with him at all?

For some years now the importance of community physicians has been emphasised in the *BMJ* and elsewhere and for a lesser length of time many doctors have been wondering what their role is. I imagine that I have overlooked, like others, a clear exposition of their function, but it is odd to read your leading article and to find no mention of this. It is of course an understood and accepted maxim of the Western world that voids should be filled with working parties, subcommittees, and "long, hard looks" at trivia. Administrative expansion and "quality control" are probably cheaper than quality achieved.

PETER CARSON

Cardiology Department,
City General Hospital,
Stoke-on-Trent

Intermittent claudication complicating beta-blockade

SIR,—The conclusion of Dr J. Christine Rodger and her colleagues (8 May, p 1125) that "claudication is an important and possible reversible side effect of cardioselective and non-selective beta-blockade" is a sweeping statement based on anecdotal and not on controlled evidence.

Since short reports are limited to five references, it is regrettable that these authors should choose to refer to unqualified review statements while omitting any mention of recent and important original work. The controlled trial of propranolol in intermittent claudication by Reichert *et al.*¹ provided no evidence of deterioration of occlusive peripheral arterial disease that could be attributed to propranolol, in spite of the high daily doses used (240-1600 mg). The careful discussion by these authors and their plea for more extensive, carefully controlled trials should be heeded.

Your leading article (15 May, p 1165) refers to the benign nature of claudication but emphasises the seriousness of the underlying disease, as disclosed by the mortality from coronary thrombosis. It is important, therefore, that the highlighting of dubious relationships to therapy should not prejudice the use of beta-blockade in situations where its value seems to have been established.

A. F. LANT
D. O. GIBBONS

Department of Therapeutics,
Westminster Hospital,
London SW1

¹ Reichert, N., *et al*, *Clinical Pharmacology and Therapeutics*, 1975, **17**, 612.

Older smokers

SIR,—In your leading article on this subject (10 April, p 859) the proportions of men and women in the United Kingdom who are cigarette smokers are quoted as 65% and 42%, respectively. It is stated that these figures came from the WHO Survey on Smoking and Health in the European Region 1974-75. No such