References

- ¹ McAllen, P M, and Marshall, J, *Lancet*, 1973, **1**, 1212.
- ² Abdon, N-J, and Malmcrona, R, *Acta Medica Scandinavica*, 1975, **198**, 455.
- ³ Walter, P F, Reid, S D, and Wenger, N K, *Annals of Internal Medicine* 1970, **72**, 471.
- ⁴ Goldberg, A D, Raftery, E B, and Cashman, P M M, *British Medical Journal*, 1975, **4**, 569.
- ⁵ Sharpey-Schafer, E P, *British Medical Journal*, 1956, **1**, 506.
- ⁶ Hirsch, C S, and Martin, D L, *Neurology (Minneapolis)*, 1971, **21**, 682.
- ⁷ Krasnow, A Z, and Bloomfield, D K, *American Heart Journal*, 1976, **91**, 349.
- ⁸ Driver, M V, and Selby, P J, *Electroencephalography and Clinical Neurophysiology*. In press.
- ⁹ White, P T, et al, *Neurology (Minneapolis)*, 1961, **11**, 354.
- ¹⁰ Rubenstein, J J, et al, *Circulation*, 1972, **46**, 5.
- ¹¹ *British Medical Journal*, 1976, **4**, 1158.
- ¹² Easley, R M, and Goldstein, S, *American Journal of Medicine*, 1971, **50**, 166.
- ¹³ Fairfax, A J, and Lambert, C D, *Journal of Neurology, Neurosurgery, and Psychiatry*, 1976, **39**, 576.
- ¹⁴ Van Durme, J P, *American Heart Journal*, 1975, **89**, 538.
- ¹⁵ Lindén, V, *British Medical Journal*, 1975, **2**, 87.
- ¹⁶ Sarachek, N S, and Leonard, J J, *American Journal of Cardiology*, 1972, **29**, 451.
- ¹⁷ Clarke, J, et al, *Lancet*, 1976, **2**, 508.

(Accepted 25 March 1977)

Today's Treatment

Diseases of the urinary system

Treatment of glomerulonephritis by drugs

J S CAMERON

British Medical Journal, 1977, **1**, 1457-1459

The story of attempts to treat glomerulonephritis is one of hopes raised, later to be dashed—despite the fact that we now possess some understanding of what processes underly the major forms of glomerulonephritis. The advances that we have made have been the result of fortunate accident, rather than inspired logic.

The major advances in managing patients suffering from glomerulonephritis have been the relief of oedema by powerful diuretics, the treatment of infection by antibiotics, and the control of hypertension by hypotensive drugs. These treatments are available to any patient, and the fact that they do not influence the underlying disease should not let us underestimate them; particularly since it seems almost certain that control of hypertension can prolong life and preserve renal function.

This happy story is not, however, the subject of this essay. Here, we must attempt the more depressing exercise of judging our ability to manipulate the events leading to glomerular injury in glomerulonephritis. It must be said straight away that many patients with different forms of glomerulonephritis do not need any specific treatment, even though they may be quite ill. An example is classic post-streptococcal glomerulonephritis with endocapillary proliferation within the glomeruli, but without extracapillary crescent formation. Even though severe hypertension may be seen in the acute phase, or oliguria and uraemia even need dialysis, there is no evidence that treatment of the nephritis is needed, or that full recovery cannot take place. Similarly, children and young adults who suffer repeated attacks of haematuria, and show IgA in the mesangium of their glomeruli with normal histological pattern on optical microscopy, rarely go into uraemia. One-quarter of adult

patients with membranous glomerulopathy, and at least half the children with a similar appearance, will show complete clinical resolution of the disease within two years; treatment can only harm them with its unwanted side effects. Thus, treatment in glomerulonephritis, especially when its effect is uncertain, should be reserved for groups of patients identified histologically and clinically as those most likely to do badly, and least likely to show spontaneous remission.

We know that most forms of glomerulonephritis appear to arise from the formation, circulation, and glomerular deposition of immune complexes, which can induce an inflammatory response. These events, and the resultant glomerular scarring, lead to the proteinuria, haematuria, hypertension, and loss of filtering surface that characterise glomerulonephritis clinically. The various histological appearances represent the glomerular reaction to various rates of deposition of complexes of different composition, size, and inflammatory potential. The fundamental lesion in the nephritic individual is a relative inefficiency in eliminating foreign antigens, which persist in the circulation in the presence of the antibody response, and may thus form large amounts of soluble complexes. This view of glomerulonephritis suggests that it is the result of various forms of relative immune incompetence—that is, a manifestation of immunodeficiency. A little glomerular disease results from the combination of antibody with specific antigens on the glomeruli—for instance, antiglomerular basement membrane antibody nephritis.

Antigen removal

The various therapeutic attacks on this condition are summarised in table I. Although we have identified over 30 different antigens in occasional cases of glomerulonephritis, in most patients the antigen cannot be removed as it is unknown. Many of the antigens in human glomerulonephritis may, in fact, be the common pathogens to which we are all susceptible and the major lesion, mentioned above, may lie with the host. We can, however, remove the antigen in some patients because it includes the treatment of a persistent infection. The treatment of

TABLE I—Possible modes of treatment for glomerulonephritis

Removal of antigen: From environment From patient From plasma	Alter immune balance of individual host: Suppress or remove antibody Increase antibody Add excess antigen	Inhibit or remove mediators of inflammation: Complement White cell proteases Fibrin/plasmin Platelets Kinins Lymphokines Histamine and SRS-A Prostaglandins
--	--	--

syphilis, subacute bacterial endocarditis, or infected juguloatrial shunts inserted for hydrocephalus by surgery and antibiotics is followed in most cases by resolution of disease; or at least an arrest in decline in renal function.

In malaria, on the other hand, antimalarial chemotherapy in conventional doses so as to eliminate the replicating parasite from the circulation does not arrest the renal disease. This suggests that either enough malarial antigen persists in the tissues to continue the immune events or that a secondary mechanism affecting autologous antibody against something in the glomeruli has taken over. It is also possible (although much less likely) that some other event associated with the malaria, which induces quite profound immunosuppression, determines the nephritis, and not the malarial antigen itself.

Alterations in immune balance

Specific treatments, however, are not available for most of the few identified antigens—for example, hepatitis B virus. Sometimes the antigens are endogenous (such as DNA or thyroglobulin) and cannot be eliminated. In these cases the patient's immune responses might be manipulated so as to render him incapable of forming immune complexes or of eliminating the antigen for which no chemotherapy is possible. Since glomerulonephritis is supposed to result from immunodeficiency, the best result would be to shift the patient towards greater immune responsiveness; alternatively, immunosuppression might reduce antibody formation to such a low level that, again, toxic immune complexes were not formed. This runs the risk, if the antigen is a replicating foreign organism, of increasing infection.

As we begin to understand the complexity of the immune system, immune stimulation remains in its infancy. Antibody production appears to depend on a complex interaction between several types of lymphocytes acting upon the differentiated plasma cell that produces the antibody. Drugs that "stimulate" or "suppress" lymphocyte function will have effects at present unknowable on this system. One drug, however, is already being used experimentally that appears to increase the ability to react to some antigens. This is the antihelminthic drug levamisole. It suffers from the disadvantage that it may induce agranulocytosis in some people, but drugs that can achieve

the same effect will undoubtedly be used in the future treatment of glomerulonephritis.

The alternative approach, that of depressing the immune response, has been more exploited, although with little success in primary glomerulonephritis. Possibly in some cases the nephritis would be made worse by "immunosuppression," and some evidence suggests that this does indeed happen. I think that what benefit has accrued from using immunosuppressive drugs has resulted from their ability to inhibit inflammatory responses (discussed below). Immunosuppression, however, is probably necessary to accompany the depletion of antibody or immune complexes that may be achieved by repeated plasmapheresis.

Plasmapheresis—This complex and expensive treatment has been tried experimentally in patients with the most severe forms of antglomerular basement membrane and soluble complex nephritis, associated with extensive extracapillary crescent formation round the glomeruli and severe decline in renal function. Normally the removal of antibody leads to its synthesis in increased quantity, and immunosuppression is necessary to depress this response. Plasmapheresis appears to benefit patients with anti-GBM nephritis, but it is not yet clear whether the results in severe soluble complex disease are better than chance or those obtainable by other regimens. Nor is it clear whether the benefit, if present, results from the depletion of mediators of inflammation from the circulation, rather than the removal of toxic antibody or toxic complexes.

Inhibition or removal of mediators of inflammation

Acute injury in experimental glomerulonephritis results from the interaction of complement and polymorphs. We know much less about chronic injury in human or experimental glomerulonephritis, but it appears that complement is not necessary, and that coagulation mechanisms, both fibrin and platelet-mediated, are intimately concerned in inducing, amplifying, or maintaining the injury. This knowledge has resulted in the use of drugs capable of inhibiting one or more aspects of the inflammatory response, alone or in combination. These drugs fall into several classes. Firstly, there are drugs which, as well as possessing some anti-inflammatory effects, are immunosuppressant: corticosteroids such as prednisone; mustard-like drugs such as cyclophosphamide and chlorambucil; and purine antagonists such as azathioprine. Then there are drugs that affect prostaglandin synthesis and are anti-inflammatory (although these agents all inhibit platelet aggregation and release), such as aspirin and indomethacin. Finally, there are drugs that are thrombin antagonists, such as heparin and warfarin, and platelet antagonists, such as dipyridamole, sulphinpyrazone, cyproheptadine, and, as noted above, aspirin and indomethacin again.

TABLE II—Results of controlled trials of inhibitors of inflammation in human glomerulonephritis

Form of glomerulonephritis	Drug	Duration of trial (yr)	Effect on glomerular filtration rate	Author(s)
Proliferative	Corticosteroids	2-4	None, mortality increased	Black <i>et al</i> (1970)
Mesangiocapillary	Corticosteroids (alternate day)	1	None	ISKDC (1977)
Proliferative	Corticosteroids + azathioprine	2	None, mortality increased	MRC (1971)
Proliferative	Corticosteroids + azathioprine	1	None	Sharpstone <i>et al</i> (1969)
Proliferative	Corticosteroids + azathioprine	1	None	ISKDC (1970)
Proliferative	Cyclophosphamide	2	None	Koelz <i>et al</i> (1975)
Proliferative	Cyclophosphamide	0.25	Some	Balslov <i>et al</i> (1975)
Proliferative	Cyclophosphamide	1	Some	German collab study (1977)
Proliferative	Indomethacin	2-4	Proteinuria initially diminished	MRC (1977)
Proliferative	Indomethacin	1	None	German collab study (1977)
Membranous nephropathy	Corticosteroids	2-4	None	Black <i>et al</i> (1970)
Membranous nephropathy	Corticosteroids + azathioprine	2	None	MRC (1971)
Membranous nephropathy	Azathioprine	1	None	Canadian collab study (1977)
Membranous nephropathy	Cyclophosphamide	2	None	Donadio <i>et al</i> (1974)
Membranous nephropathy	Chlorambucil	1	None, proteinuria decreased	Laguerre <i>et al</i> (1975)

ISKDC = International Study of Kidney Disease in Children.
MRC = Medical Research Council (Great Britain).
Collab = Collaborative.

CORTICOSTEROIDS AND IMMUNOSUPPRESSANTS

There are considerable data on the effect of these different agents in various forms of experimental nephritis. The results are easily summarised: no beneficial effect of corticosteroids or immunosuppressants has been described, and several reports suggest that heparin may make both soluble complex nephritis and anti-GBM nephritis worse. The only improvement recorded has been with antiplatelet agents. Several antiplatelet agents not only improved the histological appearance in various forms of experimental glomerulonephritis but also diminished proteinuria. Most of these experiments, however, concerned only mild nephritis.

Nevertheless, all these agents have been used in human glomerulonephritis alone and in combination. The pattern has been depressingly similar. Firstly, a few pilot studies suggest that a particular drug in a particular form (or all forms) of nephritis produces benefit. Then a larger uncontrolled study kindles enthusiasm, with glowing accounts of severe nephritis made better. Finally, one or more controlled studies fails to show an effect on any form of nephritis tested, with a considerable body of side effects in the treated patients. Table II summarises the controlled studies carried out to date in membranous nephropathy and in mixed proliferative glomerulonephritis, employing anti-inflammatory corticosteroid and immunosuppressive drugs (full bibliography in first reference). Apart from some short-term benefit from cyclophosphamide, not confirmed in the longer-term study from Newcastle, and diminution in proteinuria of membranous patients treated with chlorambucil, no effect has been seen in any study. Indeed, the mortality of the treated group in the Medical Research Council's trials of corticosteroids and of corticosteroids and azathioprine was *higher* than in the control group, even when only deaths from uraemia were considered. This suggests that in some patients the glomerulonephritis was made worse.

Few of these trials were ideal, and some doubt remains in several directions. In particular, the Medical Research Council trials included only few patients with membranous nephropathy; and in the corticosteroid trial the dosage was not constant, nor the course very prolonged. Others have published data that suggest an effect with six or eight months' treatment with corticosteroids, and, although other data (including our own) do not support this, a further trial of corticosteroids with or without immunosuppressant drugs seems justified. Such a trial is at present under way in the United States, but, unless this shows an effect, present evidence suggests that membranous nephropathy should not be treated. In any case, one-quarter of patients with this condition remit spontaneously, and any decline in renal function is almost confined to those with a persisting nephrotic syndrome; nor does it appear until five years or more from the discovery of the disease. If any treatment is to be given to this group of patients it should be confined to those with a high chance of a serious future course.

A further doubt arises with treating proliferative glomerulonephritis by indomethacin. The Medical Research Council trial used only 100 mg/day in patients with predominantly normal renal function, whereas reports of successful restoration of declining renal function towards normal have been with higher doses, sometimes accompanied by small doses of cyclophosphamide. Again there seems to be a case for further trial of this drug, especially in view of its antiplatelet activity and the favourable results in animal nephritis.

ANTICOAGULANT DRUGS

Anticoagulant drugs have rarely been used in human disease on their own. Usually, antithrombin and antiplatelet drugs have been given against a background of immunosuppression in the belief that only if all possible routes of glomerular injury were inhibited would an effect be seen. This toxic combination of drugs should not be given to patients whose disease does

not satisfy strict criteria for serious disease. To date, therefore, "cocktails" of immunosuppressive and anticoagulant drugs have been used only in severe proliferative glomerulonephritis, either with extensive epithelial crescent formation or a mesangio-capillary proliferative glomerulonephritis and persisting severe proteinuria. I think that these combinations should not be used for milder forms of glomerulonephritis when many years' treatment will be needed to establish their worth. Nevertheless, a trial of such treatment in membranous nephropathy is under way in Australia, in parallel with a similar trial in mesangio-capillary glomerulonephritis.

In the absence of controlled trials we are still in the second stage of evolution with these drugs—the stage of enthusiastic reports. Since some of these have emanated from our own unit—their enthusiasm tempered I hope, by caution—it is difficult for me to give a balanced account. The problem is to know whether the prognosis of the treated patients would have been, in the absence of treatment, as bad as we suppose on the basis of previous data. Usually this turns out not to have been the case, but for patients with proliferative glomerulonephritis and extensive crescent formation (> 60–70% of glomeruli) the prognosis does appear to be bad. Also, the treatment does not seem to work if decline in renal function results in the patients being anuric or needing dialysis, although we have now seen recovery in one such patient. It will be difficult to have controlled trials in these rare, extremely ill patients, but in their absence we must remember that the prognosis of the patient now includes regular dialysis and transplantation with a reasonable prospect for survival. Certainly, I would not use this treatment myself without decline of renal function over a short period and severe histological change.

Reference

- 1 Cameron, J S, in *Perspectives in Clinical Immunology, II: The Kidney*, ed J Zabriskie, H Villareal, and E L Becker. New York, John Wiley, 1977 (in press).

ONE HUNDRED YEARS AGO As long ago as 1860, I well remember, when a pupil of Mr John Tyler Featherstone, assisting that gentleman to insert a seton composed of six cords of silk into a large goitre which was compressing both in the mesial line and bilaterally the larynx of a boy aged 10. The seton remained in, as far as I remember, about eight weeks, being from time to time moved about so as to induce inflammatory action and to give outlet to the resulting pus. The case was entirely successful, and I saw the patient only a few years ago—a perfectly healthy man, with no affection whatever of the breathing or vocal functions.

Subsequently, while practising in Derbyshire, I have met with a great number of cases of bronchocele; but only a very small proportion of them could be described as of the suffocative type. In one case, that of a young woman aged 18, I followed out the same plan of treatment which I had remembered as so successful, and with good results. After the seton had been in some ten days, suppuration began, and the tumour was gradually lessening in size; but the patient was suffering from heart and kidney affection, and, after taking cold, pneumonia ensued, ending in uraemic poisoning and death seven weeks after the commencement of the treatment.

In the slighter cases, when seen early, I have generally found that the iodide of potassium ointment rubbed into the swelling once or twice a day, with the administration internally of both the iodide and bromide, or, in cases of an anaemic nature by no means uncommon, of the iodide of iron with bromide of potassium, will, if the treatment be steadily persevered in, produce a complete cure.

I may just remark that I found these cases by no means confined to Derbyshire or to a limestone formation. I had several cases under my care from the neighbouring counties of Nottingham and Stafford. (*British Medical Journal* 1877.)