in some cities in the USA,⁶ but the general level of health consciousness and the public attitude to family medicine and emergency care are so different in the USA that it is far from clear that American experience is relevant to the United Kingdom.

The number of lives that MCCUs can be expected to save in the United Kingdom is therefore small, so establishing MCCU services on a wider basis must be considered in competition with other developments that may well seem less dramatic to the public. Until patients' behaviour patterns change, the only way the MCCU concept can usefully be extended in this country is probably by equipping every ambulance that deals with emergencies with a defibrillator and training the crew to use it. The capital cost would not be great, and relatively few lives could certainly be saved for a reasonable outlay, provided that the extra cost of more highly trained ambulance crews was not excessive. The increased cost of the crews would probably be unacceptable unless the different ambulance functions were more formally separated.⁷ A few well-equipped ambulances, manned by specially trained crews, should be used in emergencies, while the "bus" functions of ambulances could be performed by simpler vehicles, manned by crews with only

basic training. MCCUs should be developed only as part of a national move towards a "two-tier" ambulance service.

We gratefully acknowledge the skill and endeavour of the crews who manned the Nottingham MCCU during the five years in which this and the two earlier studies were conducted. We also thank the ambulance controllers for co-operating in this study, and the Nottingham physicians, surgeons, and general practitioners for allowing us to record data from patients in their care. The research aspects of this study were supported by a grant from the DHSS.

References

- ¹ Hampton, J R, *British Medical Journal*, 1976, **1**, 201.
- ² Hampton, J R, Dowling, M, and Nicholas, C, *Lancet*, 1977, **1**, 526.
- ³ Hill, J D, and Hampton, J R, *British Medical Journal*, 1976, **2**, 1035.
- ⁴ Cameron, M, Wilkinson, F, and Hampton, J R, *British Medical Journal*, 1975, **1**, 384.
- ⁵ Joint Working Party of the Royal College of Physicians and the British Cardiac Society, *Journal of the Royal College of Physicians of London*, 1975, **10**, 5.
- ⁶ Cobb, L A, *et al*, *Circulation*, 1975, **52**, suppl No 3, p 223.
- ⁷ Howat, J G M, and Kontny, E L, *British Medical Journal*, 1977, **4**, 1298.

(Accepted 17 March 1978)

If I Had...

Cataract

BRIAN HARCOURT

British Medical Journal, 1978, **1**, 1121-1122

In occasional idle moments I look into my Snellen chart and realise that I can now read unaided only the 6/6 line, whereas it used to be the 6/5 line. I am very aware of the difference, and so sometimes ponder on my likely reactions if real trouble came my way and the clarity of one or both of my crystalline lenses became impaired.

Causes and symptoms

Cataract, of course, is an emotive word, and most ophthalmic surgeons sensibly tend to avoid using it with patients, unless matters have reached a stage where surgical treatment is envisaged, or unless directly questioned. They prefer to use the more innocuous, yet entirely accurate, term "lens opacities." The causes of cataract are numerous (see table), but by far the commonest is aging, and these senile changes are more frequent in diabetic patients and occur in them at an earlier age.

When the lens opacities are not congenital or the result of external agencies such as irradiation or trauma, both eyes tend to be affected. Even in such circumstances, however, initially deteriorating vision may affect one eye only, and this asymmetry may persist for some years. The symptoms are principally

painless progressive loss of distance visual acuity, for the optical requirements of reading vision are much less stringent. In particular, if the nucleus of the lens becomes sclerotic, there is often a change in refraction towards myopia as cataract develops, so that in the early stages the patient may even feel

Principal causes of cataract

Congenital		(a) Hereditary
		(b) Acquired: rubella
Acquired		(a) Senile
		(b) Secondary to ocular disorders: degenerative myopia, uveitis, retinal detachment, retinitis pigmentosa
		(c) Metabolic: diabetes, galactosaemia, homocystinuria
		(d) Traumatic
		(e) Irradiation: infrared: . . . glassblowers' cataract, x-rays and gamma rays

temporarily advantaged by being able to read without spectacles.

The same specific comments are heard from patients repeatedly: they have difficulty in recognising acquaintances in the street, which may cause social embarrassment; they fail to read bus numbers and traffic signs clearly; loss of confidence when driving is common, especially in older patients, and this is partly the result of the other main symptom of cataract—

General Infirmary, Leeds 1

BRIAN HARCOURT, MB, FRCS, consultant ophthalmic surgeon

dazzle and glare, caused by the tendency of the lens opacities to scatter the incident light within the eyes. For this reason, driving at night against the glare of oncoming headlights becomes a particular trial and hazard. When a cataract is unilateral and dense the patient may be aware that his field of vision is restricted on that side, and recognise his deteriorating ability to carry out fine manual tasks because of the loss of stereoscopic vision.

What I would do

If I were to develop lens opacities I hope that I would have an objective scientific curiosity regarding my own symptoms. Having discussed patients' glare problems so often, it would be interesting to find out at first hand whether the tinted spectacles and the clip-on anti-dazzle lenses, the eyeshades, and the broad-brimmed hats that are advocated really make much difference. Again, do colours become progressively yellower as is sometimes alleged? One of my first reactions would be to carry out urine tests to exclude glycosuria. This is imperative, and a test I am sad to say that is often omitted until patients are seen in the hospital outpatient department. I hope that I would be honest with myself regarding my continued fitness to drive; the number plate recognition test is roughly equivalent to 6/12 vision with two eyes together and best optical correction. Only three lines lost on the Snellen chart from the 6/6 standard takes a patient outside those limits, although many continue to persuade themselves otherwise.

For most people who do not drive 6/24 to 6/36 vision with both eyes together is taken as the stage at which they are likely to benefit from cataract surgery. This is because of the serious optical problems posed by removing the focusing lens from the eye. About one-third of the total dioptric power of the eye is furnished by the lens, and its removal renders the patient highly hypermetropic (aphakic). A spectacle lens of roughly +10 dioptres strength is required to refocus the eye, and this has the effect of magnifying the image size by about 33%. A contact lens has less magnifying effect (8%), and an artificial lens introduced into the eye at operation has no appreciable magnifying effect at all.

On account of this magnification and the accompanying peripheral distortion, cataract spectacles are difficult to adapt to, particularly for the elderly. In addition, if one eye has been operated on and the other has not, glasses cannot be worn that correct the vision of both eyes simultaneously; the pronounced difference in image size gives rise to constant double vision. These problems are much reduced by wearing contact lenses and are eliminated by intraocular lenses. Nevertheless, elderly patients often find contact lenses difficult to handle, especially if they have had cataracts removed from both eyes, when seeing sufficiently well to put the first contact lens in place is a real problem. These difficulties are likely to be reduced with the development of continuous-wear aphakic contact lenses that are just beginning to be a reasonable long-term prospect.¹

Operating techniques

In recent years conventional cataract surgery techniques have reached a high standard of safety and effectiveness. Fine suturing by using an operating microscope gives excellent wound closure, while relaxation anaesthesia allows perfect control of the eye during surgery. Nevertheless, the corneo-scleral wound is relatively large and takes some time to heal. Thus prominence is now given, particularly in the United States,^{2,3} to phacoemulsification—whereby the lens is fragmented by ultrasonic probe before aspiration through a finely controlled system of fluid inflow and outflow. This is done by using instruments that can be introduced through a small opening in the eye, thus shortening the time for wound healing. The introduction of intraocular lenses, however, has tended to

cancel out this advantage, as the lens cannot be introduced into the eye except through a fairly generous incision.

If I were developing cataract in both eyes I would certainly want to have at least one eye operated on at an early stage so that I could continue to work and drive. Even with a unioocular cataract I would wish to have early treatment in order to maintain stereoscopic binocular vision. If I was under 65 in the present state of knowledge I would not want to have an intraocular lens implant. Although recent results of this specialised surgery in experienced hands have been very encouraging,⁴⁻⁶ I feel that not enough is yet known about the long-term effects of the plastic within the eye for me to want to run any risk if I was still dextrous enough to handle contact lenses easily, and my life expectancy was over 10 years. I would certainly want to wear contact lenses, rather than cataract spectacles, and if I were intolerant of those I might consider having an intraocular lens implanted in one eye as a secondary procedure, although I know that such an operation is not without risk. I would put myself in the hands of a surgeon who would carry out a conventional cataract operation using the operating microscope.

My ardent desire for a skilled anaesthetist goes without saying; and I would make a special point of thanking him personally after the operation. Unless the cataract surgery was being carried out before I was 40, when extracapsular lens extraction is necessary, I would certainly not want the lens removed by phacoemulsification. It has been claimed that this allows one to be out on the golf course within a day or two of the operation. I don't play golf, and would be happy with a well-earned rest after an operation which had to last me for the rest of my life, safe in the knowledge that there was no risk that some of my corneal endothelial cells might have been permanently damaged—either by ultrasonic energy, or by excessive irrigation or instrumentation during my operation.

I would not have both eyes operated on during the same admission (which is some surgeons' routine); I would like to get used to my optical problems and to know about any possible operative complications with the first eye before I was irrevocably rendered bilaterally aphakic by the second operation. I would try hard to be sensible in the postoperative period, avoiding too much stooping, lifting, and carrying; if anything went wrong, I could then, with a clear conscience, blame everything except myself. If this is thought to be rather unadventurous on my part, I would point out that ophthalmic surgeons do tend to be cautious as well as scrupulous. It is a specialty which in general appeals more to the birdwatcher than the flank forward.

General reading

Duke-Elder, S, *System of Ophthalmology*, vol xi. London, Henry Kimpton, 1969.

References

- ¹ Pierse, D, and Kersley, H J, *Transactions of the Ophthalmological Societies of the United Kingdom*, 1976, **96**, 11.
- ² Kelman, C D, *American Journal of Ophthalmology*, 1967, **64**, 23.
- ³ Kelman, C D, *American Journal of Ophthalmology*, 1969, **67**, 464.
- ⁴ Binkhorst, C D, *American Journal of Ophthalmology*, 1975, **80**, 184.
- ⁵ Pearce, J L, and Ghosh, T, *Transactions of the Ophthalmological Societies of the United Kingdom*, 1977, **97**, 84.
- ⁶ Jaffe, N S, et al, *American Journal of Ophthalmology*, 1978, **85**, 24.

ONE HUNDRED YEARS AGO Mr W Eassie read an exhaustive paper at the Society of Arts, last week, on cremation as practised in many parts of the Continent; and in a very able manner reviewed the work that had been done in this respect in various parts of the Continent, and graphically described the apparatus and *modus operandi* of the several exponents of modern cremation. By one apparatus, cremation was completed in two hours, at the expense of three shillings. (*British Medical Journal*, 1878.)