Two years ago, in a study ("The BBC's medical programmes and their effects on lay audiences") for the GAC by the department of science and features, it was stated that "the responsibilities of this department, where medicine is concerned, are no different from those which apply to other scientific topics dealt with in our programmes." This tidy façade I thought to be a product of superficial reasoning. It takes no account of the connotations and imponderables in medicine or of the unique influence and immediacy of matters medical on the attitudes and lives of the lay public.

The approach to medicine could not have been thought out. How else to explain "It is programmes that make policy, not policy that makes programmes"?—a remarkable utterance by the head of science and features department at the Royal Society of Medicine in 1976. This slick thought-dampener, far from being concealed, was flaunted later before the GAC and was accepted. Presumably it is still dogma in medical broadcasting.

My respect and admiration for the BBC are undimmed by the arguments you have published, which tend to focus on irritating superficialities. It is deeper, structural flaws that should be examined and corrected, preferably by the medical profession and BBC together.

JOHN APLEY

Bath

Treatment of tuberculosis

SIR,—I in turn am surprised at the different emphases in treating tuberculosis in the Liverpool School of Tropical Medicine (Dr D Robinson, 22 April, p 1053).

(1) I believe that where possible treatment should be based on the results of controlled trials and that inspiration, eccentricity, or years of experience are insufficient as alternative bases.

(2) I did not assume that the effects of *Mycobacterium tuberculosis* and the syndrome of mycobacterial diseases are "the same thing." Indeed, the brief reference to opportunist mycobacteria simply drew attention to the fact that they behave differently. To discuss their treatment more fully would have required more space than the Editor would allow.

(3) I have never before heard tuberculosis referred to as a "subdisease." It must rank as the most important and fatal non-disease in history. Anyone who feels that the laboratory plays the major role in the management of this subdisease must have little practical experience of dealing with patients suffering from it. Perhaps that is why Dr Robinson is not aware that most doctors and indeed many laymen actually regard it as a fully fledged disease. Sensible clinicians do, however, work in close contact with sensible bacteriologists in managing the condition.

(4) The confusion does not appear to be in the paragraph but in the mind of Dr Robinson. I feel, and others more authoritative and experienced than myself seem to agree, that contacts should be investigated. Not to do so would allow cases to go undetected in the community.

(5) The side effects of the drugs are listed in my article and further information is available and referred to. Isoniazid-induced hepatitis is, as I stated, uncommon in Britain. My article makes clear that follow-up of patients is essential and that drug side effects are a problem to be aware of. Did Dr Robinson read it?

(6) Rifampicin has already shortened the standard regimen and the evidence is quoted. There are few contraindications to its use and it has now been used widely in pregnancy. Dr Robinson is badly out of date.

(7) Practice in Britain has now relegated strepto-

mycin to a somewhat less important role, though, as I state, it remains a first-line drug. Ethionamide and pyrazinamide are generally used as second-line drugs and both are mentioned in my article.

(8) Sputum culture is mentioned as a keystone in the diagnosis and is an important aspect of follow-up. The primary role in the management of tuberculosis, and with it the responsibility, should be that of the physician. A nurse or health worker is an invaluable aid in this but is surely not able to take primary responsibility.

I reply to all Dr Robinson's points because I believe he has over-reacted to my article and is almost entirely wrong in what he says. I do not wish your readers to be misled by his letter and would refer them back to my article and the suggested further reading to judge for themselves.

ANTHONY SEATON

Institute of Occupational Medicine,
Edinburgh

Diuretics in the elderly

SIR,—Your timely reminder of the dangers of prescribing diuretics to elderly patients (29 April, p 1092) fails to mention the problems of magnesium deficiency. That increased excretion of magnesium occurs has been known for some time¹ and symptomatic hypomagnesaemia has also been described.² The symptoms encountered in magnesium depletion include confusion, ataxia, tremor, nausea, and palpitations, all of which commonly occur in the elderly. Recently Ibrahim and Sutcliffe³ showed a high incidence of symptomatic magnesium deficiency in the elderly which responded well to replacement therapy.

As a number of elderly patients receiving diuretics may also be having digoxin it is important to remember that hypomagnesaemia is associated with an increased sensitivity to digoxin⁴ as well as with primary cardiac arrhythmias.⁵

I would therefore suggest that magnesium depletion should always be considered in elderly patients who become unwell, especially if on diuretics.

S WALDEK

University Department of Medicine,
Royal Hospital,
Sheffield

¹ Smith, W O, *et al*, *Clinical Research*, 1959, 7, 162.

² Wacker, W E C, and Parisi, A F, *New England Journal of Medicine*, 1968, 278, 772.

³ Ibrahim, I K, and Sutcliffe, R L G, *Modern Geriatrics*, October 1977, p 6.

⁴ Sellar, R H, *American Heart Journal*, 1971, 82, 551.

⁵ Chadda, K D, Edgar, L, and Gupta, P, *American Journal of Cardiology*, 1973, 31, 98.

SIR,—Once again Slow-K is quoted as a cause of ulceration of the small intestine (leading article, 29 April, p 1092). This is surely an extremely rare event and is almost always associated with pre-existing gut stricture or slow intestinal transit time, as in the reference cited by you.¹ Bearing in mind that approximately 600 million tablets of Slow-K are dispensed in the United Kingdom every year, Slow-K must be just about the best tolerated preparation on the market. It must not be confused with enteric-coated tablets.

DENIS BURLEY

Head of Medical Services,
Ciba Laboratories

Horsham, W Sussex

¹ Farquharson-Roberts, M A, Giddings, A E B, and Nunn, A J, *British Medical Journal*, 1975, 3, 206.

Sulphinpyrazone after myocardial infarction

SIR,—In response to your leading article (15 April, p 941) concerning rhythm abnormalities in the sulphinpyrazone cardiac infarction trial I should like to quote from our original paper¹:

"Although, as shown in table 2, more patients receiving placebo (111) than patients receiving sulphinpyrazone (83) had cardiac rhythm abnormalities at the first visit, deaths among patients with rhythm abnormalities (six in the placebo, and two in the sulphinpyrazone group) represented only 12% of all deaths; furthermore, the presence of abnormalities of rhythm at entry into the study did not appear to be a significant risk factor since 5.4% of placebo-treated patients with such abnormalities died as compared to 6.0% of placebo patients without them."

In addition, we have also calculated the death rates per 28-day exposure period $\times 1000$. The findings are shown in the table:

Treatment group	Rhythm abnormalities	
	No	Yes
Placebo	7.4 (38)*	6.6 (6)
Sulphinpyrazone ..	4.0 (23)	2.9 (2)
Combined	5.7 (61)	5.0 (8)

*Values in parentheses are actual numbers of deaths.

The statistical analyses employed in this trial (Cox model) permit adjustment in balances of entry criteria. The possible imbalance of rhythm abnormalities was thus adjusted for and did not account for any significant difference between unadjusted and adjusted P values.

The final report of this trial will include details of all aspects of electrocardiograms in so far as our data base will permit.

SYDNEY H KANE
for Policy Committee,
Anturane Reinfarction Trial

Pharmaceuticals Division,
Ciba-Geigy Corporation,
Summit, New Jersey

¹ Anturane Reinfarction Trial Research Group, *New England Journal of Medicine*, 1978, 298, 289.

Plasma urate changes in pre-eclampsia

SIR,—Mr W Dunlop and Mr J M Davison (25 March, p 786) query our use of plasma urate increments (25 February, p 467) to define grades of severity of pre-eclampsia. While we accept their general point that small increases of plasma urate cannot be used clinically to diagnose patients with pre-eclampsia, their claim that these changes may be "physiological" is not necessarily valid. That plasma urate levels above 350 $\mu\text{mol/l}$ (6 mg/100 ml) indicate significant pre-eclampsia has always implied that antenatal increases of plasma urate have already occurred. This is also clearly demonstrated in our recent paper. A priori in the development of pre-eclampsia there must be a time when plasma urate levels show small rises above an earlier baseline reading. The morbidity of this prodromal stage would not be expected to be high because in many women the plasma urate changes either coincide with or precede the elevation of the blood pressure. When elevation of the plasma urate occurs at or near term pregnancy-induced hypertension may not arise as the process will normally be cut short by spontaneous or induced delivery.

The problems of diurnal variability and laboratory variation in the measurement of plasma urate are real ones, as Mr Dunlop and Mr Davison clearly state. The former problem is not so important if blood is taken at a consistent time of day (in the morning in our study); and the latter problem can be reduced if repeated readings are demanded before a rise is accepted as real (three consecutive measurements in our study). By adopting these precautions we were able to identify a group with platelet count changes suggesting early consumption and consistent with the hypothesis that this is characteristic of the early onset of pre-eclampsia.

C W G REDMAN

John Radcliffe Hospital,
Oxford

JOHN BONNAR

Trinity College Unit,
Rotunda Hospital,
Dublin

Rubella vaccination

SIR,—May I raise again the question of our policy on rubella vaccination?

The Department of Health and Social Security issued a circular letter CMO 76-4 in February 1976 reaffirming the policy of directing the immunisation programme at girls between their 11th and 14th birthdays and susceptible groups of women of child-bearing age. It was admitted that this policy had not resulted in any decrease in the incidence of congenital rubella. Recent serological surveys had shown that about 10% of young women were susceptible compared with about 20% before the vaccination programme began in September 1970; the acceptance rate of rubella immunisation was between 70 and 75% and no further significant reduction in the proportion of susceptible mothers was expected until the 1980s.

It is known that the protection afforded by rubella vaccination lasts for at least eight years and probably for very much longer. In view of this, should we not reconsider changing our policy and aim at the elimination of outbreaks of rubella by the routine vaccination of all children, boys as well as girls? Obviously the girls would still receive individual protection, and the growing practice of screening pregnant women for rubella antibodies and offering immunisation after delivery to those found to be susceptible could continue with advantage.

The Joint Committee on Vaccination and Immunisation should be asked to look at this again.

L A CRAWFORD

Hertfordshire Area Health
Authority
Hemel Hempstead, Herts

Coronary heart disease and male sex hormones

SIR,—Several authors (25 February, p 472; 1 April, p 851) have recently presented indirect evidence suggesting that male sex hormones contribute to men's higher risk of coronary heart disease. However, several older investigations provide evidence against this hypothesis. Castration of teenage boys and young men apparently does not cause a significant difference in death rates from

cardiovascular disease.¹ Castration at older ages is not associated with any change in coronary atherosclerosis.² A comparison between young men who had recovered from a myocardial infarction and age-matched controls showed no significant differences in serum concentrations of testosterone or dihydrotestosterone.³ No consistent differences in the masculinity of body characteristics have been found in comparisons between young men who have coronary heart disease and controls^{3,4}; there were no significant differences in balding or in overall body hair, and observed differences in body form could for the most part be attributed to the greater obesity of the subjects with coronary heart disease. These observations suggest that male sex hormones are not a major cause of men's higher rates of coronary heart disease.

Other evidence indicates that major causes of men's elevated coronary heart disease mortality are men's higher rates of cigarette smoking and of the hard-driving coronary-prone behaviour pattern.^{5,6}

I L WALDRON

Department of Biology,
University of Pennsylvania,
Philadelphia

- ¹ Hamilton, J B, and Mestler, G E, *Journal of Gerontology*, 1969, **24**, 395.
- ² London, W T, et al, *Annals of Internal Medicine*, 1961, **55**, 63.
- ³ Phillips, G B, *Lancet*, 1976, **2**, 14.
- ⁴ Gertler, M M, and White, P D, *Coronary Heart Disease in Young Adults*. Cambridge, Harvard University Press, 1954.
- ⁵ Waldron, I, *Social Science and Medicine*, 1976, **10**, 349.
- ⁶ Waldron, I, *New England Journal of Medicine*, 1978, **298**, 57.

Cetiprin (emepronium bromide) in patients with prostatism

SIR,—In the advertisement for Cetiprin (emepronium bromide) on the inside back cover of the *BMJ* dated 15 April the manufacturers, KabiVitrum, state under "contraindications and precautions": "Whilst emepronium bromide is indicated in the early stages of prostatic enlargement with urinary frequency, it is contra-indicated in later stages with flaccid bladder and large amounts of residual urine." In *MIMS* the contra-indications listed are advanced prostatic hypertrophy, with special precautions in patients with glaucoma and gastric retention.

I wonder if many urologists would recommend suppressing bladder activity in patients with prostatism. There is an obvious danger that the reduced bladder awareness and activity achieved by this form of treatment may well lead to increased residual urine. My own view is that emepronium should be prescribed only for patients who are not in danger of developing residual urine. I would hope that a warning to this effect may be included in subsequent advertisements.

ERIC GLEN

Urodynamic Service,
Southern General Hospital,
Glasgow

* * * Mr Glen sent a copy of his letter to the manufacturers of Cetiprin, whose reply is printed below.—Ed, *BMJ*.

SIR,—Mr Glen rightly questions the use of emepronium bromide in patients with prostatic hypertrophy who are likely thereby to develop undesirable levels of residual urine. While we accept Mr Glen's comments as being con-

structive, it would, I feel, be wrong to regard prostatism, even in its earliest stages, as an absolute contraindication to emepronium therapy.

It is well known that even a small obstruction to the vesical outlet can increase bladder irritability, resulting in impaired control, frequency, and uninhibited bladder contractions. The balance between the degree of obstruction to the bladder outlet and the efficiency of the neuromuscular mechanism is important. Although large quantities of residual urine may be found in association with inefficient voiding contractions, it is possible that emepronium bromide can, by blocking involuntary uninhibited contractions and restoring normal detrusor-sphincter balance, improve bladder function and thereby reduce the amount of residual urine. I will be happy to refer Mr Glen to several urologists who hold this point of view.

ALLAN D LAUDER
Medical Director
KabiVitrum Ltd

London W5

Stricture of oesophagus associated with ankylosing spondylitis

SIR,—The letter from Mr A M Hay (4 May, p 1138) concerning a stricture of the oesophagus associated with ankylosing spondylitis is interesting but does not mention the association between spondylitis and Crohn's disease. The latter may present as dysphagia and strictures can develop.^{1,2} This may well have a bearing on this case.

A P WEETMAN

Royal Victoria Infirmary,
Newcastle upon Tyne

- ¹ Gelfand, M D, and Krone, C L, *Gastroenterology*, 1968, **55**, 510.
- ² Dyer, N H, et al, *Gut*, 1969, **10**, 549.

Is pancreatic isotope scanning worth while?

SIR,—It was interesting to have comments from Dr H J Testa and his colleagues (18 March, p 720) on our paper (4 February, p 282) concerning isotope scanning of the pancreas since Manchester must be one of very few centres with an interest in pancreatic diseases still apparently relying on the technique; indeed, can they tell us of any other in Britain or overseas?

Their own data on 200 patients confirms everyone's experience that abnormal scans are meaningless; of 107 patients with abnormal scans, 49 proved to have no pancreatic disease. Thus only 93 (the normal scans) of all their 200 studies could be of any clinical value, even if they were all correct. The Manchester false-negative rate was certainly lower than ours, but even they were only able to detect 64% of patients with a normal pancreas, which seems rather poor for a screening test. Scans of all types are in demand if they are clinically useful; indeed many are requested even when they are not. Isotope scanning has been available for many years. From our own experience and the reported survey of nuclear medicine departments it is quite clear that pragmatic clinicians have now discarded the technique; it is just not worth asking for.

There is now no shortage of alternative diagnostic techniques for the pancreas. Most are complex and expensive, but grey-scale