SHORT REPORTS

Disappearance of spermatozoa from ejaculate after vasectomy

The disappearance of spermatozoa from the ejaculate after vasectomy is said to be influenced by age, frequency of ejaculation, and time since operation.¹⁻³ Since their relative importance is not clear we examined the histories of 624 men who underwent vasectomy at the Margaret Pyke Centre from July 1971 to October 1972 to investigate the independent effects of these factors on the clearance of spermatozoa.

Methods and results

Details of data collection, surgical techniques, and methods of semen analysis have already been reported.¹ Coital frequency was recorded before and after the operation, postoperative information being obtained by postal survey one year after vasectomy. Monthly semen specimens were requested from all men until two consecutive azoospermic reports were obtained. The cumulative proportion of men who were azoospermic and its standard error were calculated at successive monthly intervals using life-table analysis.⁵ Most (54%) of the men were aged 30-39 years, 17% were aged 20-29 years,

and 29% over 40 years. The reported coital frequencies were similar before and after operation, with 68% reporting intercourse once or twice per week. Older men reported lower coital frequencies. Of those aged over 40 years 20% reported intercourse less than once a week compared with 9% at younger ages. Within six months of vasectomy 48% of men had cleared. At 12 months 88% were clear, and at 18 months 96% were clear. The remaining 4% comprised 22 men. Nine of them eventually cleared, one was reoperated on, and 12 were eventually lost to follow-up. Men in the 20-29 and 30-39 age groups cleared in a similar fashion while older men cleared more slowly. The difference in clearance rates between the younger and older men was statistically significant at all times after six months. Men aged 50 years and over were very slow to clear: 54% had cleared at 12 months and 77% at 18 months.

The effect of frequency of intercourse was examined separately for men aged 20-39 years and those aged 40-49 years. The figure shows the cumulative proportions cleared at monthly intervals by coital frequency. For the men aged 20-39 years the clearance rates were similar regardless of frequency of intercourse. For older men the higher the coital frequency the more rapid the clearance, the differences between the high and low coital frequencies being statistically significant from the fourth month onwards. The older men who reported intercourse more than three times per week, however, had clearance rates similar to those of the younger men.

Comment

These results suggest that age and coital frequency have a complex relationship to the clearance of sperm after vasectomy. Coital frequency seems to have little effect on clearance at ages less than 40 years, whereas at older ages the more frequent the intercourse the more rapid the clearance. But a coital frequency of more than three per week seems to be associated with rapid clearance regardless of age.

When counselling older men with low coital frequency the possibility of a delay in clearance of spermatozoa should be discussed. Nevertheless, most specimens classified here as "not clear" contained only a few non-motile spermatozoa. Since there may be considerable delay in becoming azoospermic more needs to be known about whether fertilisation is possible with such low sperm counts. Postvasectomy sterility may well be achieved earlier than azoospermia. If so, semen analysis should be begun sooner after vasectomy than is now recommended⁴ and the criteria for clearance altered.

¹ Freund, M, and Davis, J F, *Fertility and Sterility*, 1969, **20**, 163.

² Marshall, S, and Lyon, R P, *Journal of Urology*, 1972, **107**, 815.

³ Spencer, B, and Charlesworth, D, *British Journal of Surgery*, 1976, **63**, 477.

⁴ Barnes, M, *et al*, *British Medical Journal*, 1973, **4**, 271.

⁵ Armitage, P, *Statistical Methods in Medical Research*. Oxford, Blackwell Scientific Publications, 1971.

(Accepted 11 October 1978)

Department of Obstetrics and Gynaecology, Guy's Hospital, London SE1 9RT

R P MARWOOD, MSc, MRCOG, lecturer

Department of Epidemiology, London School of Hygiene and Tropical Medicine, London WC1E 7HT

VALERIE BERAL, MB, MRCP, lecturer

Cumulative proportions cleared at monthly intervals after vasectomy by the reported frequency of intercourse one year after operation (a) for men aged 20-39 years, (b) for men aged 40-49 years.

Non-motile sperms persisting after vasectomy: do they matter?

Seminal assays after vasectomy show in many cases that motile sperms have disappeared but non-motile sperms are still present. Whether these non-motile sperms can cause pregnancy is controversial.¹ Some clinicians²⁻³ advise patients in such cases that they are no longer fertile. Others⁴ insist that tests should be repeated until azoospermia is complete. If the first view is correct many men are being subjected to unnecessary testing and delay. Discussion in the past has been confused by pregnancies occurring after vasectomy as the result of spontaneous recanalisation of the vas. This can largely be

prevented by using the Schmidt technique of vasectomy, which virtually precludes the possibility of recanalisation.^{2,5} To test the hypothesis that non-motile sperms do not cause pregnancy we followed up after one year 200 men in whom the presence of residual non-motile sperms in postvasectomy seminal assays had been disregarded.

Patients, methods, and results

All vasectomies were carried out under local anaesthesia, using the Schmidt technique of light diathermy of the endothelium of each cut end of the vas without ligation then closing the sheath over only one end. Men were asked to leave a specimen of seminal fluid within four hours of collection after at least 10 ejaculations after their vasectomy. The specimen was mixed and 30 to 50 high-power fields examined. If sperms were present a count was made and any motility noted. If no sperms were found the specimen was centrifuged and re-examined. So long as no motile sperms were found, even though non-motile ones were present, the patient was advised that he was "clear" and that no further contraception was necessary. After 12 to 15 months all men were asked by letter, and if necessary by telephone, if their vasectomy had been successful in preventing pregnancy and offered a further seminal assay free of charge.

During the period under review 461 men submitted specimens for seminal assay. There were no motile sperms in the specimens of 200 of these 461 men but some non-motile ones remained. In most the non-motile sperms were only occasional, but in 30 the count was $0.5 \times 10^9/l$ or more and in two it was $24 \times 10^9/l$. Specimens had been submitted within four weeks of vasectomy by 44 men (22%), between four and eight weeks by 97 (48%), and after more than eight weeks by the remainder. Replies were obtained from 190 of the 200 (95%) on follow-up. No man reported the occurrence of a pregnancy during the year after his vasectomy. Forty-eight men (24%) submitted a further specimen for seminal assay. There were no motile sperms in any, but in five (10%) an occasional non-motile sperm could still be found.

Comment

Our hypothesis is that after vasectomy non-motile sperms do not cause pregnancies, and this hypothesis has not been disproved to date. Insisting on complete azoospermia must produce anxiety for some patients and their wives as well as increased costs. We suggest that absence of motile sperms in a postvasectomy seminal assay gives sufficient assurance of sterility, provided a fresh specimen is examined, and that non-motile sperms can be disregarded.

We thank Dr David Sugerman for assaying the semen of the patients.

¹ Staff of the Margaret Pyke Centre, *British Medical Journal*, 1973, **4**, 216.

² Edwards, I S, *Medical Journal of Australia*, 1977, **1**, 814.

³ Yeates, W K, *Proceedings of the Royal Society of Medicine*, 1973, **66**, 4.

⁴ Jackson, L N, *British Medical Journal*, 1973, **4**, 589.

⁵ Schmidt, S S, *Journal of Urology*, 1973, **109**, 296.

(Accepted 7 November 1978)

19 Gerrale Street, Cronulla 2230, and the Sutherland Hospital, Caringbah, NSW Australia

IAN S EDWARDS, MB, visiting associate obstetrician and gynaecologist
JOHN L FARLOW, FRCS, FRACS, visiting surgeon

Time and tides in Graves's disease: their implications in predicting outcome of treatment

Early clinical reports leave little doubt that, left untreated, the hyperthyroidism of Graves's disease undergoes a variable course with cycles which may vary from a few weeks to many years.^{1,2} Ord and McKenzie¹ noted that spontaneous relapses after spontaneous recoveries were not uncommon, and they quote Möbius as speaking of relapse as "the rule." Serial early (20-minute) uptake measurements under conditions of triiodothyronine (T3) suppression provide a means of following fluctuations in thyroid stimulation without interruption of treatment.³ We made a longitudinal study of these changes by frequent uptake measurements in a group of thyrotoxic patients undergoing a standardised course of treatment with carbimazole and T3.

Patients, methods, and results

Serial 20-minute ¹³²I uptake measurements were made on 44 patients with Graves's disease undergoing an 18-month course of treatment with carbimazole and T3. Carbimazole was given initially in a dose of 10 mg four times a day, reduced stepwise to 5 mg thrice daily according to response. T3 was given continuously in a dose of 20 µg four times a day. Uptake measurements were made before treatment (before and after T3, 20 µg four times a day for 8 days) and during treatment monthly for six months and two-monthly thereafter. Clinical outcome was judged exactly six months after withdrawing drugs (that is, 24 months from diagnosis).

The patterns of change in uptake in six of the 44 patients are shown in the figure. The restoration of normal uptake during the early months of treatment

Examples of iodine uptake patterns during the treatment of thyrotoxicosis with carbimazole and T3 (20 µg dose at 20 minutes). Outcome: E = euthyroid; R = relapse.

in patient 92 was the pattern seen most often in the 22 patients (50%) from the whole series who remained clinically euthyroid six months after withdrawal of drugs at month 18. Alexander *et al*⁴ likewise found that patients whose (20-minute) uptake fell by more than 50% during the first six months of treatment tended to retain normal uptake values on the last day of treatment and again when retested two months later. Nevertheless, four of the five remaining patients relapsed within six months of withdrawing treatment despite wide differences in behaviour of the underlying disorder. Patient 23 characterises the often erratic variation in uptake seen among those in the series whose uptake failed to normalise within 18 months and who mostly relapsed afterwards. It might be argued that the only fundamental difference between patients 23 and 92 was in time-scale, and that since the tendency of the former patient's uptake to fall with time was significant on regression analysis ($r^2 = 0.73$, $P < 0.05$) this patient would eventually have remitted had observations continued (at about the 28th month by extrapolation). Patients 89 and 27 were similar in that their uptakes were normal (albeit non-suppressible) at the time of diagnosis, but they subsequently rose despite control of thyroid hormone levels with carbimazole and suppression of TSH by T3. In both cases the uptake rose significantly with time ($r^2 = 0.82$ and 0.63 , $P < 0.05$, respectively), and in both the uptake by 18 months had become abnormally high despite T3 suppression. Patients 41 and 37 are further examples suggesting that when Graves's disease is treated symptomatically time is the most important determinant of outcome.

Comment

Our observations suggest that the underlying fluxes of thyroid stimulation in Graves's disease occur irrespective of symptomatic control with antithyroid drugs. While the pattern set during the first six months tends in most cases to reflect progress over the next 12 there are clearly exceptions, a fact which helps to explain the in-