some journals being free for various categories of doctors, and students can often get special rates. The scope of a journal is roughly indicated by counting the different sorts of articles. If a journal is just concerned with straight-forward medicine and surgery it scores zero on scope, but if it has articles on other topics, it scores accordingly.

Efficiency

Any system for keeping up to date must be easy to operate. It must be easy to put information in and easy to get it out. Most systems are good at one but not at the other. For example, if all you do is to subscribe to a journal or two it's easy to put information into your store: you just add your latest issue to the shelf. But it is hard to get information out. Even if the journal provides a good index it covers only a limited number of issues of that particular journal, so if you want to know what you have got on malaria you may have to consult many indexes. Alternatively, if you keep up to date by buying successive editions of well-known texts it is easy to retrieve information, provided they have good indexes, but there is no place for incoming information, which then gets tucked away and forgotten about.

The system of perpetual revision, explained above, gets round these difficulties by displaying together, in one place, all your information on, say, malaria, referred to by one index heading. By making additions to the index in the "husk" book you can refer to particular articles. Everything is together, all in one place. Adding to the husk index is

especially important when two interesting articles appear together on the same page (a rare conjunction). You would not normally think of consulting your section on pontine tumours to find out about tennis elbow, so if the articles appear on the same page the only access is through a suitably adapted index.

A weekly exercise

Sitting at my desk, once a week, with scalpel, glue, and the latest *BMJ* to hand, I take about an hour to bring out a new edition of *A Companion to Medical Studies*. It is important never to get behind: going through old issues is boring, and boredom is what undermines attempts to keep up to date. I throw away this week's *BMJ* this week, whether or not I have read it. It seems such a waste to throw it away without having read it that I usually do read it in time, and thereby, I fondly imagine, I keep up to date.

One objection to the medical scrapbook might be that it would expand very fast and soon become unmanageable. All I can say is that since 1975 my scrapbook seems to have advanced only 2 cm towards the end of its box. So there is no need to despair at the rate of increase of medical knowledge: it inches forward almost indiscernibly.

(Accepted 24 April 1979)

Personal Therapeutics

Insulin

P J WATKINS

British Medical Journal, 1979, 1, 1548-1550

A 14-year-old boy received the first injection of insulin in 1922 in Toronto, and R D Lawrence was first treated with insulin in London in 1923. Only short-acting soluble insulins were available until the 1930s, when protamine insulins were introduced (isophane and protamine zinc insulin), and in the 1950s insulin zinc suspension became available. The introduction of highly purified insulins in the 1970s has resulted in a bewildering number of insulins—no fewer than 18 are now available in the United Kingdom (table).

In this article I shall mention a selection of insulins for starting and maintaining treatment of insulin-requiring diabetics, but shall not consider emergency treatment. Insulin still needs to be given by subcutaneous injection, and experimental infusions of insulin intravenously or subcutaneously, implantable capsules of insulin, and oral insulin preparations are not yet suitable for routine use.

Aims of insulin treatment

Insulin is given firstly to save life and alleviate symptoms, and secondly to achieve the best possible control of diabetes with

blood glucose concentrations being maintained as near to normal as possible to minimise long-term complications. The first aim is relatively easy to attain, and in some elderly patients or in those who lack motivation or ability it is the only aim, because attempts at fine control are liable to cause disabling hypoglycaemia. In these patients a single daily injection of insulin will often suffice, but twice daily injections are usually needed to obtain very good diabetic control.

Insulin regimens

The rapid action of soluble insulin makes it the insulin of choice for initiating treatment in most of those with acute-onset diabetes, using two or more injections daily. For those less acutely ill it is often better to start treatment with a medium-acting insulin, either once or twice daily, since these insulins probably cause less abrupt swings of blood glucose concentration. Soon after diagnosis many patients with acute diabetes enter a stage of partial or more rarely complete remission, when quite small doses of almost any insulin are adequate to maintain good control. Sometimes insulin may even be withdrawn for a while—a practice that is not encouraged because after a few months the need for insulin is almost inevitable.

The best insulin regimen for maintaining good, long-term control is a mixture of a medium-acting insulin with a soluble insulin preparation taken morning and evening. During pregnancy, when even finer control is required, three injections may be needed—a soluble and medium-acting insulin mixture

is given in the morning, followed by soluble insulin before dinner and medium-acting insulin at bedtime. This is a useful regimen that helps to eliminate the hyperglycaemia before and after breakfast which is often a problem. For patients who lack the motivation or ability to mix insulins in a syringe because of old age, poor intelligence, or blindness, fixed mixtures of short- and medium-acting insulins (Rapitard MC; Mixtard) are useful.

Types of insulin preparation available in UK, according to animal source and purity

Type	Animal source	Purity
Soluble insulin*	Beef	
<i>Neutral soluble insulins</i>		
Actrapid MC†	Pork	Purified
Leo Neutral†	Pork	Purified
Nuso	Beef	
<i>Isophane insulins</i>		
Isophane NPH*	Beef	Purified
Leo Retard†	Pork	
<i>Insulin zinc suspensions</i>		
Semilente	Beef	Purified
Semitard MC	Pork	
Lente†	Beef	
Lentard MC†	Beef and pork	Purified
Monotard MC†	Pork	Purified
Ultralente	Beef	Purified
Ultratard MC	Beef	
<i>Mixed insulins</i>		
Rapitard MC†	Beef and pork	Purified
Leo Mixtard† (Neutral 30%, Retard 70%)	Pork	Purified
Leo Initald† (Neutral 50%, Retard 50%)	Pork	Purified
<i>Other insulins</i>		
Protamine zinc insulin	Beef	
Globin zinc insulin	Beef	

*Author's preference of conventional insulins.

†Author's preference of purified insulins.

‡Insulin mixtures.

MC = Monocomponent (highly purified) insulin.

Once-daily injections of either these insulin mixtures or a medium-acting insulin (isophane or Monotard MC) alone are useful in these patients, and have the advantage of preventing disagreeable nocturnal hypoglycaemia, which is both a danger and a fear, especially to elderly patients living alone.

Selection of insulin

Choice of individual insulin preparations is based on their duration of action, animal source (pig or beef), purity, and strength.

DURATION OF ACTION

Although insulin preparations may be broadly classified as having short, medium, or long durations of action, the effect varies substantially from one patient to another—for example, in some, soluble insulin has a longer action than isophane. This is probably due to different rates of absorption from the injection sites and to variable binding with insulin antibodies, and the persistence of small amounts of endogenous insulin may complicate matters. Duration of action in individual diabetics can be determined only by trial and error, and accurate assessment is greatly helped by self-monitoring of blood glucose concentrations.

I believe that the most useful medium-acting insulins are isophane and Monotard MC. Semitard insulin usually has too short an action and Ultratard is too long-acting, and even the conventional lente insulin (a mixture of semilente and ultralente) often causes tiresome nocturnal hypoglycaemia because of its relatively long action. Protamine zinc insulin no longer has a place in treatment because of its excessively long effect and because excess protamine partially converts added soluble insulin to protamine zinc insulin.

ANIMAL SOURCE AND PURITY

Insulin extracted from pigs is less antigenic than that from beef, and highly purified insulin less antigenic than conventional insulin. Highly purified pork neutral soluble insulin and insulin zinc suspensions are of relatively low antigenicity, but that of beef or isophane highly purified insulins has not yet been determined in the long term.

Many claims have been made for the advantages of highly purified insulins. Although the lower antigenicity may be desirable, there is still no evidence that insulin antibodies are harmful and none to associate them with diabetic complications. The most positive benefit is the virtual absence of fat atrophy, or its reversal if it has already occurred using conventional insulins, especially if the new insulin is injected into and around the hollows. The localised, red, itching patches at the injection sites that are common soon after starting insulin are said to be less common when highly purified insulins are used. Generalised allergy is rare but still occurs even with highly purified insulin, and desensitisation may be required. In the rare cases of insulin resistance some patients respond to the use of the highly purified insulins with a sharp reduction in dosage.

Unless patients have encountered the problems already described, their treatment need not be changed to the new, highly purified insulins. Nevertheless, some patients should receive highly purified insulins from the start of treatment. I believe that childhood diabetics should be treated with these insulins, and they are probably an advantage in patients needing insulin temporarily—for example, in pregnancy or at the time of acute illness or major surgery—since insulin allergy may develop if it is needed again in later years.

When treatment is changed from conventional to highly purified insulins or from beef to pig preparations a decrease in insulin dose is often needed, especially in those whose total daily dose is more than 60 units. The change in requirement is, however, variable, ranging from a decrease of up to 60% to an increase of around 10%. It is almost impossible to predict the size of the change in any individual and therefore wise to reduce the total dose by 20-30% immediately, thereafter making appropriate adjustments. The most useful highly purified insulins are: neutral soluble porcine insulin (Actrapid MC; Leo Neutral); Leo Retard (isophane); and Monotard MC (insulin zinc suspension). Fixed insulin mixtures such as Rapitard MC and Mixtard are sometimes useful.

INSULIN STRENGTH AND LABELLING

All insulins are marketed in the United Kingdom at both 40 U/cc and 80 U/cc; about two-thirds of patients use the 80 U/cc insulin to minimise the volume of the injection. In the rest of Europe 40 U/cc is used almost exclusively. In the UK most doctors who treat diabetics would like to introduce a sole strength of insulin at 100 U/cc together with a new syringe in the interest of safety, thereby eliminating the present confusion between units and syringe marks applied to two different insulin strengths. The Canadians have already achieved standardisation with a single 100 U/cc insulin, and a change to this system is being implemented in the United States. I should also mention that the traditional colour-labelling of insulins will probably disappear during the next year or two because the proliferation of insulins with a multiplicity of coloured labels has become a useless method of identification and some new labels will add to confusion by the use of similar colours for brand and strength.

Conclusions

Although the choice of insulins is wide, a small selection will fulfil the needs of most diabetics. Of the conventional insulins, soluble and isophane, either alone or in combination, have stood the test of time. In terms of highly purified insulins, combinations

of neutral soluble insulin (Actrapid MC; Leo Neutral) with Leo Retard (isophane) or Monotard MC (insulin zinc suspension) are equally effective. Useful fixed mixtures are Rapitard MC and Mixtard. It is to be hoped that in future years the number of preparations will be reduced, all insulins will be presented in a "highly purified" form, and only one strength will be available.

Further reading

- Sönksen, P H, *Clinics in Endocrinology and Metabolism*, 1977, 6, 481.
 Alberti, K G M M, and Natrass, M, *Diabetologia*, 1978, 15, 77.
 Oakley, W G, Pyke, D A, and Taylor, K W, *Diabetes and its Management*, 3rd edn. Oxford, Blackwell Scientific, 1978.
 Saunders, J, and Rogers, M L, *Insulins Chart*, 1978. Obtainable from the London Hospital (Whitechapel).

MATERIA NON MEDICA

Martyrs, monks, and manganese

We were on top of Gebel Katherine by sunset and gazed on a glorious panorama of mountains while waiting for the monk to unlock the chapel on the summit. The chapel commemorates Saint Katherine, whose body was miraculously conveyed there after her martyrdom in Alexandria, and it contains a tiny dormitory and kitchen for pilgrims. Two French friends and I arrived after a week of climbing and tramping the wadis between the mountains of southern Sinai.

The monk arrived smiling happily, though he had walked miles and climbed a considerable height just to unlock for us, and my friends set to work on our evening meal. After a good sleep we moved on via Gebel Musa to the convent.

The convent of Saint Katherine surpasses expectation. It contains within its chapel the remains of Saint Katherine, whose bones are credited with miraculous properties. There were plenty of monks, smiling and friendly, all speaking Greek. We were taken to His Beatitude the Archbishop, and, on the arrival of a bottle of araq, we sat in a circle together with one or two senior monks. After a good dinner in the guest room we settled down for the night.

I left my friends next morning to go back to work. This entailed a three-day trip to Abu Zenima, where I hoped to hitch-hike the remaining three hours to Suez in a manganese-ore lorry. Besides myself the party consisted of two young Bedouin and a camel for me to ride. The desert scenery, chiefly mountainous, rose well above expectation. The area was unmapped, but the assurance that the Bedouin, who spoke no English, probably knew the way and that the camel was sure to was a great comfort. We spent the first night in the Feiran oasis, where I got a most friendly reception from an outlying monk who put me up for the night in his small hut. He spoke only Greek, but he produced a new bottle of araq, which we emptied during the evening, and a Greek-English industrial phrase book, which, as the bottle emptied, produced increasing mirth. I woke at dawn, and proceeded on my way after the monk had given me two loaves of newly baked bread. The next night we slept in the open, and, joined by two elderly camel drivers, I opened my large tin of pilchards in tomato—a great success. The final day's journey brought us to Abu Zenima, a small port, chiefly used in connection with the manganese mines. The engineer, Mr Smith, was most welcoming and put me up for the night. The lorry was going at 10 am and I could perch on top of the manganese ore. I then produced my gift for him—thank heaven it had survived the journey—a bottle of Scotch whisky. We settled down to drinking it. I told him where I had trained. Did I know his two sisters who had trained there as nurses? Yes, one had been sister-in-charge of my male ward; the other I had met in 1940 at Tidworth, where she was theatre sister.

Next day I perched myself on three tons of manganese ore, waved farewell to Mr Smith, and jolted back to work. To those who do not often ride camels a good rule is two hours on then one hour off. This prevents the development (though only just) of what in the medical world used to be called pressure sores.—J C HAWKSLEY (retired physician, East Kennett, Wiltshire).

Prophet in the sand

I sipped ice-cold Pastis as the sun slid slowly out of sight at the end of the hottest March day I had ever experienced. At noon it had been 44°C in the shade, but in the wasteland of the Sahel "April is the cruellest month," when temperatures over 50°C are commonplace.

Some dark smudges of cloud appeared on the eastern horizon, the harbingers of the Hamartan, the Sahara's hot, dusty wind. Within minutes the dusk had changed colour to a pink-orange. The smell of sand was in the air. I hurried indoors to close the shutters and windows

before the sandstorm hit the house. Nevertheless, inside, the light of the hurricane lamp took on an eerie glow in the dust-filled atmosphere.

Two hours of unabated storm later, I needed to check the intravenous infusion of a 6-year-old child suffering from typhoid in the nearby ward. I was wearing a black baggy tuareg suit, and after winding four metres of fine muslin turban around my head, in the style of the Bellah tribesmen, I ventured out into the night. The visibility through the narrow slit in my turban was barely five metres, but the ward was only 50 metres from the house. When I reached the ward, I turned the doorhandle only to have the wind snatch it from my grasp and fling the door against the banco wall of the ward. My entrance was even more spectacular, with a great cloud of dust surrounding me.

The sick child's parents looked aghast. The father prostrated himself at my feet crying, "Mohammed, Mohammed!" I struggled to close the door, then pulled down the lower part of my turban, and greeted them in Tamachek with "Alkharrhass."

"Ayoh, nasara!" exclaimed the mother ("It's only the white man!"), obviously much relieved that the Prophet had not indeed paid them a visit. I attended to the infusion, readjusted my turban, then staggered off into the sand-clogged dark. It took me ten minutes to find my way back to the shelter of my house.—IAN CROSS (medical officer for Save the Children Fund, Gorom Gorom, Upper Volta).

Biology tutorial in Chinese

Mr Wong was the pastor of a new village church in Malaya during the Emergency. His life was secure and his church vigorous and growing. He was unhappy, however: he had no son. His wife, though both dutiful and comely, produced only daughters.

Twice weekly I conducted a clinic at Mr Wong's church for village residents. My training included only the London DTM&H course and a year's study of the Hakka dialect of Chinese in Kuala Lumpur. My youthful conviction that education would banish superstition, tradition, and ignorance was strong. The anguish of Mr Wong and the shame felt by his wife were increasingly obvious to me at each clinic visit.

Mrs Wong became pregnant a fifth time. Hopes rose, Mr Wong treated his wife with kindness, and she blossomed. Village gossip about the pregnancy was endless. Wagers were made and the odds shifted to favour a son. After all, men conjectured, Mrs Wong could not disappoint her husband five times in a row. Hopes were dashed for all when Siak Mi, a daughter, was born. Mr Wong's anguish was profound, his wife's depression abject. Believing I could ease the anguish, cure the depression, and gain parental love for Siak Mi simply by educating Mr Wong, I asked to speak privately with him.

After tea on his veranda, I attempted to explain the biology of human reproduction. Starting at ovum and sperm, with which he seemed vaguely familiar, I found Chinese words to describe fertilisation. His interest was kindled and his questions keen. As we proceeded to meiosis, he became animated and fascinated with the drawings I made on blank spaces of his newspaper. When we had finished with the X and Y chromosomes it was patently obvious he understood that the father determines the sex of his offspring.

Our eyes met and, in the space of perhaps five seconds, science contended with culture and history in his brain. Mr Wong's face hardened, his smile evaporated, and he rose from the table. Thanking me for explaining how the sex of Caucasian babies was determined, he entered his house to scold Mrs Wong about her baby's crying.

Soon the Wong family moved to another town and I never saw them again. I later learned that Mrs Wong produced a sixth daughter. Often I have reflected on my biology tutorial with Mr Wong. He taught me a great lesson.—F F HOLMES (professor of medicine, University of Kansas).