the least inflammatory or other morbid action. This fact also makes me object to the fourth conclusion, that "the operation definitively would seem to be indicated only under the following circumstances: the certain and complete mobility of the foreign body within the joint; the persistence of the accidents caused by its presence—viz., pain, arthritis or hyarthrosis, and lameness; the failure of compression and other means for fixing the condition of the body; and the free consent of the patient."

It is true that it is necessary to be certain of the complete mobility of the foreign body within the joint. Many years ago I saw an operation attempted on a case of the description to which M. Larrey alludes, and death was the result. Again, no discreet surgeon would operate unless the symptoms incident to the presence of a foreign body in the knee-joint were frequent or permanent. Still I must express my decided opinion that in every case, unless very exceptional, the foreign body should be removed by the form of subcutaneous incision which I advocate.

Original Communications.

MEDICAL PSYCHOLOGY.

By ROBERT DUNN, F.R.C.S.E.

III. On the Psychological Phenomena or Symptoms of Disease.

[Continued from page 522.]

Loss of Speech or the Power of Utterance. Among the abnormal phenomena of the intellectual consciousness, we may now turn from the noisy ravings of delirium, in febrile and other diseases, to the consideration of an opposite state, depending upon other and varying pathological conditions—that of dumb silence; not, however, of insensibility and coma, in which the phenomena of life are reduced to a mere series of automatic movements; but to that which results from the loss of articulate speech—a lack of the power to give utterance to the thoughts, feelings, and emotions of the mind. Now, the loss of speech or power of utterance may arise from a variety of causes, and may be quite independent of any cerebral lesion, or implication of the intellectual consciousness. For, aphonia is not unfrequently the consequence of disease of the larynx or vocal organs; at other times, the loss of voice is occasioned by pressure upon, or change of structure in, the lingual and glosso-pharyngeal nerves; either at their origin or in their course. From the close and intimate connection between the corpora olivaria and the nuclei of the hypoglossal nerves, Schröder van der Kolk has been led to infer that the former are auxiliary ganglia to the latter, in the production of certain combinations of movements in the acts of articulation and speech. "For," says he, "speech and the articulation of words require such a multitude of peculiar motions of the tongue, and such an infinite number of varying combinations of its muscular movements, that it cannot appear strange that two auxiliary ganglia should be required for the performance of these functions." After stating his conviction that all motor nerves take their origin from certain groups of multipolar ganglionic cells, he goes on to observe, "that he is acquainted with no other part of the extra spinal cord or brain where the multipolar ganglionic cells are so densely accumulated as in the nuclei of the hypoglossal nerves"; and then adduces pathological evidence in support of his opinion that the corpora olivaria are the organs for the articulation of the voice. But he by no means considers the corpora olivaria as the organs through which the will directly

acts; for these he admits to be, chiefly, the corpora striata, any injury to which is invariably followed by paralysis; and he adds that, "when the organs of the will are disturbed, the paralysis of the tongue and of the neighbouring parts renders the utterance of words wholly impossible."

During the past year, an interesting case came under my notice. The patient was an intelligent man, a law-stationer, of temperate, regular, but sedentary habits, aged 61 years when he died in September last. In April 1858, he had an apoplectic seizure, which left him, for some time, hemiplegic on the right side; but from this he gradually, and to all appearance completely, recovered; so that, in the course of five or six months, he was again actively engaged in his business. In May 1859, he had a second attack; but this left permanent effects behind it. He never properly recovered the use of his arm or leg. He could sign his name, and get about the house with assistance. He experienced some difficulty in deglutition, and there was an impediment in his speech. A third seizure in June last year left him utterly helpless as to his arm and leg, unable to articulate a word distinctly, or so as to be understood, and only able to swallow liquids, and these with difficulty, threatening, at times, suffocation. Still to the last he was sensible, and easily moved to tears, and suffered from emotional excitement. No *post mortem* examination was permitted, which I much regretted. As there had been three distinct seizures, it would have been interesting to have noted the records which they had left behind them; and to have seen how far the corpora olivaria were implicated.

In another case, a young lady, of a delicate constitution, who had suffered from repeated rheumatic attacks, and who had, in consequence, disease of the mitral valves, under a severe attack of bronchitis, became suddenly hemiplegic and speechless. She soon recovered consciousness and sensibility, so that she perfectly understood whatever was said to her, and noticed all that was going on about her; but she was hemiplegic and without power of utterance. A few days afterwards, and at the time quite unexpectedly and suddenly, she expired.

Here, too, no examination was permitted. But, from the suddenness of the death, and of the loss of speech, how interesting and instructive it would have been to have seen the actual condition of the cerebral arteries, and the *locale* of the rupture, in reference to the corpora olivaria and striata; and to the observations and researches of Dr. Kirkes. How often, in private practice, I cannot help remarking, does it happen that we are denied the satisfaction of a *post mortem* examination in cases where we have had the opportunity of carefully watching the patients during life, and of noting every appreciable change in their symptoms, so that they have, in consequence, become invested with an interest in our minds, which nothing short of a knowledge of the pathological conditions can fully satisfy. In private practice, we have neither the appliances nor opportunities for pathological investigation, which the physicians and surgeons of our hospitals can command and do possess; but still it is gratifying, and it ought to be encouraging, to reflect what may be done, and what an Abercrombie actually did accomplish single-handed in the walks of private practice.

Now, in all cases of loss of speech or the power of utterance, which are of cerebral origin, there is involved functional derangement or structural change in the nervous apparatus of the intellectual consciousness; for "perfect speech—that is, the power of expressing our thoughts in suitable language—depends upon the due relation between the centres of volition and of intellectual action." (Dr. Todd.) I had a striking illustration of this, in the young woman's case to which I have had occasion so frequently to refer; for she continued

speechless so long as her perceptive and intellectual faculties were in abeyance. It was in a moment of strong emotional excitement that her power of utterance was suddenly regained. The sight of her mother in excessive anguish and distress forced open the flood-gates of her inner sensibility, volition was aroused, and the barrier was swept away which had so long kept her spell-bound in silence; like a disruptive discharge of the nervous force, she suddenly, though hesitatingly, ejaculated, W-h-a-t's the matter?

The faculty of speech, as an instrument of thought, is an attribute of, and, in consequence, must necessarily have its seat in "the nervous apparatus of the intellectual consciousness." A special cerebral organ has been assigned to this faculty. Gall was the first to locate it, in the anterior lobes of the brain, and his allocation has found advocates in many distinguished physiologists, Serres, Paul, Grandchamp, Bellhomme, Bouillard, and others. My own pathological investigations have led me to espouse it. It may be disputed, whether there is a special cerebral organ for articulate speech; but there can be none about language being, as an instrument of thought, an attribute of the nervous apparatus of the intellectual consciousness; and hence, its connection and close association with the anterior lobes of the brain, and with the transverse series of convolutions.

I brought this subject under the notice of the Royal Medical and Chirurgical Society, in a paper on a Case of Hemiplegia with cerebral softening, and in which loss of speech was a prominent symptom; read June 25th, 1860, and published in the *Lancet* Oct. 22nd and Nov. 2nd of the same year. The case was that of a lady aged 66 years, who had suffered from three attacks of apoplexy. The first, which occurred in Oct. 1844, seemed "congestive" in its character, and passed away without any other permanent consequence than this, that she continually used one word for another, not applying appropriate names to the things or persons she desired to signify. The second attack, in May 1847, left her permanently hemiplegic on the right side, the power of voluntary motion being completely abolished; and but little sensibility being preserved, though reflex movements could be excited in the lower extremity by tickling the sole of the foot. For the rest of her life, she remained altogether incapable of speech, not being able to say yes or no in reply to a simple question, and never getting beyond the utterance of the monosyllable *dat-dat*; and yet all her senses were intact; the motions of the tongue were free, and there was no difficulty of deglutition. She did not seem to have lost any of her intellectual powers; but her emotional sensibility was certainly increased. Her general health continued good up to the time of the last fatal seizure, which occurred in April 1850, without any premonitory symptoms.

At the *post mortem* examination, the upper two-thirds of the anterior lobe of the left hemisphere was found to be in a state of complete destruction, with colourless softening; while the middle and posterior lobes were sound and healthy; but the greatest change was in the ganglionic masses at their base, and in the commissural structure. The upper half of the corpus striatum on the left side was destroyed by softening; the optic thalamus was shrunken to less than half its natural size, its upper surface being greatly wasted; while, on the right side, a small and recent apoplectic clot was seen on the upper and anterior surface of the corpus striatum, the whole of the upper half of which was in a state of *ramollissement*; while on the upper surface of the thalamus also were noticed some indications of white softening. The corpus callosum was destroyed, except at its anterior and inferior reflexion, and the anterior commissure and fornix were gone. Microscopic examination of the softened parts presented an abundance of compound cells and of fatty matter with capillaries. Now, in this case, it was quite evident that, with the disorgan-

isation of the left anterior lobe, its functional power was entirely abolished; and, although the right hemisphere was healthy, and there was every evidence, from the history of the case, that it maintained and exercised a functional power, as a centre of intellectual action, still the volitional agency was wanting to give utterance to the passing thought, for the corpus striatum on the same side was not in its integrity.

In this paper, I observed, "It is never to be forgotten that the power of giving utterance to our thoughts and ideas, in appropriate language, depends upon the due relation, between the centres of intellectual action, and the encephalic motor centres through which the volitional power is exercised in speech. Thoughts and ideas may be moulded for expression in the seat of intellectual action; but the due agency of the volitional power, to give them utterance, requires the integrity of the commissural fibres, and of the motor centres, through which the volitional impulses operate in speech."

The imperfect power of articulation, to which we have our attention so constantly called, in hemiplegic patients, there can be little doubt is most generally due to some structural lesion in the motor centres of volition, that is in the corpora striata, or among their commissural fan-like fibres which radiate from them to the cerebral hemispheres.

There is not, I believe, a single instance upon record, in which the power of utterance was retained intact, however sound and healthy the great hemispherical ganglia may have been found to be, where both the corpora striata were diseased. And thus, the conflicting evidence which has been adduced as to the seat of the faculty of speech admits of a satisfactory explanation; for, as we have seen, the loss of speech, when of cerebral origin, may alike result from disease of the anterior lobes; that is, in the nervous apparatus of the intellectual consciousness; or of the corpora striata; or their commissural radiating fibres, the motor centres of volition.

I could give many illustrative instances of the sudden loss or perversion of speech which have come under my notice, as the effects of shock to the nervous system, from a want of co-ordination in the action of the nervous centres of intellectual action, and of emotion or volition; in other words, an interruption to the pre-established harmony which should obtain between the subjective intelligence and the organs of speech.

I have lately had under my care, a married woman of the nervous temperament, and great emotional susceptibility, the mother of a large family; and who, during the latter months of her last pregnancy, met with a sudden and painful nervous shock, the effect of which was to deprive her of the power of speech, and to produce giddiness and confusion of mind. When I first saw her, some hours afterwards, she had recovered the power of articulation, and regained the integrity of her reasoning faculties; she knew every one about her, and was perfectly sensible to what was going on around her; but had lost the memory of the names of those about her, and of words. She could not recollect or give the name of the commonest article of household furniture, as a chair or table, etc.; and, sensible of her inability, she frequently burst into tears. This state continued for some time after her *accouchement*. One evening, after having heard a stirring sermon from Dr. Cumming, which had greatly aroused her emotional sensibility, she was seized, on getting home, with a severe and well marked epileptic fit, of long duration, which proved salutary; for since this disruptive discharge of the nervous force, there has been a marked improvement, both mental and bodily, in her general health.

To another instance, of which I have given an account in my *Essay on Physiological Psychology*, I would briefly advert, as the lady died afterwards in an apoplectic seizure; but no *post mortem* examination could be

obtained. In her case, a day or two after the seizure in the street, the perceptive and thinking powers were regained. She knew where she was, and all the family about her, as well as myself; but the memory of words was for some time in abeyance. She could not recollect the name even of her own daughter, who was constantly with her. She had a perfect recollection of past circumstances and events up to the time of her seizure; understood whatever was said to her; felt deeply conscious of her own inability to recollect names and common words when talking, and gave expression, in consequence, to emotional distress or feeling in tears. As I have elsewhere observed, "in this case, it may be fairly inferred, that the sudden shock to the nervous system in the first instance deranged the organic actions and normal correlations of the emotional and intellectual centres. The delirium was of short continuance, coherence of mind was soon regained; and the powers of thinking and reasoning were gradually though slowly restored. But there long remained—indeed, up to the time of the death—a manifest dislocation of the memory of words, to use an expressive term of Sir Henry Holland, on the slightest emotional excitement or mental agitation. Such attacks are but too often the precursors of apoplexy, or of serious disease of the brain. She died of an apoplectic seizure.

Progress of Medical Science.

ASTHMA PRODUCED BY PRESSURE ON THE SUPERIOR VENA CAVA. A woman, 45 years of age, of moderately robust constitution, was admitted into the Hospital La Charité on the 2nd February, 1862. The principal symptom which she presented was a considerable difficulty in respiration, manifested by the frequent elevation of the ribs. The face was of a violet hue, and had a general expression of suffering. When asked to point out the spot where she felt pain, she referred to the cardiac region. The first point which attracted the attention of the physician was a considerable dilatation of the venous branches, which, starting from the subcutaneous thoracic region, proceeded towards the left and even the right subclavian. The jugular veins externally (anterior, and internal) and the veins of the head were also distended, and did not diminish in volume after repeated expirations, any more than they increased in size when the patient diminished considerably the rapidity of her respiration—a circumstance which, according to Professor Piorry, proved that the obstacle to the flow of blood was situated in the circulating organs, and not in the lungs. Application of the hand to the thoracic parietes caused no pain, but a purring thrill was perceptible. The extent of the cardiac dulness, as determined by the pleximeter, and traced with a pencil, was 5.6 inches from side to side. The form of the organ was regular. Above the heart and aorta, and a little to the left, was a very dull space of an oval form, measuring 2.8 inches from above downwards, and 2 inches from side to side. The arch of the aorta was not dilated. Under the influence of repeated deepened expirations, the heart became somewhat diminished in size, whilst the size of the tumour did not alter. No abnormal sound of any kind was heard in the cardiac region. The liver was of the normal size; it diminished somewhat when the respirations were accelerated. Percussion showed that the thyroid gland was increased in size, and that inferiorly it was continuous with the tumour. Questions addressed to the patient, and a critical examination of her symptoms, led to the following result. The patient came from Dijon; she knew many persons in that locality who were the subjects of goitre; she herself had suffered from it in her infancy. Her health was good

till three months before admission. Her symptoms increased slowly, but incessantly; and her chief source of suffering was difficulty of breathing. Professor Piorry's diagnosis was, *Asthma, caused by the pressure of an intrathoracic tumour upon the superior vena cava.* Was the tumour an aneurism, or a tuberculous or cancerous mass? or was it a tumour of the thyroid gland? Whatever it was, it compressed the vena cava superior; and hence the dilatation of the veins, the stasis of the blood, the difficulty in breathing, etc. Professor Piorry, however, had several times seen the thyroid body extend beyond the mediastinum, so as to compress the aorta and vena cava. The patient died three days after her admission to the hospital; and the diagnosis of M. Piorry was verified by the results of the *post mortem* examination. The superior vena cava was compressed by a tumour consisting of the hypertrophied and cancerous thyroid gland. (*Gaz. des Hôp.*, and *Ed. Med. Journal.*)

LOSS OF AN EYE FROM THE BITE OF A LEECH. Professor Von Graefe relates, in the second part of the seventh volume of the *Archiv für Ophthalmologie*, the case of a delicate little girl, of five years of age, who, some days before he saw her, had on account of headache been ordered a leech to the right temple. The eye itself before this application had been quite healthy, and its vision perfect. He found the interior chamber filled with blood; the lower half of the conjunctival sac, including the edge of the cornea, ecchymosed; in the lower half of the cornea, nearly a line from the limbus conjunctivæ, a wound, the edges of which and neighbourhood were somewhat swollen, and presented a greyish opacity. The form of the wound, on close inspection, left no doubt that it was the result of the bite of a leech. It came out that, in making the application ordered, the leech had slipped away, had crept into the eye, and there, under considerable pain to the child, had accomplished its work of destruction. Unfortunately, perception of light was completely lost, apparently in consequence of total separation of the retina by hæmorrhagy. As he had never in such a case of hæmophthalmos, from whatever cause arising, observed a restoration of sight when once quantitative perception of light had for more than some hours been lost, he gave the most unfavourable prognosis. A few times afterwards, the blood in the anterior chamber had partly disappeared, the eyeball was already boggy, and he had no doubt that phthisis oculi had commenced. Dr. Von Graefe observes that the case, while it serves as a warning against the careless application of leeches in the vicinity of the eye, is interesting in another respect, that the bite was not within the extent of the conjunctiva, but nearly a line from it, and in a part which in the normal state is deemed destitute of vessels. How far the leech filled itself with blood, and how long it sucked, he could not determine. The conjunctival ecchymosis terminated abruptly at the limbus conjunctivæ, from whence, in the direction of the wound, there was only a non-vascular infiltration. It is also possible, he observes, that no blood escaped by the wound, but that the strong suction alone caused the intra-ocular and extra-ocular hæmorrhage. (*Glasgow Medical Journal.*)

POISONING BY APPLICATION OF STRYCHNIA TO THE PUNCTA LACRYMALIA. In the *American Medical Monthly* for December, Dr. Charles Schuler has an article upon the above subject. In a case of amaurosis he applied strychnia by Langenbeck's method, and the most alarming symptoms of poisoning supervened. He concludes his article by calling attention to the bearing of this case upon medical jurisprudence. 1. From five to fifteen *centigrammes*, either of pure strychnia or one of the salts of strychnia placed on the internal angle of the eye of a sleeping man, would be sufficient to destroy life speedily and silently. 2. The detection of the