vascular mechanism, which latter is also an important protective mechanism. The only favourable feature has been that the patient is apparently protected in some way from the absorption of toxic products. Whether this is due to the disturbance of vascularity or to flavine rendering toxic products non-toxic has not been determined. But the employment of flavine and other dyes as if they supplied royal roads to success must be regarded as retrograde steps. However promising this line of inquiry may be for the future, the introduction of fresh substances will only be justified after some such adequate comparative trial as we have done our best to carry out, holding no brief for any particular form of treatment.

CONCLUSIONS.

In the light of the foregoing facts the following points stand out clearly:

The flavine treatment of wounds is associated with—

1. Small formation of pus.
2. Slow epithelial ingrowth.
3. Delay in all the processes of repair.
4. Lingering of organisms on the wound surface.
5. Some diminution in the local and general reaction to infection.

In conclusion, we wish to acknowledge our great indebtedness to Surgeon-General Sir George Makins, to whose foresight and influence the surgical observation hut owes its existence, and to Sir Arthur Sloggett, Director-General of Medical Services in France, for according the necessary facilities for these investigations, as well as to the British Red Cross Society for provision for them. We have also to acknowledge the willingness with which Colonel H. E. Cree, A.M.S., assented to the necessary arrangements being made for the surgical observation hut attached to the general hospital under his command.

REFERENCES.

- ¹ Ligat, Clinical Notes, BRITISH MEDICAL JOURNAL, January 20th, 1917; Pilcher, Clinical Notes, *Journal of the Royal Army Medical Corps*, March, 1917. ² M.S. Report on use of flavine in greater solution than 1 in 1,000, by McNee. ³ Carrel and Dehelly, *Le traitement des plaies infectées*, p. 142. ⁴ Flemming, *Lancet*, September 1st, 1917. ⁵ Bashford, *British Journal of Surgery*, January, April, 1917.

THE TREATMENT OF PARAPLEGIA FROM GUNSHOT OR OTHER INJURIES OF THE SPINAL CORD.

BY

COLONEL A. W. MAYO-ROBSON, A.M.S., C.V.O., C.B.,
D.Sc., F.R.C.S.,

CONSULTING SURGEON TO THE SOUTHERN COMMAND.

NOTHING can be more deplorable than the condition of an otherwise healthy man suffering not only from paralysis of the lower extremities, but also of the sphincters of the bladder and rectum.

At present the condition is considered incurable and the patient is condemned to a life of helplessness and misery. But is it incurable? I trust to be able to show that there may be hope for some of these cases by surgical treatment.

The spinal cord is made up of a series of local nerve centres, together with a large amount of conducting material resembling nerve structure which conveys impressions to and from the brain. If continuity is lost, all voluntary action of the nerve centres below the seat of injury is destroyed.

If we can by operative means remedy paralysis due to nerve destruction by transplanting new nerve tissue or part of the spinal cord of one of the lower animals, as I proved possible some years ago,¹ why should we not be able to obtain return of function in a damaged spinal cord by excising the injured section and transplanting into the gap thus made a portion of the spinal cord of a recently killed rabbit, or perhaps better, of a sheep or calf?

I have for some time believed it to be feasible, and I now deliberately propose that such an operation should be tried in cases otherwise incurable.

With due care in asepsis it should be possible to perform the operation without danger to life; and as the paralysis in any suitable case would be complete, the condition of the patient should not be made worse by the operation,

even if it failed to establish continuity. In a case of simple fracture of the spine it may be done as a primary operation, when it is clearly established that the cord is divided or damaged beyond natural repair. If the cord is simply divided, possibly simple suture is all that may be required.

One of the following cases proves the possibility of successful suture, but in the case of gunshot wound the injury will be septic, and it is necessary that the wound should be healed and all trace of septicity absent before the operation I propose can be safely done.

Some years ago I performed laminectomy in the mid-dorsal region in a boy suffering from tuberculous disease of the spine and paraplegia. He ultimately made a complete recovery, and in due course returned to school, he being able to walk, and ultimately even to run about. Unfortunately the spinal support with which he had been furnished was, after a time, discarded, of course without advice, and in time he developed a well marked hunchback, which his fellow schoolboys considered it good fun to ride on. One day, while giving a boy a ride on his back, he fell and fractured his spine at the seat of weakness. This was immediately followed by paralysis.

I saw him shortly after the accident, and cut down on the cord, which I found torn nearly across from behind forwards, the separation behind being in width about half an inch, a small part of the front of the cord only holding together. On straightening the spine the divided ends came into good apposition, and were fixed with fine catgut sutures. The theca of the cord was then closed by a continuous catgut suture, and the rest of the wound was united. The patient was put up at once in plaster-of-Paris in the straight position. There was very little shock, and recovery from the operation was uninterrupted. The function of the lower part of the cord gradually returned, and ultimately he regained power over the sphincters, and recovered the use of his lower extremities.

This case proves that suture of the severed cord can be successfully performed.

During the past six months it has been my duty as well as my pleasure to visit officially the Duchess of Connaught's Canadian Red Cross Hospital at Clivedon, and through the kindness of Colonel Mewburn I have seen a case there that bears out the ideas I had previously held as to the possibility of transplantation to secure continuity of the divided spinal cord.

Acting on the knowledge that I had successfully transplanted the spinal cord of a rabbit into the human subject, the surgeons in charge of a case of gunshot injury of the spine with wound of the cord and consequent paraplegia have performed a most successful operation. The spinal cord was partly divided and otherwise seriously damaged over a limited area by a bullet which had lodged in the spinal canal. The bullet was removed, and after the wound had become germ-free, the edges of the injured part of the cord were pared and a section of the recently removed spinal cord of a rabbit was transplanted into the partly divided cord of the patient.

I have seen the patient on two occasions since the operation, and at my last visit the paraplegia was gradually passing off; function showed signs of returning in the lower extremities and the sphincters were regaining power—in fact, the case promises to be a brilliant success.*

Colonel Mewburn has kindly forwarded me an account of the condition of the patient on November 24th, 1917. Sensation has returned from above the nipple line to 3½ in. below that line. There is a general soreness over the rest of the paralysed parts indicating the beginning of return of sensation in the paralysed parts. There is a marked tendon patella reflex, and both legs have sensation when being tapped. Both limbs give a marked continuous clonus. The patient says that he is conscious when his bowels are moving and when the parts are soiled. Bed-sores are healing, and some have quite healed. All the improvements have dated from the time of operation two months ago.

These cases have encouraged me to write this short paper so that others may give a trial to the method I have advocated and so make an attempt to cure some of the deplorable cases of paraplegia up to the present time considered to be incurable.

I have just seen in one of the hospitals of the Southern Command a case of fracture of the seventh dorsal vertebra from gunshot injury in April of this year. The wound has healed soundly and the patient is in good general health except that he is completely paraplegic, including loss of use in the sphincters of the bladder and rectum. The condition of the bladder has been relieved by the

* This case will doubtless be reported later with full details by the surgeons in charge.

performance of a suprapubic cystotomy. I have advised operation, and I hope shortly to help my colleague, under whose care the man is, to perform laminectomy to be followed by excision of the damaged area of the cord and transplantation of one or more sections of a rabbit's cord into the gap, in order to try to secure conduction of impressions between the upper and lower segments of the man's spinal cord.

There are, I feel sure, many similar cases of invalided soldiers in various institutions throughout the country who would doubtless be willing to undergo operation even if it gives the barest possibility of success.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, September 13th, 1890, and October 31st, 1896.

An Address

ON THE

NECESSITY FOR EDUCATION IN RADIOLOGY
AND ELECTRO-THERAPEUTICS.

DELIVERED TO THE ELECTRO-THERAPEUTICAL SECTION OF
THE ROYAL SOCIETY OF MEDICINE.

BY

G. HARRISON ORTON, M.A., M.D.,

PRESIDENT, ELECTRO-THERAPEUTIC SECTION, ROYAL SOCIETY OF
MEDICINE; RADIOLOGIST TO ST. MARY'S HOSPITAL.

(Abstract.)

DR. ORTON began by referring to the views expressed by Dr. Thurstan Holland in his presidential address to the Röntgen Society,* and in his letter in the JOURNAL of September 15th, 1917, giving an account of the steps taken by the Surgeon-General of the United States Army to organize military radiographic work; as a result of a conference of leading radiologists a number of schools in charge of experts were established for the teaching of radiographic work, which medical officers intending to do x-ray work were required to attend for three months; the x-ray examination of recruits, especially with reference to chest conditions, was organized, and methods for the localization of foreign bodies were considered.

Dr. Orton said that the organization for military x-ray work had been carried out in this country without consultation with leading radiologists, with possibly one exception, and added that the same remark applied to electro-therapeutics. In March, 1917, the council of the Electro-therapeutic Section of the Royal Society of Medicine appointed a small committee, with Sir James Mackenzie as chairman, to consider the best manner in which to deal with questions concerning the status of medical men holding positions as radiologists and electro-therapeutists, and the steps which should be taken for the promotion and teaching of radiology and electro-therapeutics. After full consideration had been given to the matter at several meetings the following letter was addressed to the Vice-Chancellor of the University of Cambridge on May 26th, 1917:

Dear Sir,

The call for some organization of the members of the medical profession who practise radiology and other branches of electro-therapeutics is urgent, and there is great need that some steps should be taken to establish these subjects on a proper footing both in regard to the public and the profession generally. At present no standard exists; there is no organized teaching of the subjects, and those medical men who have taken up this branch of medicine have had to pick up their knowledge almost entirely by experience—at the expense of their patients. The result of this haphazard state of things is well depicted by Captain Thurstan Holland in his presidential address before the Röntgen Society—the lack of knowledge on the part of those who have been responsible for placing totally untrained men, both lay and medical, in charge of most of the military x-ray departments has led to endless tragedies, while in civil practice much suffering would be saved if competent radiologists were in charge of the departments. In civil general hospitals,

where the x-ray department is properly organized, fully 10 per cent. of all the patients who enter the doors of the hospital find their way to this department, and this in spite of the fact that those in charge are so over-burdened with routine work that there is no time for research or the undertaking of new branches of diagnosis and treatment.

As the outcome of a discussion at the Council Meeting of the Electro-therapeutic Section of the Royal Society of Medicine on March 16th of this year a small committee was formed to consider—

(i) The best means of raising the status of radiology and electro-therapeutics to their rightful place among the branches of medicine, and to attract the better class of medical men to take up these special subjects.

(ii) To suggest what steps should be taken to provide teaching in radiology and electro-therapeutics.

At this meeting, after a prolonged discussion, a sub-committee was elected to go into this matter. After considering several schemes the subcommittee suggested that it was not advisable to start any programme of teaching in London until it was known if facilities could be offered for granting a diploma at Cambridge or some other university.

It was decided to approach Cambridge University in the matter, as it was stated by one of the members that funds would be found not only to provide for the necessary expenses, but also to equip, maintain, and eventually to endow a Research Hospital and Chair if these facilities were granted by this University.

It is intended that the scheme shall be started on broad lines, in order that development may not be hampered, the objects being:

(i) To provide a standard in these subjects.

(ii) To provide the teaching that will be necessary, together with the facilities for hospital experience.

So far as it has advanced at present the scheme of the Subcommittee provides for:

A. A central organization in London, housed in an institute, where teaching facilities would be available in the scientific part of the curriculum, the practical experience being available in the large hospitals. Later, if necessary, centres could be started in other towns.

B. A post-graduate diploma in one of the older universities where a research hospital would be built. This hospital would be devoted to the scientific investigation of the many physiological, pathological, and physical problems that are waiting to be worked out. This hospital would be available for those who wished to work on various problems, and especially for those who were investigating with a view to the M.D. thesis.

Appended you will find outlines of: A, The London Scheme; B, the Cambridge Scheme:

A. The London Scheme.

A representative board, elected annually, would be responsible for this scheme.

1. The intention is to provide a central institute in London, equipped for general organization, and also for post-graduate teaching, where the necessary appliances would be available for demonstrations, and where all lectures on scientific subjects could be given.

2. Arrangements would be made for teaching in physics, chemistry, pathology, and clinical medicine, and for lectures by specially qualified medical men in those branches of radiology in which they had specialized. Special lectures and demonstrations in allied subjects would also be arranged, and whenever distinguished colleagues from abroad visited this country they would be invited to deliver lectures.

3. In conjunction with the university which grants the diploma, the board would draw up the general lines of teaching, not only for the London school, but for any other schools that might be formed.

4. The board would arrange for students to hold appointments as clinical assistants in the various x-ray and electrical departments that would be available in the London hospitals.

5. The board would arrange centres throughout the country in close touch with the central institute, where it would be possible to arrange for post-graduate teaching.

6. The institute would be the centre for the spread of knowledge of these subjects and of the work done in this country; and, if possible, the existing x-ray journal would be utilized as its official organ.

In the future it is probable that many overseas workers will be attracted to this country, and a strong

* Journal of the Röntgen Society, January, 1917, vol. xiii, No. 50. BRITISH MEDICAL JOURNAL, November 18th, 1916, p. 696. A fuller expression of Dr. Holland's view was contained in an address published in the JOURNAL of March 3rd, 1917, p. 285.