Summary

An account is given of five former professional boxers, four of whom developed a chronic cerebral disorder in later life.

Two (Cases 1 and 5) were both locally considered to be punch-drunk. However, the former had an absent septum pellucidum and the other was an alcoholic. Necropsy in the latter showed foci of softening in the cerebral and cerebellar cortex.

Two (Cases 2 and 3) suffered from cerebral atrophy. Both had cavum septi pellucidi, and in one (Case 2) the leaves of the septum seemed to be perforated.

One (Case 4) was not physically disabled, but his behaviour was violent, his cerebrospinal fluid protein was raised (60 mg./100 ml.), and air encephalography suggested that there was a perforation in his septum pellucidum.

The possible relationship between these lesions and boxing is discussed. There is a paucity of pathological observations of the so-called punch-drunk syndrome, but before the existence of the syndrome is denied, as in some quarters it has been, follow-up studies of former professional boxers will be necessary.

I thank Dr. A. S. Bligh, Department of Radiology, Cardiff Royal Infirmary, for his co-operation and advice. Dr. James Bull, the National Hospital, Queen Square, kindly offered suggestions and helpful criticisms. Dr. William Phillips invited me to investigate Case 3, and Dr. E. Payne, Department of Pathology, Welsh National School of Medicine, examined the brain of Case 5. The necropsy on the latter was performed by Dr. G. S. Andrews, St. Woolos Hospital, Newport, Mon, who sent me the brain.

REFERENCES

- Basu, B. N. (1935). *J. Anat. (Lond.)*, **69**, 394.
 Bell, W. E., and Summers, T. B. (1958). *Neurology*, **8**, 234.
 Brandenburg, W., and Hallervorden, J. (1954). *Virchows Arch. path. Anat.*, **325**, 680.
 Critchley, Macdonald (1957). *Brit. med. J.*, **1**, 357.
 Dandy, W. (1931). *Arch. Neurol. Psychiat. (Chic.)*, **25**, 44.
 Davidoff, L. M., and Epstein, B. S. (1955). *The Abnormal Pneumoencephalogram*, 2nd ed., p. 371. Kimpton, London.
 Dolgop, V. B. (1938). *Arch. Neurol. Psychiat. (Chic.)*, **40**, 1244.
 Dyke, C. G., and Davidoff, L. M. (1935). *Amer. J. Roentgenol.*, **34**, 573.
 Forster, E. (1933). *Rev. neurol.*, **6**, 1122.
 Gibson, J. K. (1924). *Anat. Rec.*, **28**, 103.
 Grahmann, H., and Ule, G. (1957). *Psychiat. et Neurol. (Basel)*, **134**, 261.
 Hahn, O., and Kuhlbeck, H. (1930). *Fortschr. Röntgenstr.*, **41**, 737.
 Hochstetter, F., and Sitzber, B. (1925). *Akad. Wiss. Wien*, **132**, 1.
 Holbourn, A. H. S. (1943). *Lancet*, **2**, 438.
 Kaplan, H. A., and Browder, J. (1954). *J. Amer. med. Ass.*, **156**, 1138.
 McCown, I. A. (1959a). *Ibid.*, **169**, 1409.
 — (1959b). *Amer. J. Surg.*, **98**, 509.
 Martland, H. S. (1928). *J. Amer. med. Ass.*, **91**, 1103.
 Neuberger, K. T., Sinton, D. W., and Denst, J. (1959). *Arch. Neurol. Psychiat. (Chic.)*, **81**, 403.
 Pudenz, R. H., and Shelden, C. H. (1946). *J. Neurosurg.*, **3**, 487.
 Savain, W. P. (1946). *Radiology*, **46**, 270.
 Schwidde, J. T. (1952). *Arch. Neurol. Psychiat. (Chic.)*, **67**, 625.
 Sintesco, S., and Mihailesco, N. (1938). *Bull. Soc. Psychiat. Bucharest*, **3**, 53.
 Tenchini, L. (1880). *Boll. Sci. med.*, **2**, 65.
 Victor, M., Adams, R. D., and Mancall, E. L. (1959). *Arch. Neurol. (Chic.)*, **1**, 579.
 Vinken, P. J., and Strackee-Kuijer, A. (1957). *Folia psychiat. neerl.*, **60**, 226.

The Research Defence Society (11 Chandos Street, Cavendish Square, London W.1) has published No. 16 in its Conquest Pamphlet Series, *The Use of Laboratory Animals in Assessing the Safety of Food Additives*. (Price 6d.)

STUDIES OF THE RETINAL CIRCULATION WITH FLUORESCIN*

BY

C. T. DOLLERY, M.B., M.R.C.P.

J. V. HODGE,† M.D., M.R.A.C.P.

AND

MORAG ENGEL, S.R.N., A.R.P.S.

From the Department of Medicine, Postgraduate Medical School, Hammersmith Hospital, London

[WITH SPECIAL PLATE]

The retinal circulation is unique in being the only part of the vascular system available for direct inspection. Observation of the retina in systemic diseases like hypertension and diabetes, in which the small blood vessels are liable to damage, has become a widely accepted routine. Much can be learned from clinical ophthalmoscopy, which reveals features characteristic of various diseases affecting the retina and its circulation. Finer details have been revealed by pathological studies, particularly post-mortem injection of the retinal vessels (Ashton, 1951) and digestion studies (Kuwabara and Cogan, 1960). These techniques have demonstrated abnormalities in vessels too small to be observed during life, and have provided new information concerning the pathogenesis of exudative lesions of the retina.

A new technique has been described by Novotny and Alvis (1961) which promises to reveal during life some of the fine details previously seen only in necropsy studies. A retinal camera was modified to photograph the passage of intravenously injected fluorescein through the retinal vessels. This method shows details of the blood flow through the retinal vessels, including variable rates of flow in different vessels, and also shows spreading patches of fluorescein which seem to indicate increased vascular permeability in certain areas.

In this paper we report experience based on 60 studies of abnormalities of the retinal circulation using the fluorescein method modified in some minor respects.

Method

The apparatus used was basically the same as that described by Novotny and Alvis. A Zeiss retinal camera was modified for fluorescence photography by the insertion of a blue glass filter (Kodak Wratten No. 47B) in the common pathway of the incandescent and electronic flashlight beams, and a green filter (Kodak Wratten No. 58) immediately in front of the camera body. For normal photography the blue filter could be withdrawn by a control on the side of the camera, and this was also used for a quick check on the position of the eye between fluorescence photographs. This was not always necessary, as with good dark adaptation of the observer and the use of the brightest incandescent-light setting, the fundus could be visualized satisfactorily by blue light alone.

Fluorescence photographs were taken on Ilford HPS film (ASA rating 800) which was force-developed in

*Work supported by the Medical Research Council.

†In receipt of grants from the Nuffield Foundation and from the Medical Research Council of New Zealand.

Kodak D76 developer for 25 minutes at 75° F. (23.9° C.). Half-plate prints were made from these negatives for preliminary study. For comparative purposes colour photographs of the retina using Kodachrome film (ASA rating 10) were taken on each patient prior to the injection of fluorescein. Colour and fluorescence photographs were magnified to the same degree and carefully superimposed in order that composite drawings could be prepared. By this means accurate location of lesions in both types of photograph was possible. Maximum dilatation of the pupil was needed for both colour and black-and-white photography, and this was achieved by using 0.5% cyclopentolate and 10% phenylephrine eye-drops instilled 20 to 30 minutes before the examination.

The injections were given through a small "polythene" catheter, which was usually inserted in the ward before the patient came to the retinal-camera room. The skin was anaesthetized and a suitable thin-walled needle was inserted into a median antecubital vein. Approximately 50 cm. of thin polythene (P.E.60) was passed through the needle in a central direction so that the tip lay in the region of the superior vena cava. The intravenous needle was then removed, the end of the polythene tubing was attached to a syringe containing heparinized saline, and both tube and syringe were taped to the forearm. This method took only slightly longer than an ordinary venepuncture and had several advantages. The arm could be moved freely with the venous catheter in place, as there was no risk of its being dislodged or kinked by movement, and repeated injections could be given without discomfort to the patient. Additionally, the more central site of injection allowed a more compact bolus of dye to reach the retinal vessels and increased the intensity of the fluorescence.

As might be expected, a fast circulation time favoured a more compact "dye curve" and better-quality photographs; a slow circulation time (as in heart failure) or other circulatory abnormalities (as in valvular incompetence or intracardiac shunts) produced a dispersed dye curve and weak fluorescence over a longer period. Results were particularly favourable in patients with severe anaemia because of the short circulation time and less absorption of the fluorescence by the blood. The catheter-to-tongue circulation-time was measured with saccharin solution before injection of fluorescein. This measurement was not essential, but it did give an indication of the appearance time of the dye and of the quality of the photographs that might be expected.

Usually three injections were made at one examination, each consisting of 5 ml. of a 5% solution of fluorescein obtained commercially ("fluorescite") or more recently prepared and sterilized in the hospital pharmacy. Photographs were taken at an appropriate interval after the first appearance of the dye in the retinal vessels and thereafter at intervals of 14 to 16 seconds, which was the time required for the recharge of the electronic flash. Since most of the dye had passed through the retina before a second photograph could be taken, the timing of the first was deliberately varied in order to study different phases of the retinal circulation. The whole examination was carried out as rapidly as possible, since increasing fluorescence of the optic media interfered with definition of photographs taken after 10 minutes or so from the first injection.

Fate of Injected Fluorescein

The injection of three 250-mg. doses of fluorescein causes a yellowish discoloration of the skin, and patients were warned to expect this. The dye is rapidly excreted in the urine, which becomes a bright yellow colour during the first few hours after injection. The discoloration of the skin fades in 6 to 12 hours, but the urine remains slightly fluorescent for 24 to 36 hours. The rate of disappearance of plasma and urine fluorescence in one patient following a dose of 1,000 mg. is shown in Fig. A.

FIG. A.—Semilogarithmic plot of fall in plasma concentration and urinary excretion of fluorescein after a dose of 1,000 mg. intravenously. Patient aged 50 with normal renal function.

The rate of loss of fluorescein from the plasma is rapid even in the first few minutes after injection while photographs are being taken. This may be important in the understanding of the mechanisms whereby some retinal lesions become brightly fluorescent. The rate of loss from the plasma was observed in one patient by injecting intravenously 375 mg. of fluorescein mixed with 10 μ c. of 131 I-labelled albumin. Blood samples were taken at frequent intervals for 20 minutes, and the concentration ratio of fluorescein and labelled protein was compared at each time taking the ratio in the injection as unity (Fig. B). After only 10 minutes the volume of distribution of the fluorescein was four times at great as that of the 131 I.

It proved difficult to measure the concentration in the red cells because of the loss of sensitivity of the

FIG. B.—Ratio of plasma fluorescein concentration to plasma 131 I albumin concentration for 20 minutes after an intravenous injection of a mixture of the dye and labelled protein.

fluorimeter in the presence of only 1% of laked red cells. Absorption of the emitted green fluorescent light by the red haemoglobin probably reduces greatly the amount of light transmitted to the camera from the eye and partly explains the excellent results achieved in patients with anaemia.

Normal Retinal Circulation

The passage of dyed blood through the retinal vessels produces a continuously changing sequence of events conveniently described in distinct phases. These phases may, however, overlap in different parts of the retina owing to local variations in the speed of passage through the retinal circulation and individual variations in the time-concentration curve of the dye.

Early Arteriolar Phase.—Within a second or so of its first appearance at the disk the fluorescein spreads rapidly outwards in the retinal arterioles. During this phase the background of the retina and the veins remain dark (Special Plate, Fig. 1).

Late Arteriolar Phase.—The peak concentration of the dye usually reaches the arterioles two to three seconds later and the whole arteriolar tree is then brightly fluorescent. From this stage capillary and early venous filling can be seen, and many photographs show fluorescence of all major vessels (Special Plate, Fig. 2). Once capillary filling begins an irregular narrow dark zone outlines on either side of the arteriolar blood column. The dark area lies between the vessel lumen and the edge of the capillary net and appears to be mainly the thickness of the arteriolar wall. As the arterioles empty the undyed blood is seen first flowing down the central part of the large vessels near the disk, demonstrating that flow is laminar.

Capillary Phase.—Filling of the capillaries is shown by background fluorescence already well developed in the late arteriolar phase. This tends to be patchy at first but very soon becomes generalized. Often capillary filling appears as a distinctly reticular pattern conforming to the expected appearance of the capillary net. Individual capillaries can be identified on some of the best photographs.

Early Venous Phase.—During the stage of early arteriolar filling some fluorescent blood passes rapidly across the capillaries into the small veins. At the stage of late arteriolar filling fluorescein is already entering the main retinal veins from tributaries near the disk. This blood layers in as a bright line along the walls of the veins where streaming remains in many cases right up to the disk, providing confirmation of the laminar flow reported by Novotny and Alvis. Nipping of the veins at arteriovenous crossings destroys this streaming beyond the point of nipping, presumably by setting up turbulence. The earliest venous filling takes place from the macular region (Special Plate, Fig. 1) at least in part due to the shorter circulation pathway. In consequence the peripheral layers of fluorescent blood in the main superior and inferior temporal veins are usually unequal in width, the wider column being on the side of the vein nearest the macula. Where two veins showing stratified flow unite a third central column of dyed blood is formed by the coalescing of streams from adjacent walls of tributary veins (Fig. C; Special Plate, Fig. 2). Occasionally a dark band may be seen beside the lumen of the vein narrower than that seen beside the arterioles.

Late Venous Phase.—By 15 to 20 seconds from the first appearance of the dye the arterioles lose their

fluorescence and laminated flow in the veins reverses its earlier pattern—that is, the main superior and inferior temporal veins now receive more fluorescent blood from peripheral parts of the retina and show a wider band of fluorescence on the side away from the macula. Central streaming of dyed blood can also be seen in some veins. The brightness of venous fluorescence then fades but does not disappear completely, as recirculation of dyed blood allows all major vessels to be seen faintly for some time. In the normal retina most background fluorescence clears by the late venous phase except at the disk, where it persists for several hours. It is not clear whether the disk fluoresces because of a higher local concentration of the dye due to greater capillary permeability or even avidity for fluorescein, or whether dye at uniform concentration in the tissues occupies a thinner layer in the retina than at the disk. The latter explanation seems more probable.

FIG. C.—Schematic drawing of the development of laminar flow in retinal veins and its destruction at a site of arteriovenous "nipping" (see text). Drawn as from a film negative—that is, black outlines indicate bright fluorescence.

Abnormalities of Blood Vessels

Irregularities of calibre, segmental narrowing, and occlusion of vessels are shown clearly by the fluorescein technique which usually only confirms appearances seen on direct ophthalmoscopy. Abnormal twisted and tortuous vessels are seen in some diabetic fundi and occasionally in the fundi of hypertensive patients (Special Plate, Fig. 3). The extreme vascularity of retinitis proliferans shows as an immediate fluorescent blush and other areas of neovascularization as meshes of tiny vessels, many of them too small to be seen with the ophthalmoscope (Special Plate, Fig. 4). There are greatly increased numbers of small vessels on the edge of the disk if there is papilloedema (Special Plate, Fig. 9) in agreement with the findings of Ashton (1951) in necropsy material. Spreading pools of fluorescence appear round many areas of new vessels, suggesting that they are excessively permeable to fluorescein. In papilloedema the disk becomes brightly fluorescent, a finding confirmed in three patients with malignant hypertension and one with a cerebral tumour. The disk of the patient with a cerebral tumour remained fluorescent for 16 hours.

Microaneurysms appear in fluorescein photographs as sharply defined dots of varying size usually lying near small branch vessels and sometimes linked to them by minute channels. In two diabetic patients the aneurysms bulged from the walls of branch veins (Special Plate, Fig. 4). Microaneurysms fill with fluorescein with the rest of the capillary bed during the late arterial phase but they empty more slowly than the rest of the vascular bed. Superimposition of the fluorescein and colour photographs shows that the lesions visible by ophthalmoscopy are much fewer in number than those seen in the fluorescein pictures. Furthermore, many lesions con-

sidered clinically as "microaneurysms" do not fluoresce and appear only as black dots. Many of these look like small blot haemorrhages, but some may be thrombosed aneurysms. In some cases small haemorrhages lie close to true microaneurysms as though blood may leak from them.

As expected, microaneurysms are seen most frequently in the retinae of diabetic patients, but they also occur in a small proportion of hypertensive patients. Micro-

aneurysms in hypertensive patients are seen in one or more of three circumstances: (1) as isolated scattered lesions; (2) near the disc edge, associated with increased numbers of small vessels in papilloedema (Special Plate, Fig. 9); and (3) in the vicinity of cotton-wool patches, where they sometimes lie circumferentially outside the visible exudate and give the whole lesion an annular appearance on the fluorescein photograph.

A similar pattern of aneurysms in relation to "soft" exudates has been described in the eyes of patients with malignant hypertension removed after death (Ashton, 1959). Lesions of this type are shown in Special Plate, Fig. 9, and Fig. D. However, many soft exudates do not have any visible local vascular abnormality.

Several large aneurysms in an area of abnormal small vessels and typical microaneurysms were unexpected findings in a patient with benign hypertension (Special Plate, Figs. 5 and

FIG. D.—Outline drawing prepared by superimposing the photographs shown in Special Plate, Figs. 7 and 9. Microaneurysms in the fluorescence photograph are drawn as black dots, and soft exudates which were visible in the earlier colour photograph (Fig. 7) as stippled areas.

6). Background fluorescence was absent in an area enclosed by these lesions, as if the capillary bed was closed in that area. The vascular pattern resembles that described by Ashton as characteristic of soft exudates, but the retinal appearance and the relatively mild hypertension make this unlikely. The lesion may have been near an exudate which had cleared with antihypertensive treatment or it may be the result of occlusion of a branch retinal vein. Another patient who had a branch venous occlusion with anastomotic vessels had a number of microaneurysms in the territory of the obstructed vein.

Retinal Exudates

Most "soft" exudates fluoresce brightly in the first few minutes after the dye injection and some fluorescence remains in them after the first transit. Often the fluorescence spreads over a wider area than the visible exudate. Exudates were studied in seven patients with hypertension, one with pernicious anaemia, and one with subacute bacterial endocarditis. Two of the hypertensive patients were studied on five separate occasions and comparison of the photographs shows that a fluorescent area can precede a visible soft exudate and remain after the exudate clears (Special Plate, Figs. 7 to 10).

The background blush due to capillary filling is normal in many soft exudates but some remain dark (Special Plate, Figs. 11 and 12). Curiously localized bright fluorescent spots are seen in some patients with

LEGENDS TO SPECIAL PLATE: C. T. DOLLERY ET AL.

FIG. 1.—Man aged 40 with moderately severe hypertension. Fluorescence photograph taken 9 seconds from time of injection showing complete filling of arterioles with very early capillary and venous fluorescence, especially near the disk, where transit is more rapid.

FIG. 2.—Same patient as Fig. 1. Fluorescence photograph taken 12 seconds from injection. All major vessels are well outlined, laminar flow in veins can be seen, and the background is brighter because of capillary filling.

FIG. 3.—Woman aged 49 with hypertension and renal failure. Fluorescence photograph showing tortuous small vessels in the disk region arising from both arterioles and venules. Many of these were invisible by ordinary ophthalmoscopy.

FIG. 4.—Man aged 49 with a five-months history of diabetes and a positive family history. Fluorescence photographs show meshes of grossly abnormal small vessels. Moderate numbers of microaneurysms are seen, some as discrete fluorescent dots and others clustered along the course of small venules. Increased background fluorescence is already occurring in some areas of abnormal vessels.

FIG. 5.—Woman aged 57 with moderate "benign" hypertension. Reproduction of colour photograph showing a localized area of hard exudates and haemorrhages in a fundus which had only hypertensive vascular changes elsewhere.

FIG. 6.—Same patient as Fig. 5. Fluorescence photograph of same region showing saccular aneurysms, dilated venous loops, microaneurysms, and tortuous small vessels surrounding a dark area of apparent avascularity. The largest aneurysm was invisible clinically.

FIG. 7.—Woman aged 32 with malignant hypertension. Reproduction of colour photograph showing numerous exudates and haemorrhages in the vicinity of an oedematous optic disk.

FIG. 8.—Same patient as Fig. 7. Reproduction of colour photograph of the same region taken after two weeks' antihypertensive treatment. Note disappearance of soft exudates in several sites, particularly above and below the disk, and reduction in size of others.

FIG. 9.—Same patient as Fig. 7. Fluorescence photograph taken on same date as Fig. 8. Numerous microaneurysms can be seen close to the disk and in areas of soft exudate. Above the disk a circularly arranged group of microaneurysms occupies the site of a soft exudate which has cleared between the taking of Fig. 7 and Fig. 8. Increased numbers of small vessels are seen around the edge of the disk, which is already showing abnormal bright fluorescence (see also text, Fig. C).

FIG. 10.—Same patient as Fig. 7. Fluorescence photograph taken 10 minutes after Fig. 9 showing bright fluorescence of disk and soft exudates, including those above and below the disk which were no longer visible clinically.

FIG. 11.—Man aged 32 with malignant hypertension. Reproduction of colour photograph showing an area of soft exudate and haemorrhages.

FIG. 12.—Same patient as Fig. 11. Fluorescence photograph taken during first transit of dye showing bright spots from which fluorescence spread outwards subsequently. Some of these lie within or close to visible exudates. Note also the absence of background capillary filling in the region of two soft exudates which failed to show the expected bright fluorescence at a later stage (see text). Haemorrhages can be seen as black areas in this figure.

malignant hypertension before treatment is begun. These light up during the first transit of the dye and form the centre of a spreading pool of fluorescence. Comparison with colour photographs shows that these lesions sometimes lie close to a visible exudate but may occur in areas of retina which appear normal. The nature of the "bright spots" is unknown but presumably they are areas of local acute vascular damage.

"Hard" exudates in the eyes of diabetic and hypertensive patients do not fluoresce. Retinal haemorrhages obscure the passage of the dye in the underlying retina and appear as dark patches on the photographs.

Discussion

It is clear from the work of Novotny and Alvis and from our own observations that the technique of fluorescence photography can be usefully applied to the study of both normal and abnormal retinæ. Patterns of retinal blood flow, abnormalities of blood-vessels, microaneurysms, and soft exudates can all be examined by this method. The technique is relatively simple, and in our experience so far has proved entirely safe.

Confirmation has already been possible of some findings previously reported only in necropsy material. Thus the frequency of retinal microaneurysms and their association with soft exudates, papilloedema, and venous occlusion, and the increase in small vessels on the edge of the disk in papilloedema, are findings which have now been demonstrated both in pathological specimens (Ashton, 1951, 1959) and on fluorescence photography. Our studies of microaneurysms by this technique show how few of these lesions can be seen by ophthalmoscopy. Most true microaneurysms are invisible except when outlined by the fluorescein method, and many of the red dots diagnosed clinically as microaneurysms do not fill with fluorescein. Reports of the disappearance of these lesions must therefore be accepted with caution if they are based solely on ophthalmoscopic study: many of the lesions described are probably punctate haemorrhages and not microaneurysms. Routine mapping of such lesions in diabetic patients is made much more accurate by fluorescence photography, a study on which we are now engaged in co-operation with Dr. D. J. Scott.

The demonstration of microaneurysms in relation to some soft exudates fits in closely with the observation of Ashton (1959) on necropsy preparations. Many soft exudates do not show microaneurysms, and Ashton in a personal communication has confirmed that this is also true of his material.

Most of the soft exudates studied have shown abnormal fluorescence which persists for several hours after the injection of the dye. This appears either as a diffuse "blush" or as brilliant point sources of fluorescence which lie within or near the edge of exudates. In both types fluorescence spreads outwards, often extending beyond the limits of the visible exudate (Special Plate, Fig. 10). These findings may be partly explained by an increase in the capillary permeability to fluorescein in the region of the exudate, but as fluorescein normally leaves the circulation rapidly (Fig. B) this hardly explains the contrast between the area of the exudate and the surrounding retina. Other possibilities, such as an increase in the thickness of the layer of tissue fluid or even local affinity for the dye, require further study.

Some soft exudates did not fluoresce, and Novotny and Alvis also observed this. Two exudates which did

not fluoresce on the first examination did so subsequently. It is not clear why these differences should exist between lesions which appear identical on ophthalmoscopy. The age of the exudate does not appear to be an important factor, as some exudates fluoresce at all stages of their development. It has been stated that soft exudates are basically ischaemic in origin and that branch arteriolar occlusions may sometimes be found in relation to them (Friedenwald, 1952; Leishman, 1957). Circulatory changes may explain why some exudates fluoresce and others do not, because partial arteriolar occlusion might cause ischaemic damage to the capillary bed with increased permeability to fluorescein, while more complete obstruction could prevent access of fluorescein to the damaged area. However, we have not found convincing evidence of vascular changes in the vicinity of many exudates, and other explanations are possible. Simple pressure-oblation of the capillaries by a large exudate has been postulated (Ashton, 1959) and deflection of large vessels by growing exudates has been observed in the hypertensive retina (Dollery and Hill, 1961). Transmission of fluorescent light from the region of the capillary bed might be prevented by the high optical density of an unusually thick or opaque exudate. We are still studying this problem. Whatever the mechanism responsible, it is of interest that fluorescence can precede a visible lesion and persist for a time after it has cleared.

Apart from the theoretical interest of such observations the more exact documentation of retinal lesions by this technique may well be of practical importance in the future as a prognostic or even as a therapeutic guide. More work is required, and especially serial studies in suitable individual patients, in order to establish the natural history of these lesions.

Summary

A retinal camera has been modified so that it can be used to photograph the passage through the eye of intravenously injected fluorescein. The dye passes rapidly out along the arteries, through the capillaries and layers into the veins from small tributaries as individual bright streams along the outer walls. Mixing of the various streams of blood is slow except where arteriovenous nipping sets up turbulence in the vein. Laminar flow has been demonstrated in both arteries and veins. In the best photographs individual capillaries can be seen, and a thin dark line, mainly the thickness of the vessel wall, separates the edge of the capillary net from the lumen of the vessel. Vascular abnormalities which were invisible with the ophthalmoscope were demonstrated in some patients with hypertension and others with diabetes. Often such abnormal vessels appear particularly permeable to fluorescein.

Greatly increased numbers of small vessels, together with microaneurysms, have been demonstrated on the disk edge in papilloedema. The oedematous disk becomes brightly fluorescent.

Microaneurysms have also been demonstrated in areas of venous obstruction and in the vicinity of some soft exudates, as well as more diffusely scattered through the retina in diabetic patients. Most true microaneurysms are invisible clinically and can be demonstrated only by the fluorescein technique. Many lesions diagnosed clinically as microaneurysms are probably small punctate haemorrhages.

Most soft exudates become fluorescent, and in some the fluorescence precedes the visible exudate and persists after it has disappeared.

We would like to express our thanks to Professor J. McMichael, F.R.S., for his interest and encouragement, and to Professor N. Ashton, of the Institute of Ophthalmology, for valuable suggestions and criticism.

REFERENCES

- Ashton, N. (1951). *Brit. J. Ophthalm.*, **35**, 189.
 — (1959). *Lancet*, **2**, 625.
 Dollery, C. T., and Hill, D. W. (1961). In *Hypertension: Recent Advances*. Lea and Febiger, Philadelphia.
 Friedenwald, J. S. (1952). *Ophthalmic Pathology: An Atlas and Textbook*. Saunders, Philadelphia.
 Kuwabara, T., and Cogan, D. G. (1960). *Arch. Ophthalm.*, **64**, 904.
 Leishman, R. (1957). *Brit. J. Ophthalm.*, **41**, 641.
 Novotny, H. R., and Alvis, D. L. (1961). *Circulation*, **24**, 82.

SOME BIOCHEMICAL CHANGES IN THE TRANSPLANTED KIDNEY

A PRELIMINARY REPORT

BY

M. A. WILLIAMS, M.Sc.

H. M. TYLER, B.Sc.

MARY MORTON, B.Sc.

A. NEMETH,* M.D. (Szeged)

AND

W. J. DEMPSTER, F.R.C.S.

Department of Surgery, Postgraduate Medical School
of London

Homotransplanted kidneys of the dog are known to survive for several days before their function is abruptly terminated by the onset of an irreversible anuria. Possible causes of this anuria were discussed but no satisfactory explanation was put forward (Dempster, 1955), although at this time a reaction of the kidney involving a tubular metabolic upset was thought possible. Homotransplanted kidneys in general show little histological alteration from the normal apart from some interstitial oedema and a plasma-cell infiltration. Recent investigations into the source of the cellular infiltration (Fowler and West, 1960, 1961; Porter and Calne, 1960; Dempster and Williams) indicate that a variable number of these cells originate in the host and may be implicated in the rejection process.

The available data on the physiological performance of homotransplanted kidneys (Dempster, 1953a; Fedor *et al.*, 1959; Tyler *et al.*, 1962) give little indication of any pathological processes which could explain the sudden anuria.

In an attempt to elucidate further the sudden terminal events, a series of biochemical studies have been undertaken on transplanted kidneys. The respiratory enzymes succinic dehydrogenase and malic dehydrogenase and the glycosidic enzymes β -glucuronidase, β -galactosidase, and β -glucosaminidase have been measured quantitatively in normal, autotransplanted, and homotransplanted kidneys. Preliminary studies suggested that transplants possessed much lower enzyme levels than normal kidneys. Factors such as the interstitial oedema and

*Honorary Research Assistant, Wellcome Trust Travelling Fellow.

cellular infiltration, which may give rise to apparent changes in kidney enzyme levels, have been assessed by extending the preliminary studies to include enzyme histochemistry and microchemical analyses. Enzymes studied histochemically include succinic- and malic dehydrogenases, D.P.N.H. (reduced diphosphopyridine nucleotide) diaphorase, cytochrome oxidase, acid phosphatase, and β -glucosaminidase.

Chemical analyses included D.N.A. (desoxyribose nucleic acid), R.N.A. (ribonucleic acid), protein, and methionine. Hydroxyproline and hexosamine (representing collagen and mucopolysaccharide respectively), and the wet weight to dry weight ratio of the tissue were also estimated.

Materials and Methods

Animals.—All dogs used in these experiments were greyhound bitches with an average weight of about 22 kg.

Surgical Techniques.—Kidneys were transplanted by techniques previously described (Dempster, 1953a, 1954).

Removal of Kidneys.—Functioning autotransplants were removed between the second and seventh days after transplantation. Homotransplants have been divided into two main groups—those which were removed between the second and fourth days while still functioning, and those which proceeded to oliguria. Oliguric kidneys have been further divided into two subgroups according to the day that they became oliguric. The first subgroup includes those kidneys becoming oliguric on the third, fourth, fifth, or sixth day; the second subgroup includes kidneys becoming oliguric later than the sixth day. The longest survival was 17 days.

Assessment of Oliguria.—Kidneys were adjudged to be truly oliguric when a sudden decrease in daily urine output occurred together with a poor response to successive infusions of 500 ml. of 0.9% saline and 300 ml. of 2% saline. This condition was often accompanied by a high blood-urea level and a toxic syndrome (Dempster, 1953b).

Enzymology.— β -glucuronidase, β -galactosidase, and β -glucosaminidase were estimated by hydrolysis of phenolphthalein glucuronide, *o*-nitrophenyl- β -D-galactoside and *p*-nitrophenyl- β -D-glucosaminide respectively (Findlay *et al.*, 1958; Levvy and Marsh, 1959; Conchie *et al.*, 1959). Malic and succinic dehydrogenases were estimated manometrically (Umbreit *et al.*, 1957). All enzymes were estimated on complete homogenates of kidney cortex, the media for homogenization being water for the hydrolases and 0.25M sucrose for the dehydrogenases.

Histochemistry.—Small blocks of kidney were frozen in liquid oxygen and sections cut at 6 microns in a Bright cryostat at -15° C. Succinic dehydrogenase, malic dehydrogenase, and D.P.N.H. diaphorase were demonstrated by the M.T.T. method of Pearse (1960), cytochrome oxidase by the method of Burstone (1960), and acid phosphatase by the method of Burstone (1958). β -Glucosaminidase was demonstrated using naphthol AS-LC-N-acetyl- β -glucosaminide as substrate (kindly supplied by Dr. D. Janigan).

Histology.—Material for paraffin sections was taken immediately on removal of the kidney and fixed in 10% formol-saline. Sections were stained with haematoxylin and eosin or methyl-green-pyronine.

C. T. DOLLERY *ET AL.*: STUDIES OF THE RETINAL CIRCULATION WITH FLUORESCEIN

FIG. 1

FIG. 2

FIG. 3

FIG. 4

FIG. 5

FIG. 6

FIG. 7

FIG. 8

FIG. 9

FIG. 10

FIG. 11

FIG. 12