- Everson, T. C. (1964). *Ann. N.Y. Acad. Sci.*, **114**, 721.
 — and Cole, W. H. (1959). *J. Amer. med. Ass.*, **169**, 1758.
 Fishman, W. H., and Farnelant, M. H. (1953). *Endocrinology*, **52**, 536.
 Franks, L. M. (1954). *J. Path. Bact.*, **68**, 603.
 — (1956). *Lancet*, **2**, 1037.
 Gardner, W. U. (1946). *Cancer Res.*, **6**, 493.
 Ghaleb, H. A. (1961). Ph.D. Thesis, University of London.
 Heinbecker, P., Rolf, D., and White, H. L. (1943). *Amer. J. Physiol.*, **139**, 543.
 Hewlett, J. S., Hoffman, G. C., Senhauser, D. A., and Battle, J. D. (1960). *New Engl. J. Med.*, **262**, 1058.
 Hooker, C. W., and Pfeiffer, C. A. (1942). *Cancer Res.*, **2**, 759.
 Horning, E. S. (1952). *A.R. Brit. Emp. Cancer Campgn*, **30**, 60.
 — (1954). *Brit. J. Cancer*, **8**, 627.
 — (1956a). *Ibid.*, **10**, 678.
 — (1956b). *Z. Krebsforsch.*, **61**, 1.
 — and Whittick, J. W. (1954). *Brit. J. Cancer*, **8**, 451.
 Jacobson, L. O., Goldwasser, E., Fried, W., and Plzak, L. (1957). *Nature (Lond.)*, **179**, 633.
 Kirkman, H. (1959). *J. nat. Cancer Inst., Monogr.* No. 1.
 — and Bacon, R. L. (1949). *Anat. Rec.*, **103**, 475.
 — (1952a). *J. nat. Cancer Inst.*, **13**, 745.
 — (1952b). *Ibid.*, **13**, 757.
 Kochakian, C. D., and Fox, R. P. (1944). *J. biol. Chem.*, **153**, 669.
 Korenchevsky, V., and Hall, K. (1944). *J. Path. Bact.*, **56**, 543.
 — and Ross, M. A. (1940). *Brit. med. J.*, **1**, 645.
 Lattimer, J. K. (1942). *J. Urol.*, **48**, 778.
 Lavender, A. R., Pullman, T. N., Rasmussen, H., and Aho, I. (1961). In *The Parathyroids*, edited by R. O. Greep and R. V. Talmage, p. 406. Thomas, Springfield.
 Lipschütz, A., and Vargas, L. (1941). *Endocrinology*, **28**, 669.
 Longley, L. P. (1942). *J. Pharmacol. exp. Ther.*, **74**, 61.
 Ludden, J. B., Krueger, E., and Wright, I. S. (1941). *Endocrinology*, **28**, 619.
 MacKay, E. M. (1940). *Proc. Soc. exp. Biol. (N.Y.)*, **45**, 216.
 McQueen-Williams, M., and Thompson, K. W. (1940). *Yale J. Biol. Med.*, **12**, 531.
 Matthews, V. S., Kirkman, H., and Bacon, R. L. (1947). *Proc. Soc. exp. Biol. (N.Y.)*, **66**, 195.
 Miller, H. C., Woodruff, M. W., and Gambacorta, J. P. (1962). *Ann. Surg.*, **156**, 852.
 Naets, J. P. (1958). *Nature (Lond.)*, **182**, 1516.
 Page, I. H., and Bumpus, M. (1960). In *Clinical Endocrinology*, edited by E. B. Astwood, **1**, 591. Grune and Stratton, New York.
 Pasqualini, R. Q., and Imbriano, A. (1947). *J. Amer. med. Ass.*, **135**, 374.
 Prentiss, R. J., Hollander, F. G., Mullenix, R. B., Feeney, M. J., and Howe, G. E. (1962). *Calif. Med.*, **97**, 235.
 Richardson, F. L. (1957). *J. nat. Cancer Inst.*, **18**, 813.
 Riches, Sir E. (1963). *Ann. roy. Coll. Surg. Engl.*, **32**, 201.
 Sakula, A. (1963). *Brit. J. Dis. Chest*, **57**, 147.
 Selye, H. (1939). *J. Urol.*, **42**, 637.
 — (1940). *J. Pharmacol. exp. Ther.*, **68**, 454.
 — (1941). *J. Urol.*, **46**, 110.
 — and Rowley, E. M. (1944). *Ibid.*, **51**, 439.
 — Stone, H., Nielsen, K., and Leblond, C. P. (1945). *Canad. med. Ass. J.*, **52**, 571.
 Smithers, D. W. (1962). *Clin. Radiol.*, **13**, 132.
 Stanbury, S. W. (1963). In *Hormones and the Kidney*, edited by P. C. Williams, p. 165. Academic Press, London.
 Welsh, C. A., Rosenthal, A., Duncan, M. T., and Taylor, H. C. (1942). *Amer. J. Physiol.*, **137**, 338.
 West, C. D., Damast, B. L., Sarro, S. D., and Pearson, O. H. (1956). *J. biol. Chem.*, **218**, 409.
 Whisenand, J. M., Kostas, D., and Somers, S. C. (1962). *West. J. Surg.*, **70**, 284.
 White, H. L., Heinbecker, P., and Rolf, D. (1949). *Amer. J. Physiol.*, **156**, 67.
 Winternitz, M. C., and Waters, L. L. (1940). *Yale J. Biol. Med.*, **12**, 705.
 Zak, F. G. (1957). *J. Mt Sinai Hosp.*, **24**, 1352.

Relief of Post-operative Pain: Comparison of a 25% Nitrous-oxide and Oxygen Mixture with Morphine

G. D. PARBROOK,* M.B., F.F.A. R.C.S.; G. A. D. REES,† M.B., F.F.A. R.C.S.

G. S. ROBERTSON,‡ M.B.

Brit. med J., 1964, **2**, 480–482

Nitrous oxide has had a wide use as an analgesic in midwifery and in dental surgery, but its use does not appear to have been investigated for the relief of post-operative wound pain.

We describe here a study designed to compare the relative potency of nitrous oxide and morphine for relief of pain in patients during the first 48 hours after upper abdominal operations. Post-operative pain in such patients restricts breathing to a marked degree, and the relief by analgesics of this restriction can be assessed by simple tests of vital capacity in the manner described by Overholt (1930), Bromage (1955), and Masson (1962). Previous work on volunteers (Chapman *et al.*, 1943; Sonnenschein *et al.*, 1948; Delisle Burns *et al.*, 1960; Dundee and Moore, 1960; Parkhouse *et al.*, 1960; Robson *et al.*, 1960; Dundee *et al.*, 1962) suggested that 25% nitrous oxide in oxygen was the most suitable concentration for this study.

Material and Methods

The investigation was limited to patients under the age of 60 who had operation wounds in or extending to the upper abdomen. Prior to operation the patients were seen, the nature of the trial was explained, and their co-operation obtained; vital capacity and peak expiratory flow readings were then

measured and the nitrous oxide and oxygen breathing apparatus was demonstrated.

If vital capacity is measured before operation and again after operation, when the patient is fully recovered from the anaesthetic, its diminution gives some measure of the impairment of respiratory movement. This can be taken as some measure of the degree of pain. If this pain is then relieved by an analgesic the vital capacity increases in proportion to the efficacy of the drug. A perfect analgesic will restore the vital capacity to virtually the pre-operative value, whereas a drug having no effect will make no difference to the vital capacity. We therefore used changes in vital capacity, and, to a less extent, those in peak expiratory flow rate, as a way of assessing the analgesic effects of nitrous oxide and morphine.

Vital Capacity.—This was measured by the Wright respirometer used with a mouth-piece and Ruben valve, and with a nose-clip applied. After practice readings the mean of three readings was noted. Although the Wright respirometer is less accurate than bulkier apparatus for vital capacity measurement, it was thought to give satisfactory comparative readings since the same respirometer was used in every case.

Peak Expiratory Flow Rate.—The Wright peak flow meter was used and the mean of three readings taken after practice "puffs."

25% Nitrous Oxide in Oxygen Mixture.—The mixture was delivered to the patient from a specially calibrated Walton 5 machine using a reservoir bag and a non-rebreathing circuit with a Ruben valve. The "pressure" control was adjusted so that the reservoir bag did not collapse during inspiration while

* Senior Registrar, Department of Anaesthetics, the Royal Infirmary and University of Aberdeen.

† Registrar, Department of Anaesthetics, the Royal Infirmary and University of Aberdeen. At present Senior Registrar, Department of Anaesthetics, Morriston Hospital, Swansea.

‡ Registrar, Department of Anaesthetics, the Royal Infirmary and University of Aberdeen.

the patient breathed through a mouthpiece. A nose-clip was worn to prevent air dilution of the mixture. The gas mixture delivered was regularly checked with a Beckman paramagnetic oxygen meter, and found to lie between 24 and 26% nitrous oxide in oxygen.

Post-operative Procedure.—The vital capacity and peak expiratory flow measurements were repeated the day after operation, at a time when no opiate had been administered in the previous five hours. The 25% nitrous oxide mixture was then given for a period of 10 minutes, immediately after which the readings were repeated. About 15 minutes later morphine was administered intramuscularly (1 mg./stone (0.16 mg./kg.) body weight) and further recordings were made one hour later. The same procedure was repeated some 11 hours later.

Results

Vital Capacity Results.—The results may be presented most clearly in the form of the respiratory restoration factor (R.R.F.) of Bromage (1955). This represents the improvement of the readings as a percentage of the improvement which one would see if the vital capacity was to return to the pre-operative value, and may be expressed as follows:

$$\text{R.R.F.} = \frac{\text{Analgesic V.C.} - \text{Pain V.C.}}{\text{Pre-operative V.C.} - \text{Pain V.C.}} \times 100$$

Fig. 1 shows the close agreement of the initial and repeat readings when they are expressed in this form. Analysis of the total readings gives the following results:

Mean R.R.F. Percentages			Residual Standard Deviation	Standard Error of Mean Difference
N ₂ O	Morphine	Difference		
19.17	10.84	8.33	12.52	2.80

FIG. 1.—Showing close agreement of the initial and repeat readings.

The residual standard deviation, which measures the variation in repeat measurements on the same patient, is the relevant index of the scatter of observations for the comparison of the effects of nitrous oxide and morphine. The ratio of the mean difference to its standard error gives a *t* value of 2.98 ($P < 0.01$). The 25% nitrous oxide in oxygen was therefore significantly more potent than morphine. The advantage was also significant for initial and repeat tests separately, whether calculated from R.R.F. or absolute values. Two sources of bias existed. Firstly, the vital capacity readings improved at an average rate of 1% per hour in this trial and consequently the results after morphine are 1% higher than they should be. Secondly, the Wright respirometer gives a slightly high reading if a denser gas is used in place of air, and laboratory checks showed that the recordings with a 25% nitrous oxide mixture are about 1% too high. As these two sources of bias were small, equal, and opposite, no corrections were made to the results.

Duration of Analgesia After 25% Nitrous Oxide.—After the administration of this mixture to the patient the duration of analgesia was checked by repeat vital capacity readings. On 20 occasions the results were repeated 10 minutes after nitrous oxide and on 10 occasions each they were repeated 5 and 15 minutes after nitrous oxide. The results (Fig. 2) indicated that significant analgesia persisted for five minutes and that the effect was entirely lost after 15 minutes.

FIG. 2.—Duration of analgesia after 25% N₂O mixture.

Combined Use of 25% Nitrous Oxide and Morphine.—On 15 occasions the administration of nitrous oxide was repeated one hour after morphine (1 mg./stone (0.16 mg./kg.) body weight) had been given. It was found that the combined use of these analgesics was more potent than either one by itself as judged by the vital capacity results ($P < 0.05$) (Fig. 3). These

FIG. 3.—Showing greater potency of combined therapy.

figures contrast with results following the combined use of mild analgesics, where no increased benefit occurred (Wolff *et al.*, 1941).

Peak Flow Results.—Not all patients could produce a recordable peak flow after operation, and initial and repeat recordings were obtained in 15 patients. Higher average readings were obtained with nitrous oxide, but were significant on the second occasion only ($P < 0.05$), and the combined use of nitrous oxide and morphine in 15 patients (Fig. 4) gave peak flow R.R.F. recordings significantly higher than either treatment by itself ($P < 0.05$). The peak flow recordings showed greater variability than those for vital capacity and they appeared to be less satisfactory as a test for analgesic potency. Masson (1962) and Parkhouse and Holmes (1963) showed that little change can be detected in post-operative peak flow readings after conventional narcotics.

Side-effects.—In nearly all patients a tendency to feel sleepy and appear sleepy was noticed after nitrous oxide. No other side-effect was noted. The gas appeared to be readily accepted

by the patients, but in one patient the administration of the mixture had to be discontinued prematurely because the mouth-piece produced a tendency to retch.

FIG. 4.—Post-operative readings.

Discussion

This pilot study suggests that a 25% nitrous oxide in oxygen mixture will be of value in the relief of post-operative pain.

This mixture could be provided either from a special machine or, perhaps more conveniently, from premixed cylinders (Tunstall, 1961, 1963; Cole, 1964). Short-term combined therapy of nitrous oxide and oxygen mixture and narcotic may, by producing better relief from pain, facilitate coughing and vigorous physiotherapy.

A recent study (Nunn and Payne, 1962) has suggested that many patients are mildly hypoxic in the first 24 hours after operation, and it may be that continuous treatment with nitrous oxide and oxygen mixtures warrants further investigation to relieve both anoxia and pain. The main indication for analgesia after operation is during the initial 24 to 48 hours (Parkhouse *et al.*, 1961), and this period appears to be within the safety margin for reversible bone-marrow depression which occurs on more prolonged therapy with higher concentrations of nitrous oxide (Lassen *et al.*, 1956; Green and Eastwood, 1963). The problems involved in the clinical application of a 25% nitrous oxide and oxygen mixture in post-operative pain relief are the subject of a further study.

Summary

The use of vital capacity and peak expiratory flow rate readings to estimate the degree of pain after upper abdominal surgery is described. A mixture of 25% nitrous oxide and oxygen was administered to 20 patients and a significant degree of pain-relief was obtained lasting for a few minutes after administration was stopped. In terms of vital capacity and peak flow the improvement was better than that obtained by morphine (1 mg./stone (0.16 mg./kg.) body weight). The combined use of nitrous oxide and morphine gave better results than either agent alone. The value of a 25% nitrous oxide mixture as an aid to physiotherapy and for prolonged use after abdominal operations is discussed.

We are indebted to the British Oxygen Company Limited, who lent us the Walton 5 machine, and to Mr. P. F. Jones and Mr. J. Kyle, who kindly permitted us to carry out the investigations on patients under their care. Our thanks are also due to Dr. W. Brass, senior lecturer in the Department of Statistics, for his assistance, and to Dr. W. N. Rollason, Director of the Department of Anaesthetics, for his encouragement and guidance in this work.

REFERENCES

- Bromage, P. R. (1955). *Brit. med. J.*, **2**, 589.
 Chapman, W. P., Arrowood, J. G., and Beecher, H. K. (1943). *J. clin. Invest.*, **22**, 871.
 Cole, P. V. (1964). *Anaesthesia*, **19**, 3.
 Delisle Burns, B., Robson, J. G., and Welt, P. J. L. (1960). *Canad. Anaesth. Soc. J.*, **7**, 411.
 Dundee, J. W., and Moore, J. (1960). *Brit. J. Anaesth.*, **32**, 453.
 ——— Nicholl, R. M., and Black, G. W. (1962). *Ibid.*, **34**, 158.
 Green, C. D., and Eastwood, D. W. (1963). *Anesthesiology*, **24**, 341.
 Lassen, H. C. A., Henriksen, E., Neukirch, F., and Kristensen, H. S. (1956). *Lancet*, **i**, 527.
 Masson, A. H. (1962). *Anesth. Analg. Curr. Res.*, **41**, 615.
 Nunn, J. F., and Payne, J. P. (1962). *Lancet*, **2**, 631.
 Overholt, R. H. (1930). *J. Amer. med. Ass.*, **95**, 1484.
 Parkhouse, J., Henrie, J. R., Duncan, G. M., and Rome, H. P. (1960). *J. Pharmacol. exp. Ther.*, **128**, 44.
 ——— and Holmes, C. M. (1963). *Proc. roy. Soc. Med.*, **56**, 579.
 ——— Lambrechts, W., and Simpson, B. R. J. (1961). *Brit. J. Anaesth.*, **33**, 345.
 Robson, J. G., Delisle Burns, B., and Welt, P. J. L. (1960). *Canad. Anaesth. Soc. J.*, **7**, 399.
 Sonnenschein, R. R., Jamison, R., Loveseth, L. J., Cassels, W. H., and Ivy, A. C. (1948). *J. appl. Physiol.*, **1**, 254.
 Tunstall, M. E. (1961). *Lancet*, **2**, 964.
 ——— (1963). *Brit. med. J.*, **2**, 915.
 Wolff, H. G., Hardy, J. D., and Goodell, H. (1941). *J. clin. Invest.*, **20**, 63.

Effect of Circulating Antibody to Insulin on Serum Levels of Insulin-like Activity in Rats, Guinea-pigs, and a Diabetic Patient

NAGUIB SAMAN,* D.M., M.R.C.P.ED.; RUSSELL FRASER,† M.D., F.R.C.P.

Brit. med. J., 1964, **2**, 482–485

Our recent attempt (Samaan *et al.*, 1962) to make the fat-pad bioassay for insulin more specific by incorporating an antibody in the assay procedure revealed that most but not all of the insulin-like activity (I.L.A.) in peripheral serum was "atypical"—that is unaffected in its *in vitro* activity by an antibody to insulin. Subsequent studies have confirmed that while the pancreas probably secretes insulin into the bloodstream in "typical" form—that is, one whose bioassay effect

is antagonized by an antibody to insulin—the liver transforms some of this typical insulin into the other, or atypical, form which constitutes most of the I.L.A. in peripheral serum (Samaan *et al.*, 1962a, 1962b, 1963). It is surprising that the guinea-pigs in whose sera this antibody is raised show no hint of diabetes mellitus, for their glucose-tolerance tests are strictly normal (Wright, 1961, 1963; Mann and Smith, 1963) even in the presence of the very high titre of potent antibody to insulin known to circulate in these animals. At first sight this does not seem consistent with the reports of other experiments in which injecting serum from these animals into other species has been followed within hours by the development of a full-blown

* Research Fellow of British Medical Research Council at the Department of Medicine, Postgraduate Medical School of London.

† Professor of Endocrinology, Department of Medicine, Postgraduate Medical School of London.