Lupus erythematosus cells obtained from the patient.

T. LEHOCZKY *ET AL.*: SKELETAL MUSCLE GLYCOGENOSIS IN IDENTICAL TWINS

FIG. 1

FIG. 2

FIG. 3

FIG. 1.—Haematoxylin and eosin stain; cross-section, low-power view.

FIG. 2.—Same as Fig. 1, high-power view.

FIG. 3.—Hale's method and P.A.S. stain (combined with hyaluronidase); cross-section, high-power view. Arrows indicate the mucopolysaccharide.

two cases, it is apparent that abnormally high oestrogen excretion and gynaecomastia were coincident and probably related.

Subject 15, on the other hand, had a comparatively low oestrogen excretion during the presence of gynaecomastia, and this was increased to a value within the normal range after the condition had subsided. This case might be analogous to the gynaecomastias of puberty, where pituitary activity, uninhibited by the normal adult secretion of oestrogen, results in the release of hormones which stimulate mammary growth. Subsequent to recovery of oestrogen secretion to normal levels, the pituitary is once again controlled and the release of mammatrophic hormones is depressed.

Discussion

Of the 15 cases of gynaecomastia studied, three can be explained on the basis of abnormally low oestrogen production, two cases were probably due to pubertal increases in pituitary activity, and five were consequent upon increased oestrogen secretion. In the remaining five cases some other cause of the increased breast growth must be sought.

Breast growth in cases with excessive oestrogen excretion is not unexpected, nor is its regression following the reduction of oestrogen excretion to a normal level. It is perhaps surprising that only one-third of the cases investigated came into this category. In only one case was the excessive oestrogen shown to be due to a tumour, although this possibility is often first considered as an explanation for mammary hypertrophy in the adult male. Apparently (Subject 14) the endogenous oestrogen can vary over a sufficiently wide range that transient gynaecomastia may be observed. Such fluctuations in oestrogen

production might well be related to variations in emotional or physical stress.

In the absence of methods for the estimation of pituitary secretion it is difficult to extend knowledge of the circumstances in which gynaecomastia may occur. Non-endocrine aetiological factors are important, but the nature of these is at present obscure. It should be accepted, however, that adult male mammary growth may be due to a number of fundamentally different reasons. Oestrogen excess is the least common of these, but, on account of its possible association with neoplasia, the one most desirable to exclude.

Summary

The urinary excretion of oestriol, oestrone, oestradiol, pregnanediol, pregnanetriol, 17-KS, and 17-KGS is reported for 11 normal men and 15 males with gynaecomastia.

Abnormally high oestrogen excretion was found in five of the patients with gynaecomastia. Three patients were excreting less oestrogen than is normal. In the remaining seven patients oestrogen excretion was within the normal range.

In two patients regression of mammary hypertrophy was associated with reduction of an unusually high oestrogen excretion, while in one other patient regression was accompanied by increase in oestrogen excretion to normal from a subnormal level.

REFERENCES

- Daly, D. W., Dossett, J. A., and Jull, J. W. (1963). *Brit. J. Surg.*, **50**, 816.
Jull, J. W., and Dossett, J. A. (1964). *Brit. med. J.*, **2**, 795.

Anterior Radiculopathy and Lupus Erythematosus Cells : Report of Case*

GUY GARGOUR,† M.D. ; KEITH MacGAFFEY,‡ M.D. ; SIMEON LOCKE,§ M.D. ; MAX D. STEIN,|| M.D.

[WITH SPECIAL PLATE]

Brit. med. J., 1964, **2**, 799-801

Collagen diseases as a group and systemic lupus erythematosus in particular are receiving increasing attention as their protean manifestations become more familiar. Affection of the nervous system in lupus erythematosus has been reported. In particular, Heptinstall and Sowry (1952), Siekert and Clark (1955), and Scheinberg (1956) have presented cases of involvement of peripheral nerves early in the course of the disease. The current case is unique in that findings of an anterior radiculopathy in the presence of positive lupus erythematosus preparations and positive serum fluorescent antibody reactions occurred in the absence of any of the known constitutional manifestations of systemic lupus.

Case Report

A 32-year-old white married woman was admitted to hospital in April 1963 with a chief complaint of progressive weakness and aching of her limbs. She had been in good health until April 1959, when, following the delivery of her third child, leg cramps, initially attributed to pregnancy, persisted for a period of six months. In

1959 she was involved in a minor car accident and sustained a "whiplash" injury. From that time she experienced occipital, suboccipital, and neck pain. Accompanying these she noted progressive weakness of her legs and "heaviness" of her body. One year prior to admission her grasp became weak. She had episodic blurring of vision associated with mild discomfort in the right eye. Ophthalmological examination revealed no abnormality. Persistent fatigability, lack of energy, and increased desire for warmth appeared.

In July 1962, because of persistence of headache and weakness of the legs, she was admitted to another hospital, where bilateral carotid and bilateral vertebral angiograms were performed because of papilloedema. These studies revealed no abnormality. Lumbar puncture disclosed fluid under "increased pressure" with a protein

* From the Department of Medicine, New England Deaconess Hospital, Boston, Mass., U.S.A.

† Medical Resident, New England Deaconess Hospital, Boston, Mass., U.S.A.

‡ Chief Resident in Medicine, New England Deaconess Hospital, Boston, Mass., U.S.A.

§ Neurologist, New England Deaconess Hospital, Boston, Mass., U.S.A.

|| Physician, Quincy City Hospital, Quincy, Mass, U.S.A.

of 640 mg./100 ml. Cervical myelogram was unremarkable. Repeat lumbar puncture disclosed a spinal fluid protein of 208 mg. After discharge from hospital, increasing paraesthesias of the hands were noted, with the appearance of a coarse tremor in the fingers and a tendency to "drop things." Persistent fatigue, aching of legs and shoulders, and menstrual irregularity occurred. In February 1963 repeat lumbar puncture revealed a spinal fluid protein of 1,080 mg. There was no antecedent history of allergic disorder.

Physical examination in April 1963 disclosed a well-developed, slightly obese, alert, co-operative, intelligent woman with a blood-pressure of 100/70 mm. Hg, a pulse of 80, and temperature of 99° F. (37.2° C.). The heart and lungs were unremarkable, the thyroid was normal, and no abdominal organs were palpable. Neurological examination revealed an absolutely normal mental status. The visual fields were full and the optic disks were of high colour, with blurred margins, but were not elevated. Confrontation fields revealed slight enlargement of the blind spot bilaterally. Mild proptosis was present on the right. The extraocular movements were normal. There was no nystagmus. The pupils were 4 mm. each and reacted to direct and consensual illumination. Facial strength and facial sensation were intact and the lower cranial nerves were unremarkable. Gait was wide-based, insecure, with slight foot-drop bilaterally. There was marked proximal weakness of the legs. The patient was unable to rise from a chair without using her hands, and could not go up or down stairs without support. A coarse, 4 to 5 per second, irregular, arrhythmic, asymmetrical tremor of the outstretched hands was present bilaterally. There were no cerebellar signs. All deep tendon reflexes were absent. The plantar response was bilaterally flexor. Sensation was intact to touch, pin, position, and vibration, although she complained of easily induced paraesthesias if she maintained one posture for short periods of time.

Laboratory Examination

The haemoglobin was 12.7 g., with a haematocrit of 37% ; white-cell count 6,100/c.mm., with a normal differential count ; total eosinophil count 125/c.mm. Sedimentation rate was 38 mm./hour. Reticulocyte count was 2.2%, and platelet count was normal. Serum iron was 65 mg./100 ml., with an iron-binding capacity of 293 mg./100 ml. Bleeding, clotting, and prothrombin times were normal. Blood Hinton reaction was negative. Bone-marrow biopsy revealed normal morphology of cellular elements, with no evidence of tumour or tubercles. Bone-marrow culture for fungus and acid-fast bacillus was unremarkable. Urine concentrated to 1027, and revealed no abnormality except for 1 to 6 white blood cells per high-power field. Urine electrophoresis was negative. Blood-urea nitrogen was 17 mg./100 ml., and 24-hour creatinine clearance 67 ml./minute, with a 24-hour volume of 670 ml. X-ray examination of chest, skull, and lumbo-sacral spine revealed no abnormality. Electrocardiogram and electroencephalogram were within normal limits. One hour postprandial blood sugar was 72 mg./100 ml., and a glucose-tolerance test was normal. Protein-bound iodine was elevated to 18.8 µg./100 ml. as a result of the antecedent myelogram. R.B.C. ¹²⁵I uptake was 15.6 µg./100 ml. Serum chemistry determinations revealed a calcium of 9.4 mg./100 ml. and serum protein of 7.5 g./100 ml., with an A/G ratio of 1.2. Serum electrophoresis revealed a moderate elevation of the gamma-globulin. Watson-Schwartz test was negative.

Lumbar puncture revealed an opening pressure of 210 mm. Cuff dynamics were free. Fluid was moderately xanthochromic and contained 2 red cells, 4 lymphocytes, a protein of 1,000 mg./100 ml., a glucose of 69 mg./100 ml., and a gamma-globulin of 4.5% of total protein as determined by the zinc sulphate method. Closing pressure after removal of 10 ml. was 160 mm. Cultures for tubercle bacillus and fungi were unremarkable. No tumour cells were seen. Electrophoretic pattern of spinal fluid revealed an albumin content of 66% of total protein. Alpha-1 globulin was 2%, alpha-2 globulin 4%, beta-globulin 10%, and gamma-globulin 17% of the spinal fluid protein. The gamma-globulin was double peaked, suggesting an unusual globulin. Serum latex fixation test was negative. Serum fluorescent antibody preparation was positive on one occasion and negative on a second. Lupus erythematosus preparations were found positive on three occasions (see Special Plate). Urinary 17-ketosteroid, 17-hydroxysteroid, and follicle-stimulating-hormone levels were within normal limits. Biopsy of the left biceps and the left rectus medialis revealed normal muscle with no evidence of vasculitis or denervation atrophy. A small branch of the anterior femoral

cutaneous nerve obtained at biopsy revealed no abnormality when stained by standard techniques.

Treatment

Treatment was started with 80 mg. of prednisone daily in divided dose. By the sixth day, in addition to a state of well-being, the patient noted subjective increase of strength. Objective evidence of mild improvement was present by the eighth day. After 10 days at high dosage the amount of prednisone was reduced over a 14-day period to 10 mg. in divided dose. Improvement of muscle strength continued at that level. When medication was reduced to 5 mg. daily, the patient reported increasing weakness by the third day. When returned to a regimen of 10 mg. daily, strength improved once again. Six months after beginning of treatment improvement continued. Lumbar puncture, performed in October 1963, six months after beginning treatment, revealed a spinal-fluid protein of 630 mg./100 ml. Neck and shoulder pain has returned, but in June 1964 strength was good enough to allow the patient to care for her home and family, in contrast to her state prior to treatment.

Comment

Involvement of the central nervous system in terminal cases of lupus erythematosus was originally reported in 1872 by Kaposi. Garcin's (1955) survey takes account of the researches by Keil (1933), Jarcho (1936), Klemperer *et al.* (1941), Cluxton and Krause (1943), and Daly (1945). Heptinstall and Sowry (1952) were the first to offer pathological verification of the peripheral neuropathy of systemic lupus. Although the clinical description of the neurological involvement in the case they report is suggestive of a proximally placed lesion, they attribute the disorder to the vascular involvement of peripheral nerves. Two years later Tumulty (1954) reported that cord lesions occurred but were uncommon. Ascending myelitis with bladder involvement was reported in Case 2 of Siekert and Clark's (1955) series, and Scheinberg (1956) described a case of polyneuritis in an 18-year-old girl as the most prominent manifestation of systemic lupus.

In the case reported here the clinical signs were of a diffuse radiculopathy affecting the motor system almost exclusively. The easily induced paraesthesias suggest that the sensory roots were not entirely spared, but clinical examination failed to give objective evidence of irreversible impairment of posterior-root function. That the motor involvement was not a peripheral polyneuropathy in the generally accepted sense but rather a radicular syndrome is suggested by its diffuse nature, the greater affection of proximal than distal musculature, and the excessive elevation of cerebrospinal fluid protein, which indicates that the process, whatever its nature, was in close proximity to the subarachnoid space. The weakness, areflexia, and high spinal fluid protein in the absence of a significant cellular reaction raised the diagnosis of infective polyneuritis. If one accepts the stringent criteria restated by Osler (1960) for the Guillain-Barré syndrome, it is clear this patient does not fit that category. Here, as in Guillain-Barré syndrome and other disorders in which spinal fluid protein reaches extremely high levels, papilloedema may occur. It has been suggested that this results from external hydrocephalus as a result of occlusion of the Paccianian granulations by protein coagulum.

In many features this case is reminiscent of the case reported by Austin (1958) in his review of chronic polyneuropathies. The possibility that one of the causes of this ill-defined group may be lupus erythematosus must be considered, although the finding of L.E. cells may not be specific (*New Engl. J. Med.*, 1963). The conjunction of a slowly evolving anterior radiculopathy in the presence of a high spinal-fluid protein, positive lupus erythematosus preparations, positive serum fluorescent antibody reaction, and a good response to steroid medication suggests that this peripheral neurological disorder may be an

unusual manifestation of subacute lupus erythematosus without systemic evidence of the disease. It is not possible, on the basis of the evidence at hand, to decide whether the radicular syndrome results from a vascular lesion or is a manifestation of a primary reaction of nerve root or its myelin.

Summary

The case of a 32-year-old woman with a chronic polyneuropathy manifest as an anterior radicular syndrome is reported. Weakness evolved during a three-and-a-half-year period and was associated with high spinal-fluid protein, positive lupus erythematosus cells, and positive serum fluorescent antibody reaction. There was no evidence of systemic lupus. The possi-

bility that lupus presents as an indolent disorder of nerve roots is suggested.

REFERENCES

- Austin, J. H. (1958). *Brain*, **81**, 157.
 Cluxton, H. E., and Krause, L. A. M. (1943). *Ann. intern. Med.*, **19**, 843.
 Daly, D. (1945). *J. nerv. ment. Dis.*, **102**, 461.
 Garcin, R. (1955). *Rev. neurol.*, **92**, 95.
 Heptinstall, R. H., and Sowry, G. S. C. (1952). *Brit. med. J.*, **1**, 525.
 Jarcho, S. (1936). *Johns Hopk. Hosp. Bull.*, **59**, 262.
 Kaposi, M. (1872). *Arch. Derm. Syph. (Wien)*, **4**, 36.
 Keil, H. (1933). *Arch. Derm. Syph. (Chic.)*, **28**, 765.
 Klemperer, P., Pollack, A. D., and Bachr, G. (1941). *Arch. Path.*, **32**, 569.
New Engl. J. Med., 1963, **268**, 1019.
 Osler, L. D. (1960). *Ibid.*, **262**, 964.
 Scheinberg, L. (1956). *Ibid.*, **255**, 416.
 Siekert, R. G., and Clark, E. C. (1955). *Neurology (Minneapolis)*, **5**, 84.
 Tumulty, P. A. (1954). *J. Amer. med. Ass.*, **156**, 947.

Medical Memoranda

"Stiletto Heel" Penetrating Fractures of the Skull

Brit. med. J., 1964, **2**, 801-802

Recently we have seen two cases of penetrating fracture of the skull resulting from blows with a shoe with a stiletto heel. The cases are of interest not only because of the nature of the weapon but because they show that serious damage may be present under relatively trivial-looking scalp lacerations. The second case demonstrates, in addition, the remarkable powers of recovery of the brain after serious injury.

CASE REPORTS

Case 1.—A man aged 31 years was admitted on 27 October 1963. In a fracas in a public house the previous night he had been struck on the head with a stiletto heel. He fell to the ground but rose immediately and began fighting back. In the morning he complained of headache and came to the hospital.

On examination he was fully conscious and alert and showed no abnormal neurological signs. Radiographs of the skull revealed a small stellate depressed fracture in the left parietal area and there was a small scalp laceration over this. He was operated on straight away by Mr. A. G. MacIntyre. The head was shaved and the laceration excised to reveal the fracture. It involved the brain. The fracture was excised and necrotic brain sucked out.

During the next day he had two epileptic fits, starting with convulsive movements of the right face and arm and culminating in unconsciousness and incontinence of urine. On recovering consciousness he was dysphasic and had an upper motor-neurone right facial paresis. He recovered completely on phenobarbitone and Epanutin (phenytoin sodium) and was discharged from hospital on 7 November 1963.

At the out-patients clinic on 11 February 1964 no neurological abnormality was detected. An electroencephalogram showed abnormal activity in the left temporo-parietal area.

Case 2.—A woman aged 36 years was involved in a street fight at 10.30 p.m. on 11 January 1964. She was hit on the head with the stiletto heel of a shoe by another woman. She went to a hospi-

tal where a small scalp wound was sutured and she was allowed home. She walked home, a distance of about a mile, and then began dragging her right leg. She became noisy and confused and lost her speech. On admission to the same hospital she was aphasic and had a right hemiplegia. A mitral systolic murmur was heard over the praecordium and cerebral embolus was considered as a possible diagnosis. Radiographs of the skull, however, revealed a depressed fracture.

She was transferred to the Neurosurgical Unit on 13 January, nearly 48 hours after the injury. On examination she was found to be alert but unable to talk and was distressed by this. By squeezes of her left hand, we were able to elicit satisfactory "yes" and "no" answers in reply to simple questions. (She is right-handed.) She had a complete right hemiplegia. Radiographs of the skull showed a small depressed fracture of comminuted type in the left parietal parasagittal area.

She was taken to the operating theatre immediately. The head was shaved. A small circular laceration, the size of a stiletto heel, with crushed infected edges, situated 2 in. (5 cm.) behind the coronal suture, very close to the midline on the left side, was found and excised. A circular hole in the outer table of the skull, about 8 mm. in diameter, was now seen. The fracture—fragments of which, together with some hair, had penetrated the brain—was excised and necrotic brain sucked out. Profuse bleeding from the sagittal sinus was controlled by application of gelatin sponge. A ventricular cannula was passed into the brain and 20 ml. of clot and necrotic brain was aspirated from the parietal lobe. Post-operatively *Staph. albus* and *Cl. welchii* were cultured from the aspirate, but fortunately no signs of infection developed.

Ten days after the operation she began to talk, at first in single words and later in complete sentences. The hemiplegia slowly recovered, improvement starting, in order, in the face, hand, and leg. She was discharged on 26 February. At that time she was talking normally, was able to read, and could walk with the aid of a calliper. Subsequently she made a full recovery.

COMMENT

It is well known that stiletto heels leave marks on all surfaces save very hard ones. It has been calculated that the force exerted on the area of surface in contact with the heel with a woman's weight on it is more than that exerted by an elephant's foot on the surface it treads on. These facts explain the lethal nature of such a shoe if it is used as a weapon on the head. The force of the arm is concentrated through the heel on a small area. A penetrating fracture may thus easily result. The