from which wisdom could derive ; and we profess to despise technology though we see it all around us.

Understanding of Man

The university whole-time clinical departments have helped the scientific understanding of disease, but it can hardly be said that they have helped the understanding of man. Patients are selected to illustrate the scientific aspects of medicine in which the teachers are interested. Nobody studies man himself. Even the psychiatrists often seem to be as afraid of human emotion as the rest of us. Future generations, paying tribute to the medical advances of our time, will say: "Strange that they never seemed to realize that the real causes of ill-health were to be found largely in the mind, and that even in 1965 there was hardly a teacher who could talk about sex except in biological terms" (which I must say takes most of the interest out of it).

A leading article in the *Observer* last Christmas said:

"Over the last 70 years the workings of the unconscious mind have been systematically studied (almost entirely outside academic institutions) and the knowledge gained offers for the first time an intelligible explanation of man's difficulties in following his own good intentions.

"It might have been expected in an age devoted to science and troubled by human destructiveness that the same sort of organized study that has been applied to the control of matter would have been applied to the control of human behaviour. . . .

"Can it be that this is the frontier where man's heart fails him ; is it possible that he finds it easier to look into the composition of atoms than into the basis of his own emotions ? Is it that his undignified inner feelings concerning sex and violence disturb him too deeply so that he prefers to study almost anything rather than this ?"

This I believe is where the universities have failed, and will go on failing if they recruit their teachers solely from those who would reduce the whole of medical knowledge to a series of hard scientific data. In all this I must share the blame as a medical teacher. Like others, I have been too involved in my own tasks and have too late come to see some of the absurdities and shortcomings of medical education.

The danger of saying these things is that I may be overheard by the universities. The danger is not that they will disapprove of me, but that they may agree with me but will find a way out in characteristic academic fashion and turn even the study of human behaviour into something in which man as a statistic is recognized but from which man as a personality is excluded, and find new reasons for facing anything rather than the facts of life.

Medical Education and Medical Practice*

ALASTAIR HUNTER,† M.D., F.R.C.P.

Brit. med. J., 1965, 2, 552-557

Medical education has recently been so much debated that you may wonder that a Bradshaw Lecturer should wish to say anything about it. But in spite of everything that has been written and said there is much confusion and much dissatisfaction with the existing state of affairs. Teachers in a number of branches of medicine, especially perhaps in psychiatry and social medicine, frequently draw attention to the archaic character of the present system. Articles and correspondence in the medical press and even in the leading national newspapers express both professional and public concern at what appears to be a growing separation between education and the needs of practice. The teaching hospitals are frequently criticized and represented either as the last strongholds of the dying authority of Harley Street or as clinical museums where university professors practise and teach medical science in defiance of the principles of good doctoring, and where the investigation of rare pathological conditions takes precedence over the care of those who are really sick.

Curiously enough, most of the criticisms have been directed against undergraduate education, which even now seems to be regarded as the beginning and the end of a doctor's training, while graduate and postgraduate education attract little attention. The failings of the preregistration year have passed almost unnoticed, and the need for special training for all forms of practice, including general practice, has, until recently, been scarcely recognized. For the most part this criticism indicates a failure of the profession as a whole and sometimes of teachers, to agree upon the objectives of the various stages in a doctor's

training, and to recognize the changes which modern medicine is forcing upon undergraduate education. The traditional fallacy that doctors on qualification must be fully fashioned for practice leads many to complain of their lack of clinical experience as house officers and of their failure to understand the subtleties of patient management as trainees and assistants in general practice. It is said that they have been taught too much science and too little medicine. There are many and divergent demands for reform, proposals for new curricula, and for new outlooks in teaching. Over and above these doubts about the effectiveness of the course, it is possible to discern a more radical difference of opinion about the function of a doctor. Distinctions made between individual and social medicine seem to picture the doctor of the future as a benevolent father figure concerned more to regulate the environment and to shepherd his flock through the hazards of life than to diagnose and treat their diseases.

Conflict Between Ideals and Technical Demands

There must always be a conflict in university medical education between the ideal of a broad education in science and the technical demands of a medical career. It is first felt in the preclinical course where there are two general patterns. One is designed as the first stage of an undergraduate course in the faculty of medicine ; the other is an honours degree course in natural science which includes enough of the preclinical subjects to satisfy the requirements of a 2nd M.B. examination. The course in the medical faculty is generally shorter and occupies five or six terms. It is devoted largely to anatomy and physi-

* The Bradshaw Lecture delivered to the Royal College of Physicians of London on 8 February 1965.

† Dean, St. George's Hospital Medical School, London.

ology, and its factual content in those subjects is at least equal to that of the natural sciences course. Biochemistry is included, but often as a part of physiology, and pharmacology is a rather uneasy third subject.

The natural sciences course as exemplified at Cambridge at least allows a wider choice of subject. Although limited by the predominance of anatomy and physiology in the first two years, it does give medical students opportunities for studying thoroughly subjects such as biochemistry and experimental psychology as well as pharmacology and pathology. At present the majority of students complete the course in three years and have the advantage of a rather leisurely approach to their subject, while a reasonable number complete it in two years and spend a third year studying a single subject in depth. Both courses have difficulties and disadvantages. The Cambridge course is designed primarily to teach biological science. It has been considerably affected by changes in the organization of the natural sciences tripos and is at the moment being redesigned as a separate medical sciences tripos. Some students find it too theoretical and complain of its remoteness from medicine, but generally it has a quality of excellence, especially for those studying Part II in a single subject, which compensates for any lack of relevance. Even so, there must be a danger that the proper scientific bias of the new tripos will become increasingly difficult to reconcile with the need to give medical students a broader background of knowledge.

The course in the faculty of medicine, at least in London, shows considerable signs of strain. Biochemistry is expanding separately from physiology and claiming an independent part in the 2nd M.B. examination. Pharmacology tends to be something of a poor relation, and psychology, genetics, and statistics are virtually omitted. The basic course has little room for manoeuvre and little chance to teach science except for the few who spend a further year studying anatomy or physiology for a B.Sc. degree. For many it has a rather breathless character culminating in the formidable obstacle of the 2nd M.B. Facts learned for the occasion are not always mastered, and many of them seem to have been rather thankfully forgotten in the short interval between the examination and the beginning of clinical work. The ultimate effect is hard to judge. Students do not always appreciate the potential importance of the course until they have had some experience of disease, and clinical teachers who hope to found their introductory teaching upon a sound knowledge of normal structure and function are apt to be disappointed. The fault lies surely in the rigid separation of preclinical and clinical teachers and teaching and in the attempt to expose medical students to a number of separate academic disciplines in which they are generally interested only in so far as they provide a stepping-stone to medicine.

Attempted Reforms

Attempts to correlate preclinical with clinical teaching have been made with varying degrees of success. Several new courses have been devised. But the essential requirement of any new curriculum is that it should find room for hitherto neglected subjects and remain sufficiently flexible to accommodate an inevitable increase in their number in the future. Another important question is whether to align the preclinical course with the science departments of the university or with the clinical departments of the hospital. In spite of its separation from clinical work a three-year degree course in biological science is attractive, but it must be obviously directed towards medicine if it is to hold the interest of the majority of medical students. Where the university and its medical school and hospital are close to one another, as is the case in provincial schools, teachers in all departments should be able to co-operate in overcoming specific difficulties. But circumstances will almost certainly favour a course in the faculty of medicine for the majority.

The role of the preclinical subjects in such a course is primarily to give students enough information about human function, structure, and behaviour to enable them to begin to learn about disease, and only secondly to teach them science. Close association between clinical and preclinical teachers both in research and in teaching is essential throughout the whole course if the present artificial divisions are to be abolished. In London some separation from the other faculties of the university is inevitable, and I appreciate the difficulties for preclinical departments. Physiologists, biochemists, and psychologists have affinities which should enable them to find satisfactory academic interests in clinical science. The interests of anatomists seem to be more in the field of human biology, but they might find a profitable link with departments of experimental pathology and surgery. Clinicians would certainly benefit by the association.

Reconsideration of Traditional Notions

Such a view clearly challenges two assumptions which have hitherto dictated the pattern of undergraduate medical education. The first is that the preclinical period is the only part of the course when medical students gain any insight into scientific method, and the second that a thorough knowledge of anatomy and physiology is a necessary and almost sufficient preparation for clinical work. Both are open to question. Some knowledge of anatomy and physiology is of course necessary, but it could well be in outline, and need not be exhaustive. Biochemistry has become equally important for medical students, while normal psychology, genetics, statistics, and possibly some elements of sociology cannot be neglected in future. Some rearrangement of the course is inevitable, and, unless it is to be further lengthened, and thereby made less suitable for medical students, some of its content will have to be transferred to B.Sc. courses or allowed to overlap with clinical teaching. Furthermore, clinical science at least uses scientific standards of measurement and scientific methods of inquiry and assessment of evidence. Already by the investigation of disease it has made valuable contributions to human physiology which have established it as a reputable university discipline, and, in fact, as the basic discipline of a medical course for the majority. Medical students certainly gain if they can spend their formative years with students of other faculties, but the advantages can be overemphasized. Medical education is in a sense liberal and humane, and medical students tend to become personally mature through contact, however indirect, with the work of the hospital. Even in multifaculty colleges they tend to have a life of their own which is robust and not altogether uncivilized, and their extracurricular activities are varied and stimulating. Generally they have small Latin and less Greek, but a culture based on Merseyside with undertones of Freud and Picasso is not a bad substitute and is a valid part of the new Humanism.

I hope that I have not given the impression of undervaluing the preclinical subjects or of being unsympathetic to the difficulties of those who teach them. My only concern is to ensure that what they have to contribute should be applied in the most effective way. The expansion of knowledge and the changing pattern of medicine itself must force us to reconsider long-accepted values. I am not sure that this is fully recognized in the universities. Some of the proposals for new medical schools seem to be ignoring entirely the advances in clinical science which have caused the most radical alteration in medical teaching. It is to be hoped that these schools will at least appoint clinical professors at the same time as their non-clinical staff before they perpetuate the existing separation irrevocably.

Preclinical Rigidity

The effectiveness of the preclinical course is hampered by the rigidity of its academic structure. The clinical course is

fragmented by dispersal among a variety of specialized professional departments. It lacks any real sense of direction, though it is vaguely committed to a policy of training for practice, particularly general practice. It is divided into a series of apprenticeships in the major specialties and their subdivisions, which have multiplied as professional organization has become more complicated. Even now, schools are periodically subjected to requests for additional instruction in subjects as varied as normal psychology, chronic rheumatic diseases, rehabilitation, family planning, and first aid for the injured. Many of them are of undoubted professional importance, but their relevance to undergraduate education is by no means clear. The programme, if there is one at all, is generally devised without reference to the teaching practices of individual schools. Often it is only too clear that, following the traditions of professional apprenticeship, teaching undergraduates is regarded as a necessary stage in the establishment of a professional specialty. The process is essentially one of ribbon development, and carries the inevitable risk that an unplanned drift to the periphery will obscure the central purpose. The real problem in medical education is whether clinical medicine should be taught as a technical discipline, or whether it should revert to a series of apprenticeships designed to cover the complete range of professional activity, and to provide a comprehensive experience. In fact, the vast increase of medical knowledge has determined the issue, and medical schools have already been forced to concentrate upon what Thomas Lewis called theoretical as opposed to vocational training, and to aim at a rational understanding of disease rather than a particular preparation for practice. The difficulties of achieving this objective are constantly underrated or tacitly ignored. The generally didactic character of pre-clinical teaching often leaves students singularly unprepared for clinical work. Many are still waiting to be taught and are not yet ready to learn by making accurate observations for themselves. Those who watch their progress closely commonly find that they take at least a year to acquire a sound clinical technique only to lose it again after the rather disconnected tour of the special departments which usually follows the initial period of ward work.

An Essential Quality

The unique function and skill of a doctor is the ability to identify and evaluate the signs and symptoms of mental and physical disorder in a patient. It is a technical skill which must be learned in the undergraduate period, otherwise it is unlikely to be learned at all. It is founded upon a substantial body of knowledge, but it can be properly understood only by studying patients. Diagnosis can no longer be taught as a matter of recognition, but as a precise assessment of clinical phenomena as evidence of disordered function. Problems must be studied in depth and much use made of laboratory methods of investigation which often demand an understanding of fairly complicated technical methods. Many have criticized this teaching as being too scientific, and have feared that the whole man will be lost in the analysis of the patient. Others have objected to what they regard as a specialist-based education with too much use of laboratory methods, and to undergraduate education being based upon observations drawn from rare and advanced cases which are believed to constitute the majority of those in teaching hospitals. Suggestions have been made that non-teaching district and regional hospitals would provide a better experience for students and future practitioners. The Annual Meeting of the British Medical Association in 1963 passed a number of resolutions in this sense, and recommended that teaching should once more concentrate on the art of medicine.

I do not share these views and believe that the best specialists are generally able to see their subject in relation to medicine as a whole. But with the rapidly changing technical processes of medicine the specialist sets standards to-day which will become

general to-morrow. The general practitioner, whom some prefer as a model for students, and for that matter the general physician, will exist as reputable professional figures only in so far as they can master their essentials. Medical students cannot be streamed in the early stages of their career, and must be taught by the highest available standards to ensure that every practising doctor shall be thoroughly trained in sound clinical method. Even in the interests of producing more doctors quickly it would be wrong to set two standards in training deliberately.

Theoretical Teaching

Thomas Lewis believed that undergraduate teaching would come to depend more upon whole-time teachers, while Illingworth has found the apprenticeship system unsuitable for teaching technical science. I believe that in a modified form it is valuable, partly because students learn more from continued observation than from occasional demonstrations, and partly because access to plenty of patients contributes much to their personal development. But clinical teaching to-day must be time-consuming and will often require co-operation between teachers in more than one branch of medicine. It cannot be left to chance and must always be more than a by-product of professional activity. It is no reflection upon the standards of non-university hospitals or of those who work in them to say that unless they were reorganized very much on the lines of existing university hospitals they would be unable to give medical students the foundation of clinical science which they need to-day, and will need even more in the future. The argument that undergraduate teaching is based too largely upon rare and difficult cases and neglects the common diseases is also, I believe, a mistake. My own experience is that the differences between the kind of patient seen in teaching hospitals and in other general hospitals of comparable size has been greatly exaggerated, but in any case the argument is educationally fallacious. Patients with complicated diseases having gross physical signs are particularly suitable for undergraduates because these signs are of fundamental importance and are fairly easily identifiable. Convenience apart, the chief reason for teaching undergraduates in hospitals at all is that most of the patients admitted require a high standard of medical care and often fairly full investigation, and thus enable students to understand the methods upon which the diagnosis and treatment of even more common and trivial diseases and of disease in its early stages depend. Teaching hospitals ought to provide as wide an experience of disease as possible, but their aim should be primarily to emphasize clinical methods and not to be comprehensive. Provided that they are large enough to develop research interests and maintain a reasonable turnover of patients their clientele should not differ appreciably from that of any other large hospital. They should undoubtedly accept district responsibility.

Relevant Application of Disciplines

There is a widespread notion that learning to analyse phenomena and to evaluate them by accurate scientific methods leads students to neglect the personal side of medicine. I think that it springs in part from a particularly English distrust of science as something necessarily cold and inhuman. I hope that I should be among the last to underrate the importance of the personal side of medicine or of the more intangible values to which Platt has referred, but in medicine wisdom is founded on knowledge and personal care and understanding cannot make up for lack of technical competence. If they disguise it they are dangerous. The art of medicine, like any other form of art, must comprehend the relevant application of scientific or technical disciplines. There is no evidence whatsoever that technical mastery dehumanizes medical students provided that

it is acquired in a place where medicine is practised with humanity, and where teachers accept personal responsibility for the care of their patients. But the problem is far more complicated and involves a change in the attitude of doctors to their patients and of medicine to the community as a whole. The old reticence about disease has been dispelled by the pressures of a better-informed society. It is generally acknowledged that some explanation of illness is not only due to a patient but is a part of his treatment. But the matter is not for the doctor alone. Journalism, radio, television, and other methods of communication are bringing a new image of medicine to the public, but general education in elementary human biology remains grossly inadequate, and many who now take a close interest in doctors and disease are familiar with little more than medical jargon. How much can they really understand and what do they want to know? Students are not in a position to do more than learn the rudiments of communication, because it is essentially a part of individual therapy, but they can at least become aware of the problem.

Personal problems are usually a natural reaction of patients and their relatives to sickness and its emotional and economic consequences. Many of them cannot be dealt with by doctors alone and need the assistance of social workers. But their understanding and their management in so far as it is possible are generally a matter for common sense. To separate them off from the problems of organic illness as social or personal medicine is to complicate them in a way which is in practice generally unnecessary. Educationally it is mistaken and under-rates medical students who are for the most part normal young people for whom motives of personal service are stronger than scientific curiosity. Their main need is to obtain a rational understanding of disease, but they can and do develop personal understanding from their contacts with patients. During 25 years of trying to teach them and learning from them I have seldom been disappointed with their standards of personal care.

Psychiatry

Two well-established professional specialties, psychiatry and social and preventive medicine, have recently claimed to be included as undergraduate disciplines. Both have some popular and political appeal, and some of those who support them most strongly do so for emotional reasons and because they believe in a muddled sort of way that they will be a corrective to too much science. The new awareness of psychiatry is timely. For long it was relatively, if not wholly, neglected. There were a number of reasons, such as the isolation of mental hospitals and the dominating influence of neurology, and above all a sense of mystique, and what to many seemed fantasy attaching to some concepts of psychopathology and psychotherapy. Professional reactions were seldom lukewarm, and the scepticism of some was matched only by the rather over-enthusiastic commitment of others. It was small wonder that teaching did not universally prosper.

Fortunately a more rational outlook is emerging and clinical academic departments are now recognized as having an equal part to play in education alongside those of organic medicine. Their development is often hampered by the difficulty of providing adequate in-patient departments. None the less the introduction of in-patient clerking in my own school has so radically changed the attitude of students to psychiatry and improved their understanding and handling of psychiatric problems that I am convinced that it is a necessary part of undergraduate education. But the problem of what to teach remains. At least we can agree that sound methods of case-taking must be learned and that the signs of a depression or of a dementia are as important to a doctor as those of a raised venous pressure.

Ellis and, more recently, Lewis in his *Harveian Oration* have clearly analysed the place of psychology and psychiatry

in undergraduate teaching in a way which must commend itself to teachers in other branches of medicine. Lewis rejected the introduction of psychotherapy, but there are undoubtedly those in this country as well as in the U.S.A. who would wish to introduce it. Some even plead for a more calculated personal involvement between student and patient, in place of the attitude of detachment which is believed to underlie the ordinary doctor-patient relationship. I would hesitate to subscribe to such a view. Students are certainly aware of the difficulties of making personal contact with patients but generally approach them with sympathy and a readiness to listen. To involve them in a more complicated process would distract them from the already difficult task of observing and interpreting symptoms and signs. In the event, doctors determine their personal methods of practice only when they have responsibility for the care of their patients, and, as Ellis has observed, the relationship of a student to his patient must be very different from that of a doctor.

The importance of academic units in psychiatry is not confined to their role in teaching. Without a clearer knowledge of the mechanism of mental disorder psychiatric teaching is likely to remain to some extent descriptive and perhaps intuitive. For some time ill-digested notions of psychosomatic medicine have blighted clinical thinking and confused diagnosis. As a practising physician I have come to believe that the chances of overlooking mental disease in patients with physical symptoms are at least equalled by the generally far more serious risk of missing organic disease. Depression and anxiety in those suffering from serious disorders such as early heart failure and cancer are all too easily mistaken for the primary diagnosis, and I feel equally certain that an appreciable number of those receiving prolonged and unsuccessful psychotherapy are suffering from less dangerous but equally disabling disease. Our whole thinking and teaching about so-called functional and organic disease is in need of revision.

Social and Preventive Medicine

Social and preventive medicine has been enthusiastically acclaimed in some quarters as a new form of medical practice. It encompasses a number of activities ranging from public health and community welfare to the much-discussed discipline of epidemiology. Its exact position in undergraduate education is ill-defined and probably limited. There is no doubt that medical students ought to be broadly familiar with the part which medicine and individual doctors play and should play in the community. They must also understand the administrative machinery for controlling communicable disease and for removing the major hazards from the environment. But a detailed knowledge is clearly a matter for postgraduate study. Leaving aside the public health aspects of the subject, I believe that the main contribution of social medicine to undergraduate education must be in the fields of epidemiology and social pathology. The wider understanding of aetiology which they encourage complements the more limited study of disease which is the concern of organic pathology. The statistical ascertainment of the prevalence of disease and its relationship to individual and environmental factors contributes an important technical method which is of use in medicine generally. Epidemiology is valuable also as a bridge between medicine in hospitals and in the community, and offers almost unlimited opportunities for co-operation between general practitioners, specialists in public health, and social workers generally and those working in a number of university departments.

Recently a different concept of social medicine has been advocated. It seems to assign to the doctor of the future a more positive role as a sort of prophylactic counsellor seeking out disease and the roots of disease in individuals and in the community. A different professional organization would pay as much attention to the environment as to the patient himself, and periodic medical examination backed by exhaustive labora-

tory investigations would take the place of ordinary consultations. Many of the technical methods are still untried and must be used with critical discernment. If diseases such as tuberculosis can largely be abolished by screening tests, others such as coronary artery disease, where early diagnosis is difficult and where the aetiology is unknown and probably complex, are harder to prevent. Physical and psychological stress can undoubtedly cause disease, but they are so much part of the process of living that I doubt whether they should ever become subject to any but the slightest medical direction. Positive health is a nebulous concept too near to happiness and spiritual welfare for which doctors are only indirectly responsible.

My purpose is not so much to air my prejudices against particular forms of preventive practice but to show that prevention, like cure, is an objective of medicine which depends for its attainment upon a knowledge of disease and its causation and of the methods of recognizing it in individuals. The rest is a matter of administration, propaganda, and ultimately of social and political action.

General Practice

Much of the criticism of undergraduate education comes from those in general practice. Suggestions have been made that, being increasingly deprived of the care of patients with organic disease, general practice should turn more towards psychiatry and social psychiatry, and that medical students should be taught more about these subjects and less about medical science and organic disease. I don't believe that the majority of general practitioners, who can and do give first-class all-round care to their patients, would agree. If their function is really to be altered and their training in medicine is to be seriously weakened, it would be better to be frank about it and to call them something other than doctors. As Hunt has stated, the essence of general practice is comprehensive and continuing help covering the greater part of illness and much more besides. Surely its proper preparation must be a thorough grounding in the disciplines which I have been discussing. The claim that academic teaching departments for what is essentially a postgraduate professional organization should be established in undergraduate schools is surely a contradiction. General practice will not survive by teaching medical students but by the quality of care that it can give to patients and by its ability to provide doctors with a satisfying professional life.

The clinical responsibilities of general practitioners, whether single-handed or in partnership, have always been heavy. Much of their time is spent in dealing with relatively trivial problems, and, like slip fieldsmen in a timeless Test match, they can spend long periods almost in technical idleness before meeting an emergency which may tax their resources to the limits. Even if they can find time to read adequately and to attend postgraduate courses they are apt to be divorced from the first-hand knowledge of new developments, particularly in treatment, which are readily available to those working in hospital.

General practice is unlikely to survive in its present form or in continued isolation from the hospital services. It is not only a question of providing laboratory services and trained social help, but of breaking down professional isolation. Contacts with specialists in consultation, or appointments as clinical assistants in special clinics, are helpful but limited, and as yet no system of appointments in hospital has been found which does not conflict with the duties of practice. It is not always possible or desirable for teaching hospitals to sponsor individual practices, as has been done successfully in Edinburgh, though departments of epidemiology could provide centres where general practitioners can satisfy their academic interests by co-operating in suitably designed research projects with academic workers. But the main need of general practi-

tioners, as indeed of all other practitioners, is continuing education. They should come freely to hospitals and take an active part in clinical discussions. They should be welcome in wards and out-patient departments, and should make a practice of personal consultation with consultants or their deputies on patients when they are returned to their care. Students would gain far more and would understand the specific rule of the general practitioner far better by being present and by following suitable patients with them after discharge than they could from any haphazard system of attachment. Tentative efforts along these lines have so far met with limited success. There is little difficulty in principle, but only in practice, which in the end must resolve itself in terms of time and money. Both would be well spent.

Undergraduate Course

I have pleaded that the purpose of undergraduate education is to provide a sound understanding of the technical disciplines of medicine. It is pertinent to ask whether our present system is best ordered to achieve it. I have already mentioned the difficulty of integrating the preclinical subjects. The clinical course is still dominated by the ideal, stated in the General Medical Council's Charter, of safe and efficient practice in all branches of medicine and by a compartmental tradition of teaching and learning.

A certain amount of compartmentalism is inevitable, but there is no longer any need for it to be made more rigid by separate examinations in each of the subjects. These have an almost unlimited scope covering the work of the whole curriculum. Theoretical and vocational aspects have almost equal weight in an examination which is still regarded as much a test of professional competence as a qualification for a university degree. It is still possible, though no less so than it was, for a candidate, after a course extending over four and a half years, to fail in a single part in spite of a competent performance in the remainder.

Much of the trouble lies in the interpretation of the General Medical Council's Charter. If the Council cannot alter it, it can at least follow up the liberal spirit of its last recommendations and give a more positive direction to education by redefining the purpose of the various stages of training.

Many believe that the undergraduate medical course could be shortened without becoming ineffective. A six-year course is difficult to justify, and even four and a half or five years of the London course is long for any first degree. Some of the time spent in a rather desultory round of unrelated clinical appointments and in preparation for badly spaced examinations could almost certainly be saved. But there are difficulties. The acquisition of responsibility for practice under supervision marks a new phase in learning, but it is a critical period in a doctor's development and must not come before he is personally mature. Many students are saturated with "A" level teaching at school and would undoubtedly benefit by spending a year in some profitable way before beginning their course. The reintroduction of the premedical subjects to form a three-year integrated course in human biology would have some advantages, but I doubt whether it is necessary or advisable for the majority who have studied science at school. Shortening the course would require a fairly drastic reorganization of the curriculum, and much that is at present included would be transferred to postgraduate courses or to a degree course in preclinical and clinical subjects. If the course were to be shortened the educational purpose of the preregistration period would have to be re-emphasized. Close supervision is necessary and a much wider variety of experience than is generally obtained. Rotating appointments in a number of specialties, including psychiatry, should be added to a sound general appointment in medicine, and, later, opportunities should be provided for working overseas and in general practice.

Conclusion

The undergraduate period is the beginning of medical education; its task is to lay the foundation of further education, both graduate and postgraduate, and to prepare students for practice under supervision. The foundation is a rational understanding of mental and physical disease, and of the methods of identifying them in individuals with a view to treatment and prevention. New therapeutic opportunities will demand even higher technical standards in the future, and the problem will be to achieve them while ensuring that education remains as humane as I believe it does to-day. Without such a foundation doctors will be merely purveyors of welfare. With it they can learn to practise efficiently according to their temperament and their personal beliefs. Some will envelop their patients in a benevolent paternalism, others will pursue a philosopher's stone for positive health and universal prevention, but many will be content to practise medicine well, and perhaps, as

Parkinson has said, "will accompany their patients on the second mile, and to the end of the road."

BIBLIOGRAPHY

- Balint, M., *Lancet*, 1961, 2, 1051.
 Brain, Lord, *ibid.*, 1964, 2, 325.
 Ellis, J. R., *ibid.*, 1956, 1, 813, 867.
 ——— *Brit. med. J.*, 1963, 2, 585.
 Herridge, C. F., *Lancet*, 1960, 2, 949.
 Hill, K. R., *ibid.*, 1964, 2, 517.
 Hunt, J. H., *ibid.*, 1964, 2, 84.
 Hutton, P. W., Williams, P. O., Graves, J. C., and Graves, V., *ibid.*, 1964, 1, 38.
 Illingworth, C., *ibid.*, 1964, 1, 283.
 Lewis, A., *Brit. med. J.*, 1963, 2, 1549.
 Lewis, T., *Lancet*, 1944, 1, 619, 649.
 Marshall, H. E. S., *ibid.*, 1960, 2, 949.
 Parkinson, J., *ibid.*, 1951, 2, 457.
 Pickering, G., *Brit. med. J.*, 1964, 2, 1615.
 ——— *Lancet*, 1964, 2, 653.
 Platt, R., *Linacre Lecture—Universities Quarterly*, 1963, 17, 327.
 Tredgold, R. F., *Lancet*, 1962, 2, 1344.
The Field Work of the Family Doctor, 1963. H.M.S.O., London.

Assessment of Postgraduate Medical Education

A Report to the Nuffield Provincial Hospitals Trust

The following report was prepared by Mr. D. H. Patey, Dr. J. O. F. Davies, and Dr. John Ellis. It is based on discussions with a special committee of the Trust, whose members were: Sir George Pickering (Chairman), Professor Hedley Atkins, Dr. John Fry, and Dr. John Revans. Mr. Gordon McLachlan edited the material, and Miss G. B. Bradfield acted as Secretary.

Brit. med. J., 1965, 2, 557–564

I. The Backdrop

This survey is yet another stage in the developments arising from the Christ Church Conference on Postgraduate Medical Education convened by the Trust in December 1961.¹

The Trust's policy in making grants was based on the general agreement reached at this conference, at which there were responsible spokesmen from all the main bodies and organizations involved in the field, that something could and should be done straight away to improve the arrangements for postgraduate medical education generally. The broad outline of requirements, unanimously agreed as urgently necessary, was based on the premises that postgraduate medical education must include the district hospital or hospital group, and be organized in regional schemes which would allow for a real association with the regional university. The immediate primary objective was to promote an educational atmosphere in the *basic local hospital* unit; to this end a consultant should be appointed by a fully representative and powerful regional committee as the "clinical tutor" responsible for the organization of teaching arrangements and the general care of those under training. At the same time *all* consultants and general practitioners should recognize their responsibilities to continuing education in the area. Equally the hospital authorities should be sympathetically and liberally disposed towards helping in the training not only of all grades of resident hospital staff but also towards the continuing education of all doctors. Among other things, this would entail the authorities seeking to provide certain necessary physical facilities. Ideally, too, attempts should be made to

develop criteria in order to set standards to which all such units should aspire. These would relate to:

- (1) standards of supervision;
- (2) quantity and variety of clinical material;
- (3) standards of records;
- (4) post-mortem service;
- (5) radiological and pathological services;
- (6) laboratory facilities;
- (7) seminars, clinicopathological conferences, etc.

At the regional level it was urged there should be a strong *regional* committee, the convener of which should be the postgraduate dean appointed by the university concerned. He should have proper accommodation and assistance and be responsible for the interrelationship between the regional hospital authority and the university regarding the arrangements for teaching (including the continuing education of general practitioners) in the region. *Nationally* it was recommended "as a matter of urgency" that some official body should be established as soon as possible representing the University Grants Committee and universities, the General Medical Council, the Departments of Health, the Royal Colleges, and the College of General Practitioners.

It was, however, recognized that in the absence of official and formal arrangements any immediate action would probably have to depend on local initiative; and to a large extent the trustees' policy was to support the promise of such enterprise at this level.

Following meetings convened by the Chief Medical Officer of the Ministry of Health and attended by representatives of all the principal professional, educational, and administrative

¹ *British Medical Journal* and *Lancet* of 17 February 1962.