After more than two years there is one survivor out of four patients who underwent surgical resection after improving with preliminary treatment, whereas five out of eight having surgery without preliminary improvement survived two years and three are still alive. Again this difference could be pure chance. It could indicate that preliminary improvement on cytotoxic drugs was of no advantage or was harmful; but it was probably at least in part due to the fact that improvement with cytotoxic drugs occurred in patients with more rapidly growing and metastasizing tumours, which were therefore less amenable to curative resection.

The pulmonary-artery-perfusion group was planned because of a remarkable result that occurred previously in one out of 25 patients treated in this way. He was a 52-year-old man who at thoracotomy was found to have an advanced oat-cell growth invading the main vessels. Three weeks later he was given one dose of 17.5 mg. of nitrogen mustard by drip into the right main pulmonary artery. The next day he felt much better, and five days later the chest x-ray picture showed marked reduction in the shadowing due to the growth. Two weeks later the x-ray appearance had returned to normal and bronchoscopy confirmed the apparent complete resolution of the neoplasm. Four further perfusions were carried out during the next year and he remained well and at work for this period. Early recurrence was first suspected after 16 months, and deep x-ray therapy was given after 20 months. He survived for two and a half years.

One patient in the present series treated by pulmonary-artery perfusion with cyclophosphamide survived over 18 months, but this was a case of squamous-cell carcinoma and may well have been naturally slow-growing. There were no other responses worthy of remark.

The place of cytotoxic-drug therapy in the treatment of carcinoma of the bronchus remains controversial. The present series suggests that in about 50% of patients the tumour will recede to some extent during high-dosage treatment with cyclophosphamide.

Poulsen's (1962, 1963) findings of a higher one-year survival rate in patients with operable squamous-cell carcinomas treated with pre-operative and post-operative cyclophosphamide than in a controlled group having surgery alone encourages further research on these lines in the early cases.

With advanced disease the remissions are usually short, ranging from two to six months, but often they are worth while from the point of view of relief of distressing symptoms. It remains to be seen whether better results can be obtained with maintenance therapy or the combination of two or more different agents, and whether, when combined with x-ray therapy, summation or sensitizing effects can be obtained.

Summary

Ninety-seven patients with bronchial carcinoma were treated with nitrogen mustard or cyclophosphamide. Of those treated with cyclophosphamide 51% responded favourably as judged by the objective evidence, compared with 33% of those receiving nitrogen mustard, but the limits of tolerance of nitrogen-mustard dosage may not have been reached. "Follow-up" treatment of patients who had improved was carried out with x-ray therapy, further intravenous injections, or pulmonary-artery cytotoxic perfusion.

We are grateful to the surgeons, physicians, and general practitioners of the York/Harrogate Area whose patients were included in the series; and also to Dr. J. Nuttall and Dr. Joan Mallender, who supervised the x-ray therapy; to Dr. Mary Catterall and Dr. J. Stoker, who carried out the respiratory physiology studies at Leeds General Infirmary; to the late Dr. G. Henry for help throughout; to Dr. J. M. Simister for advice on the cytotoxic drugs; and to Mr. Michael Curwen, of St. Bartholomew's Hospital, for much assistance and statistical advice; and to Mrs. Armitage and Mrs. Guest for the secretarial work.

REFERENCES

- Bass, B. H. (1960). *Brit. med. J.*, **1**, 617.
 Braun-Woitke, I. (1963). *Wien. klin. Wschr.*, **75**, 52.
 Flavell, G. (1962). *Brit. med. J.*, **1**, 284.
 Grenville-Mathers, R., and Trenchard, H. J. (1964). *Lancet*, **2**, 1200.
 Hatch, H. B., Bradford, J. K., and Ochsner, A. (1956). *J. Amer. med. Ass.*, **160**, 1129.
 Kent, L., and Reh, E. P. (1950). *Dis. Chest*, **17**, 190.
 Levine, B., and Weisberger, A. S. (1955). *Ann. intern. Med.*, **42**, 1089.
 Lynch, J. P., Ware, P. F., and Gaensler, E. A. (1950). *Surgery*, **27**, 368.
 Poulsen, O. (1962). *J. int. Coll. Surg.*, **37**, 177.
 — (1963). *Acta chir. scand.*, **125**, 498.
 Spear, P. W., and Patno, M. E. (1962). *Cancer Chemother. Rep.*, **16**, 413.

Prolongation of Skin Homograft Survival by Thalidomide

K. HELLMANN,* D.M., D.PHIL.; D. I. DUKE†; D. F. TUCKER,‡ B.SC.

Brit. med. J., 1965, **2**, 687-689

On immunogenetical grounds the foetus may be regarded as a homograft upon the mother, which, for reasons that are only poorly understood, escapes the usual fate of a homograft and remains unrejected. It seems possible, however, that some, perhaps most, early spontaneous abortions might be a specialized example of homograft rejection. Since a large number of foetuses expelled in such abortions are abnormal—for example, Moir (1964)—a non-toxic immunosuppressive drug might, on this idea, permit a malformed foetus to go to term that would otherwise have aborted. Most immunosuppressive drugs are, however, acutely toxic and their per-

formance in preventing homograft rejection has not been an unqualified success.

Thalidomide is known to be relatively non-toxic (Somers, 1960). If it also had immunosuppressive properties this would at once provide another aspect to explanations of its teratogenic action, and, at the same time, make it of considerable interest to those who seek to modify homograft rejection. To test whether it had any immunosuppressive properties its effect on the survival of mouse-skin homografts was examined. Mouse-strain combinations with strong ($C_{57}BL-C_3H/He$) and with weaker ($A-C_3H/He$) differences at the H-2 histocompatibility locus were used.

In previous experiments (to be published) it was found possible to modify the antigenic character of skin grafts by *in vitro* treatment prior to transplantation; thus rejection was obtained of syngeneic skin grafts previously incubated with

* Head of Cancer Chemotherapy Unit, Imperial Cancer Research Fund, Lincoln's Inn Fields, London.

† Senior Technician, Cancer Chemotherapy Unit, Imperial Cancer Research Fund, Lincoln's Inn Fields, London.

‡ Research Assistant, Cancer Chemotherapy Unit, Imperial Cancer Research Fund, Lincoln's Inn Fields, London.

allogeneic skin. Because of these findings some grafts were treated with thalidomide *in vitro* in the present experiments and some donors were given thalidomide before the grafts were excised.

Methods and Materials

The technique of skin-grafting did not differ significantly from that described by Billingham and Medawar (1951) except in the method of inducing anaesthesia. The mice each received intraperitoneally 0.1 ml. per 10 g. body weight of a freshly prepared 2.5% solution of bromethol (Avertin) in distilled water. Full-thickness skin-grafts were taken from either C₅₇BL or A strain female donors and transplanted to C₃H/He female recipients suitably prepared.

Thalidomide as a suspension in 0.5% carboxymethyl cellulose (C.M.C. in normal saline) was made up freshly just before it was given intraperitoneally. Amounts of thalidomide varied in different experiments (see Table), but the volume of C.M.C. in which they were given remained constant at 0.2 ml. Where recipients received post-operative thalidomide the first dose was administered on the day after skin-grafting, since even quite small doses were found to be toxic if given within 24 hours of bromethol administration. Dosage was continued daily until graft rejection was complete. The end-point for complete rejection was taken as that stage at which, on visual inspection, no viable graft tissue remained. Control animals received C.M.C. alone.

In some experiments, in addition to the recipients, the donors were also treated with thalidomide for varying periods before grafting. In other experiments the skin to be grafted was incubated for three hours in thalidomide (6 mg./100 ml.) in Krebs solution before transplantation. The Krebs solution had the following composition in g./l.: NaCl 6.9; KCl 0.35; KH₂PO₄ 0.16; CaCl₂·2H₂O 0.37; MgSO₄·7H₂O 0.29; glucose 1.0; and NaHCO₃ 2.1, aerated constantly with 95% oxygen and 5% carbon dioxide.

The animals were kept in groups of four or five and the bandages removed under anaesthesia 10 days after transplantation. The grafts were inspected daily until rejection was complete. The median survival times of the grafts are given together with their range. The median survival time is calculated as the time (in days) after transplantation at which 50% of the grafts have been completely rejected. The range gives the days on which the first and last grafts of a group were seen to have been completely rejected.

Results

The results of 15 experiments are given in the Table. The survival time of individual grafts in the control groups of both mouse-strain combinations was remarkably constant. In over 100 control transplantations from C₅₇ to C₃H (approximately eight animals for each of the 13 experiments) graft survival never exceeded 11 days, and in 18 control transplantations from A to C₃H (nine animals in each of two experiments) it was never more than 12 days. Complete rejection had thus occurred in both strain combinations within one to two days after removal of bandages.

When bandages were taken off, grafts on treated animals always appeared strikingly healthier and better healed than in those on controls in which often as many as half the animals in the group might already show signs of advanced graft rejection. In all treated animals, irrespective of the dosage scheme, rejection began later and advanced at a slower pace than in the controls. Rejection in controls was mostly complete within 12 to 24 hours; in the treated animals, on the other hand, the graft often broke up slowly into islands of viable-looking skin which took two to three days to disappear. Thalidomide seemed thus not only to slow down the onset of

homograft rejection, but also to make it less violent when it occurred.

Where recipients were treated the median survival time of grafts was improved if donor skin was also treated either *in vivo* or *in vitro* prior to transplantation. The best result (Experiment 10) was obtained where several days' treatment with thalidomide before transplantation of both donors and recipients was continued post-operatively in the recipients.

Treatment of donors alone (Experiments 6, 13, and 14) gave only a slight prolongation of graft survival, and treatment of recipients alone (Experiments 1, 2, 7, 8, and 9) similarly gave only a small prolongation, but donor and recipient treatment evidently summate in their effect.

No obvious toxic effects were encountered even with the fairly large doses of thalidomide given daily as in Experiment 10, and mortality rates in treated and control groups were not very different. It is possible that graft prolongations might have been obtained with smaller doses of thalidomide, since in those animals that came to post-mortem examination flecks of white solid, presumably thalidomide, were found in the peritoneal cavity 24 hours after the last injection. It is not at present possible to say what effect variation of dosage had on graft survivals.

Results of Experiments

Expt. No.	Treatment		No.	Survival Time in Days	
	Donors C ₅₇ BL	Recipients C ₃ H/He		Median	Range
0	Nil	Nil	103	10.9	10-11
1	Nil	4 hrs. post-op. 1 dose 25 mg./mouse TH	7	13.6	13-14
2	Nil	From 1 day post-op. for 14 days 5 mg./mouse/day TH	14	13.0	11-17
3	Grafts incubated for 6 hrs in 6 mg./100 ml. TH	Nil	16	13.9	12-16
4	Grafts incubated for 6 hrs in 6 mg./100 ml. TH	From 1 day post-op. for 14 days 5 mg./mouse/day TH	11	13.1	11-16
5	For 2 days pre-op. 200 mg./mouse/day TH	From 1 day post-op. for 16 days 5 mg./mouse/day TH	15	14.5	12-22
6	For 2 days pre-op. 200 mg./mouse/day TH	Nil	10	11.9	11-12
7	Nil	For 10 days pre-op. and from 1 day post-op. for 14 days 25 mg./mouse/day TH	10	11.8	11-12
8	Nil	For 10 days pre-op. 25 mg./mouse/day TH	10	11.5	11-12
9	Nil	From 1 day post-op. for 14 days 25 mg./mouse/day TH	5	12.2	11-17
10	For 10 days pre-op. 25 mg./mouse/day TH	For 10 days pre-op. and from 1 day post-op. for 16 days 25 mg./mouse/day TH	9	15.0	14-16
11	For 10 days pre-op. 25 mg./mouse/day TH	For 10 days pre-op. 25 mg./mouse/day TH	10	13.9	12-16
12	For 10 days pre-op. 25 mg./mouse/day TH	From 1 day post-op. for 16 days 25 mg./mouse/day TH	10	13.2	11-16
13	For 10 days pre-op. 25 mg./mouse/day TH	Nil	10	11.0	11-12
Donors A					
0	Nil	Nil	18	10.8	10-12
14	For 10 days pre-op. 25 mg./mouse/day TH	Nil	7	12.5	12-14
15	For 10 days pre-op. 25 mg./mouse/day TH	For 10 days pre-op. and from 1 day post-op. 25 mg./mouse/day TH	14	15.0	15-16

Discussion

With such a sharply defined temporal end-point to graft rejection in a large number of controls even small increases in length of survival assume significance. At doses that give no overt sign of toxicity, thalidomide is able to prolong mouse skin homografts across the H-2 histocompatibility barrier. This stands in marked contrast to immunosuppressive drugs such as cyclophosphamide, mercaptopurine, and methotrexate that are able to prolong graft survival in a variety of species, but only at severely toxic doses. The extent of the prolongation of skin-graft survival with thalidomide is entirely comparable

to that found with cyclophosphamide (Fox, 1964) and better than that found with Amethopterin (methotrexate) (Medawar, 1963) or mercaptopurine (Meeker *et al.*, 1960).

The mechanism by which thalidomide produces prolongation of skin-graft survival is not clear. It may be through a direct immunosuppressant action on the lymphoid system, though this is rendered less likely by the finding (see Table) that treatment of the donor before transplantation as well as the recipient increases length of graft survival. It seems more probable that, if a specific effect on the immune system is involved, thalidomide modifies the donor skin antigen or its recognition.

The findings of Roath *et al.* (1962, 1963) that thalidomide inhibited the phytohaemagglutinin-induced transformation of human lymphocytes into "blast" cells might also indicate that thalidomide could be expected to have an immunosuppressive action. Some doubts about these findings were raised by Lindahl-Kiessling and Böök (1963), though these authors gave insufficient details of their experiments to allow assessment of the validity of their criticism. In the light of the present findings this problem might usefully be looked at again.

These experiments of Roath *et al.* also led Playfair *et al.* (1963) to test oral thalidomide on skin homograft survival in mice; they, however, failed to observe any significant increase. It is difficult to know to what to attribute the disparity between their results and those reported here, since these authors too give virtually no experimental detail. It is unlikely to be due to the different route of administration which they employed, since in one experiment (not included in the Table because of excess mortality unrelated to the experiment) increased graft survival after oral thalidomide was also noticed in the two surviving mice of this experiment.

The timing of the thalidomide doses is of some importance. The best results in the present experiments were obtained when the recipients (as well as the donors) were pretreated. In this respect thalidomide is comparable to radiation, radiomimetic drugs, and cortisone-like steroids, which are most effective as immunosuppressants when given before an antigenic stimulus (Hitchings and Elion, 1963).

Timing was also found to be important by Tata (1964), who showed that thalidomide is toxic to the tadpole for a short period in metamorphosis. It therefore seems likely that thalidomide might act only at a particular time in cell division.

Because it is sparingly soluble in aqueous media and because of its instability the problem would be to have a sufficient concentration of the compound available to sensitive cells.

Although more information has been obtained with the $C_{57}BL-C_3H/He$ combination than with $A-C_3H/He$, it appears that the latter combination gives better prolongation than the former. This again might be expected if thalidomide were acting as an immunosuppressant.

Thus in a limited way the present experiments provide support for the idea that thalidomide might possess immunosuppressive properties.

Conclusion

Thalidomide prolongs skin homografts in mice at non-toxic doses. The best results were obtained when the donor graft was treated with thalidomide either *in vivo* or *in vitro* in addition to the recipient.

These findings are discussed and lend some support to the idea that thalidomide may have an immunosuppressive action.

We are much indebted to Professor L. Brent for showing us the transplantation technique he uses and to Sir Peter Medawar for a useful discussion. We would like to thank Mr. Ade Olukoya and Mr. Peter Hodgson for their technical help. We are particularly grateful to Dr. M. Perelman and Dr. S. S. Szinai, of Lilly Research Laboratories Ltd., for generous supplies of thalidomide, and also to Dr. G. F. Somers, of Reckitt & Sons Ltd.

REFERENCES

- Billingham, R. E., and Medawar, P. B. (1951). *J. exp. Biol.*, **28**, 385.
 Fox, M. (1964). *Transplantation*, **2**, 475.
 Hitchings, G. H., and Elion, G. B. (1963). *Pharmacol. Rev.*, **15**, 365.
 Lindahl-Kiessling, K., and Böök, J. A. (1963). *Lancet*, **2**, 405.
 Medawar, P. B. (1963). *Transplantation*, **1**, 21.
 Meeker, W. R., Condie, R. M., Good, R. A., and Varco, R. L. (1960). *Ann. N.Y. Acad. Sci.*, **87**, 203.
 Moir, J. C. (1964). *Munro Kerr's Operative Obstetrics*, 7th ed. Baillière, Tindall and Cox, London.
 Playfair, J. H. L., Leuchars, E., and Davies, A. J. S. (1963). *Lancet*, **1**, 1003.
 Roath, S., Elves, M. W., and Israëls, M. C. G. (1962). *Lancet*, **2**, 812.
 ——— (1963). *Ibid.*, **1**, 249.
 Somers, G. F. (1960). *Brit. J. Pharmacol.*, **15**, 111.
 Tata, J. R. (1964). *Nature (Lond.)*, **204**, 939.

L.E. Cells and Antinuclear Factors in Leprosy*

L. BONOMO,† M.D.; A. TURSI,‡ M.D.; G. TRIMIGLIOZZI,§ M.D.; F. DAMMACCO,‡ M.D.

Brit. med. J., 1965, **2**, 689–690

The suggestion has been made of the possible involvement of autoimmune mechanisms in some aspects of leprosy—namely, lepra reactions (Chini, 1961). A number of autoimmune-like reactions have been reported to occur in leprosy (Cathcart *et al.*, 1961; Rizzi *et al.*, 1962; Bonomo *et al.*, 1963). We were therefore particularly interested in investigating sera from leprosy patients for the presence of lupus erythematosus (L.E.) cells and antinuclear factors (A.N.F.) by means of the fluorescent-antibody technique.

Material and Methods

The sera of 55 unselected patients with leprosy (lepromatous type) in hospital at the Hansenian Colony, Gioia del Colle, Bari,

Italy, were tested for A.N.F. Blood specimens from 10 leprosy patients whose sera were found to contain A.N.F. were examined for L.E. cells as well.

Fluorescent Test for A.N.F.—The indirect technique described by Weir *et al.* (1961) was employed, with slight modifications (Bonomo *et al.*, 1965). Sera were tested in serial dilutions with buffered saline (pH 7.2), and those which showed fluorescent reactions in dilutions beginning from 1:2 up were

* This investigation was supported in part by Consiglio Nazionale delle Ricerche (Immunopatologia), Roma.

† Reader, Institute of Clinical Medicine, University of Bari, Bari, Italy.

‡ Fellow, Institute of Clinical Medicine, University of Bari, Bari, Italy.

§ Chief of Service, Hansenian Colony, Gioia del Colle, Bari, Italy.