- McArthur, J. W., Worcester, J., and Ingersoll, F. M. (1958). *Ibid.*, 18, 1186.
- (1961). In *Human Pituitary Gonadotropins*, edited by A. Albert, p. 210. Thomas, Springfield, Ill.
- Maqueo, M., Perez-Vega, E., Goldzieher, J. W., Martinez-Manautou, J., and Rudel, H. (1963). *Amer. J. Obstet. Gynec.*, 85, 427.
- Berra, C., Munguia, H., and Goldzieher, J. W. (1964). *Ibid.*, 90, 395.
- Martin, L., and Cuningham, K. (1961). In *Human Pituitary Gonadotropins*, edited by A. Albert, p. 226. Thomas, Springfield, Ill.
- Mastroianni, L., and Hongsonand, C. (1964). Paper presented at the Second International Conference on Intra-Uterine Contraception, 2 October 1964. Population Council, N.Y.
- Matsumoto, S., Ito, T., and Inoue, S. (1960). *Geburtsh. u. Frauenheilk.*, 20, 250.
- Mears, E. (1964). Paper presented at the Second International Congress of Endocrinology, London, 17–22 August 1964.
- (1965). "The Clinical Application of Oral Contraceptions," in *The Proceedings of the Symposium on Agents Affecting Fertility*, edited by C. R. Austin and J. S. Perry. Biological Council, Churchill, London.
- Østergaard, E. (1963). Paper presented at the Symposium on Ovarian Inhibition by Progestogens in the Woman, Brussels 2–3 March 1963. International Fertility Association.
- Overbeek, G. A., Madjerek, Z., and de Visser, J. (1962). *Acta endocr. (Kbh.)*, 41, 351.
- Parlow, A. F. (1961). In *Human Pituitary Gonadotropins*, edited by A. Albert, p. 300. Thomas, Springfield, Ill.
- Pincus, G. (1955). *Proceedings of the 5th International Conference on Planned Parenthood*, I.P.P.F., Tokyo, p. 175. I.P.P.F., London.
- (1956). *Acta endocr. (Kbh.)*, Suppl. No. 28, p. 18.
- (1959). *Vitam. and Horm.*, 17, 307.
- Rock, J., and Garcia, C. R. (1959). *Proceedings of the 6th International Conference on Planned Parenthood, New Delhi*, p. 216. I.P.P.F., London.
- Purshottam, N., Mason, M. M., and Pincus, G. (1961). *Fert. Steril.*, 12, 346.
- Rice-Wray, E., Aranda-Rosell, A., Maqueo, M., and Goldzieher, J. W. (1963). *Amer. J. Obstet. Gynec.*, 87, 429.
- Rock, J., Garcia, C. R., and Pincus, G. (1957). *Recent Progr. Hormone Res.*, 13, 323.
- Rosenberg, E., and Engel, I. (1960). *J. clin. Endocr.*, 20, 1576.
- Rudel, H. W., and Martinez-Manautou, J. (1965). Paper presented at the Meeting of the American Society for the Study of Sterility, San Francisco, 2–4 April 1965.
- Ryan, G. M., Craig, J., and Reid, D. E. (1964). *Amer. J. Obstet. Gynec.*, 90, 715.
- Schmidt-Elmendorff, H. (1965). Paper presented at the Meeting of the Mittelrheinischer Gesellschaft für Gynäkologie, Frankfurt.
- Staemmler, H.-J. (1964). *Die gestörte Regelung der Ovarialfunktion*, pp. 184–189. Springer, Berlin.
- and Staemmler, H. (1960). *Arch. Gynäk.*, 194, 183.
- Stevens, V. C., and Vorys, N. (1965a). To be published.
- (1965b). To be published.
- Sturgis, S. H. (1961). In *Control of Ovulation*, edited by C. A. Vilee, p. 213. Pergamon Press, New York.
- and Albright, F. (1940). *Endocrinology*, 26, 68.
- Taymor, M. L. (1964a). Paper presented at the Second International Congress of Endocrinology, London, 17–22 August 1964.
- (1964b). *J. clin. Endocr.*, 24, 803.
- Tyler, E. T., and Olson, H. J. (1958). *Ann. N.Y. Acad. Sci.*, 71, 704.
- Wide, L., and Gemzell, C. (1962). *Acta endocr. (Kbh.)*, 39, 539.
- Kaiser, J., and Gemzell, C. A. (1965). To be published.
- Zafartu, J. (1964). *Int. J. Fert.*, 9, 225.

Toxicity of Motor Neurone Disease Serum for Myelin in Tissue Culture

E. J. FIELD,* M.D., PH.D., M.S., M.R.C.P. ; D. HUGHES,† B.SC.

[WITH SPECIAL PLATE]

Brit. med. J., 1965, 2, 1399–1401

The serum of rabbits with experimental allergic encephalomyelitis is known to contain substances which produce demyelination in cultures of newborn rat cerebellum (Bornstein and Appel, 1961). Although the evidence is far from secure, experimental allergic encephalomyelitis is commonly regarded as a "model" for human multiple sclerosis, and serum from patients with this disease has been claimed to possess a similar demyelinating activity (Bornstein, 1963). The present work reports a high proportion of positive results from serum of cases of motor neurone disease.

Material and Methods

Cases of motor neurone disease which had been subjected to careful diagnostic scrutiny for another purpose were investigated (Vejjajiva, Foster, and Miller, 1965). Sera were divided into four or five aliquots for separate testing and stored at -65°C . for not longer than two months before use.

Explants of newborn rat cerebellum were set up on coverslips coated with rat-tail collagen and grown at 35°C . in roller tubes for 18 to 21 days. The nutrient medium comprised Simm's balanced salt solution 40% ; Parker 199 medium (Glaxo) 5% ; bovine serum ultrafiltrate (Microbiological Associates, Bethesda, Md) 25% ; human ascitic fluid 25% ; chick embryo extract (nine days *in ovo*) 5%. Glucose was added to a final concentration of about 600 mg./100 ml. and pH was maintained within the range 6.9–7.1 by adjustment with CO_2 as required. Cultures were screened in bright field or phase-contrast illumina-

tion, and those quite free from degenerative or fatty change were selected for testing purposes.

Tests were carried out in either a Maximow or an optical-glass viewing-chamber (Rosenthal, New York), maintained at 35°C . on a hot stage ; or they were carried out in chambers kept in the incubator and examined at intervals. In this way it was possible to obtain an overall view of the culture and revert to a specific field in order to record the sequence of changes photographically.

In testing each serum the following preparations were set up : (a) culture plus serum under test (25% in nutrient fluid) and (b) culture plus normal human serum (25% in nutrient fluid).

As an additional control, culture plus nutrient fluid alone was set up in many cases. Two preparations of (a) and (b) were set up for each test ; occasionally three. Altogether 101 cultures of (a), 63 of (b), and 40 with nutrient fluid alone were set up, so that more than 200 cultures were used in all. Myelotoxicity of serum is known to be complement-dependent (Bornstein, 1963), and in some early additional experiments fresh guinea-pig serum was added as a source of complement. In these cases additional controls were set up to establish the non-toxicity of the guinea-pig serum. Later, however, it was found unnecessary to introduce fresh guinea-pig serum, since direct estimation showed adequate complement levels to remain in the stored human sera. At the conclusion of a test the culture was fixed in formol-saline and the axis cylinders were impregnated by the method of Bodian.

Altogether serum was tested from 21 patients in whom the diagnosis of motor neurone disease was based upon the most stringent diagnostic criteria, five in whom the diagnosis was highly probable, and 20 normal subjects. A total of 204 cultures was examined.

* Honorary Director, Medical Research Council Demyelinating Diseases Research Unit ; Neuropathologist, Royal Victoria Infirmary and University of Newcastle.

† Technical Officer.

Results

In assessing the effects of any particular serum account was taken of the overall effects upon myelinated fibres and glial cells throughout the culture, since misleading conclusions might be drawn if attention was confined to a single selected field. Variation of effect within a given culture was often observed and might well have been related to the concentration of myelinated fibres able to fix toxic antibody. The length of time which elapsed before morphological changes occurred was noted and some assessment of toxicity estimated as the percentage number of fibres damaged at a given time. Results were then scored according to the schema in Table I. Inevitably some element of subjectivity was present in assessing minor degrees of change, so that it was decided to regard $\pm$ readings as negative. Results are set out in Table II.

TABLE I.—Myelin Destruction (% Fibres)

Time	5%	25%	50%	75%
6 hours ..	+	++	+++	++++
24 " ..	$\pm$	+	++	+++
48 " ..	—	$\pm$	+	++
72 " ..	—	$\pm$	+	+

TABLE II

Type of Sample	Score				Positive	Negative	Total
	4+	3+	2+	1+			
A. Motor neurone disease where diagnosis not in doubt (21 cases) ..	1	0	7	5	13	8	21
B. Motor neurone disease; some doubt as to diagnosis (5 cases) ..	0	1	1	1	3	2	5
C. Normal (20 cases) ..	0	1	1	0	2	18	20

The χ^2 test applied to this table (with Yates's correction for continuity) shows that $P=0.005-0.001$ —that is, the number of positive results with clear cases of motor neurone disease is different from that of normal sera.

No correlation could be observed between clinical evidence of upper motor neurone involvement (Babinski sign) and positivity of the test. Thus the most positive patient (4+) showed no Babinski sign, while some of the negative ones did do so. A typical sequence of events during the testing of a case of motor neurone disease is shown in Figs. 1, 2, and 3 (Special Plate). In Fig. 1 clearly outlined myelinated nerve fibres are seen coursing through the culture. In bright field illumination they appear as two parallel refringent lines separated by a dark line. At the beginning of an experiment glial cells (? oligodendrocytes) are pear-shaped or show two or more processes. With most sera (whether they turn out to be myelotoxic or not) degenerative changes in these glial cells begin within about an hour. They become swollen and rounded, and the clear cytoplasm contains small granules which are in constant Brownian motion. A certain number of these glial cells go on to dissolution, but in many non-toxic sera the majority may gradually revert to normal. In general the changes in glia in all sera are well marked (specially if 50% serum is used in a test) and not specific.

Fig. 2 shows the same field four and three-quarter hours after the application of 25% motor neurone disease serum. Swollen glial cells are prominent (B) and show the cytoplasmic particles referred to above. As they expand they compress myelinated fibres which run in their vicinity. Myelin sheaths in this figure are beginning to show localized expansions and some are becoming rather tortuous (A). The swollen glial cell (B) has compressed the myelinated fibre (C) which passes below it, while above B there are dilatations along a previously regular myelinated fibre (D).

At eight hours (Fig. 3) these changes have progressed. The tortuosity of fibre A is now more marked and is associated with expansion of adjacent glial cells. The myelinated fibre (C) in

Fig. 2 is now further compressed and that at D has almost disappeared, leaving myelin ellipsoids. Numerous other previously regular fibres show beading.

By 24 hours the field (Special Plate, Fig. 4) is almost unrecognizable when compared with Fig. 1. Many swollen glial cells have now disintegrated and added to the general debris scattered in the field, and myelin ellipsoids are prominent. Where isolated myelinated fibres yet remain their sheaths are distinctly thinner and less refractile than at the beginning of the test. This particular culture was observed for 69 hours, by which time marked disintegration had occurred (Special Plate, Fig. 5).

The degenerative changes occurring with motor neurone disease serum are essentially similar to those brought about by multiple sclerosis serum. The latter are, in general, faster than with motor neurone disease. In addition to myelin beading and fragmentation, a general dissolution of myelin similar to that reported by Bornstein and Appel (1961) commonly took place at the same time, the bright refractile lines seen in the bright field picture gradually becoming narrowed.

Minor changes are sometimes seen when normal serum is subjected to the same testing, and careful assessment of controls is necessary before concluding that any particular serum is toxic. Often these minor changes are temporary and reversible, while true toxic changes are progressive. Serum from two normal subjects did, however, produce well-defined toxic changes (see Table II). In some cultures which were tested with toxic sera well-defined growths of glial cells were seen on a more superficial plane than the layer of myelinated nerve fibres. The cells, which appeared to be astrocytes (though morphology is an uncertain criterion in assessing the nature of glial cells *in vitro*), seemed unaffected by the serum, which was so toxic for myelin and its immediately associated (? oligodendroglial) cells (Special Plate, Fig. 6).

Examination of Bodian preparations of the cultures which had been exposed to strongly myelotoxic serum (either motor neurone disease or multiple sclerosis) showed persistence of axis cylinders and nerve cells (Special Plate, Fig. 7), so that the inimical effect of the serum appeared very largely confined to myelin and the glial cells associated with it.

Discussion

The present work confirms the findings of Bornstein and Appel (1961) and of Bornstein (1963) regarding the toxicity of certain sera for myelinated fibres and glial cells in tissue culture. The pattern of demyelination, beginning with swelling of glial cells and myelin sheaths and progressing to the extrusion of myelin fragments while neurone bodies and axis cylinders are left relatively unaltered, is confirmed. In a later publication Bornstein and Appel (1965) reported demyelinating activity in 9 out of 15 sera from cases of amyotrophic lateral sclerosis. They point out that "the interpretation of tissue culture techniques . . . must be viewed with considerable scepticism in view of the general cytotoxic properties displayed by almost all sera, particularly when applied to short-term or young cultures." All sera in the present study were toxic to myelin-associated glial cells, as judged by their marked swelling followed by disruption in many cases. This was true even with the 25% serum tested and seemed similar to the toxic effects reported with undiluted serum by Berg and Källén (1962a, 1962b). Such effects were therefore regarded as non-specific. Recently Lumsden (1965) has reported nine "strong" positive results, five "weak," and three repeatedly negative out of 17 acute cases of multiple sclerosis, and this, too, confirms the reports summarized by Bornstein and Appel (1965).

Lumsden raises the important question "whether . . . (the antibodies at work in these tests) . . . are secondary products of the destruction of myelin." The present work confirms Bornstein's observation that myelotoxic activity is not confined to multiple sclerosis but is present, though possibly in lesser

degree, in amyotrophic lateral sclerosis. The possibility that myelotoxic antibodies may be formed secondarily to myelin destruction, arising perhaps from some non-immune cause, is a very real one in view of the finding of antibody to encephalitogenic factor in a variety of neurological conditions reported by Field, Caspary, and Ball (1963). Nevertheless, their undoubted occurrence in multiple sclerosis and experimental allergic encephalomyelitis might well be of some pathogenetic importance because of the lowered blood-brain barrier at lesions in this disease (Broman, 1947, 1964) and in experimental allergic encephalomyelitis (Barlow, 1956; Vulpé, Hawkins, and Rozdilsky, 1960; Field, 1961).

Finally, the occurrence of two examples of myelotoxicity among our 20 normal sera should be compared with the 2 out of 26 reported by Bornstein and Appel (1965). Lumsden (1965) does not report upon normals, but in view of the findings of Hashem and Barr (1963) it is possible that normal subjects may become sensitized to brain or its degeneration products if these are ingested. The possibility exists also that laboratory workers or others habitually exposed to brain might show some antibody responses. In this connexion Field *et al.* (1963) were able to demonstrate antibody to an encephalitogenic factor in 20.4% of 44 normal subjects. There is, however, no evidence that human antibodies demonstrated by the tanned-cell method parallel serum myelotoxicity.

Summary

Serum from 13 out of 21 undoubted cases of motor neurone disease showed demyelinating activity (at one in four dilution)

when allowed to act upon neonatal rat cerebellum cultures. Three out of five other cases in which the diagnosis was open to some doubt also showed this activity. Two out of 20 normal subjects also did so. All sera showed some degree of gliotoxic activity.

Myelotoxicity is not a property possessed exclusively by multiple sclerosis serum, though it may play a part in the pathogenesis of lesions.

We are much indebted to Dr. A. Vejajiva, of the Department of Neurology, Royal Victoria Infirmary, for serum specimens from cases of motor neurone disease and for providing details of the clinical condition of the patients.

We would also like to thank Miss Greta Joyce, A.M.I.L.T., for histological preparations; and Miss Marjorie Hall, A.M.I.L.T., Miss Catherine Smith, A.M.I.L.T., and Miss Christine Colligan for their valuable technical help in preparing so many cultures.

REFERENCES

- Barlow, C. F. (1956). *J. Neuropath. exp. Neurol.*, **15**, 196.
 Berg, O., and Källén, B. (1962a). *Acta path. microbiol. scand.*, **54**, 425.
 — (1962b). *Lancet*, **1**, 1051.
 Bornstein, M. B. (1963). *Nat. Cancer Inst. Monogr.*, No. 11, p. 197.
 — and Appel, S. H. (1961). *J. Neuropath. exp. Neurol.*, **20**, 141.
 — (1965). *Ann. N.Y. Acad. Sci.*, **122**, 280.
 Broman, T. (1947). *Acta psychiat. scand.*, Suppl. No. 46, p. 58.
 — (1964). *Acta neurol. scand.*, Suppl. No. 10, p. 21.
 Field, E. J. (1961). *Exp. Neurol.*, **4**, 233.
 — Caspary, E. A., and Ball, E. Janet (1963). *Lancet*, **2**, 11.
 Hashem, N., and Barr, M. L. (1963). *Ibid.*, **2**, 1029.
 Lumsden, C. E. (1965). In *Multiple Sclerosis—a Reappraisal* by D. McAlpine, C. E. Lumsden, and E. D. Acheson. Livingstone, Edinburgh.
 Vejajiva, A., Foster, J. B., and Miller, H. (1965). *Lancet*, **1**, 87.
 Vulpé, M., Hawkins, A., and Rozdilsky, B. (1960). *Neurology (Minneapolis)*, **10**, 171.

Congenital Rubella Syndrome as a Systemic Infection. Studies of Affected Infants Born in Philadelphia, U.S.A.

J. M. LINDQUIST,* M.D.; S. A. PLOTKIN,†‡ M.D.; L. SHAW,§ M.D.; R. V. GILDEN,† PH.D.;
 M. L. WILLIAMS,* M.D.

[WITH SPECIAL PLATE]

Brit. med. J., 1965, **2**, 1401–1406

Since the teratogenic potential of the rubella virus was established by Gregg in 1941 a syndrome of maternal rubella in early pregnancy followed by congenital heart disease, cataracts, and deafness in the infant has often been described (Manson *et al.*, 1960; Ingalls *et al.*, 1960; Lündström, 1962). Although this classical triad of sequelae to intrauterine rubella infection is well documented, other anomalies have received less attention. Recently, the birth of a large number of American infants with a variety of congenital abnormalities has demonstrated that foetal damage associated with maternal rubella is not confined to a few defects. The damaged infants were born six to nine months after an epidemic of rubella that was widespread in the United States from the end of 1963 to the summer of 1964 (Morbidity and Mortality Weekly Report, 1964; Plotkin, 1964; Phillips *et al.*, 1965). The congenital abnormalities which followed the epidemic included osseous lesions, hepatosplenomegaly, anaemia, and deficiency of megakaryocytes

in the bone-marrow. In addition, retardation of intrauterine growth, neonatal thrombocytopenic purpura, and rubelliform rash were seen with higher frequency than previously reported (Lündström, 1962; Berge *et al.*, 1963).

With the aid of recently developed procedures for virus isolation (Weller and Neva, 1962; Parkman *et al.*, 1962), Alford *et al.* (1964) showed that infants born after maternal rubella may excrete virus. In the recent epidemic their findings have been amply confirmed (Rudolph *et al.*, 1965; Cooper *et al.*, 1965; Plotkin *et al.*, 1965; Rubella Symposium, 1965). In several instances individuals in contact with infants shedding virus have developed rubella, thus raising an important public health and epidemiological problem (Cooper *et al.*, 1965).

The present paper reports the experience in a large city hospital in Philadelphia, Pennsylvania, during the period in which rubella-induced abnormalities were prevalent. Information is provided with respect to the incidence of congenital rubella syndrome in a predominantly negro population, the spectrum of anomalies in affected infants, the persistence of virus infection after birth, and the development of rubella antibodies.

* Department of Pediatrics, Philadelphia General Hospital.

† The Wistar Institute.

‡ The Children's Hospital of Philadelphia.

§ Department of Radiology, Philadelphia General Hospital.

E. J. FIELD AND D. HUGHES: TOXICITY OF MOTOR NEURONE DISEASE SERUM FOR MYELIN

FIG. 1.—Myelinated culture of newborn rat cerebellar cortex 21 days *in vitro*. Bright-light illumination. Twenty minutes after application of 25% serum from a case of motor neurone disease. Section maintained on heated stage at 35° C. Arrow X marks a clump of globules serving as a landmark. A, C, and D are normal myelinated fibres which are later affected. B is normal glial cell. ($\times 745$.)

FIG. 2.—Same field at four and three-quarter hours. Clump of globules (X) serves as a landmark. Same myelinated fibres are becoming tortuous (A), and this is in large measure due to expansion of glial cells. The glial cell (B) of Fig. 1 is now markedly swollen and shows granules clustered round its nucleus. The myelinated fibre (C), which passes below the cell B, is compressed. The fibre D above B, which was quite regular in Fig. 1, shows marked blebbing of myelin. ($\times 745$.)

FIG. 3.—Same field at eight hours. Note increased tortuosity of fibre A, and glial cell B beginning to degenerate. Myelinated fibre C is further compressed below the cell B, and the myelin of that at D has broken up with discrete ellipsoids. ($\times 745$.)

FIG. 4.—Same field at 24 hours, showing disintegration of many glial cells with much granular and myelin debris. Where short stretches of myelin persist they are thinned. ($\times 745$.)

E. J. FIELD AND D. HUGHES: TOXICITY OF MOTOR NEURONE DISEASE SERUM FOR MYELIN

FIG. 5

FIG. 5.—Same field at 69 hours, showing almost total disorganization of myelin. Nevertheless, axis cylinders are well preserved. ($\times 745$.)

FIG. 6.—Unharmful astroglial cells in culture which had been largely demyelinated by exposure to motor neurone disease serum (Figs. 1-5). These cells are more superficial in the culture and more towards the periphery. They are not actively associated with myelination. (Bodian stain. $\times 465$.)

FIG. 7.—Culture shown in Figs. 1-5 (motor neurone disease serum test). Nerve fibres impregnated by Bodian's method. Despite the extensive demyelination, nerve fibres (and nerve cells) are unaltered. Both thick axons (which were heavily myelinated initially) and the non-myelinated fibres are preserved, glial nuclei are heavily impregnated. (Bodian stain. $\times 1,245$.)

FIG. 6

FIG. 7

J. M. LINDQUIST *ET AL.*:
CONGENITAL RUBELLA SYNDROME
AS A SYSTEMIC INFECTION

Typical x-ray appearance of the lower limbs in rubella-syndrome infants, showing metaphyseal streaking and irregularity.