

fracture of cadmium-containing alloys, alkaline accumulators, cadmium-based pigments, or from any process where cadmium or cadmium-plated metal is heated to evolve fume. Cadmium is also used as a neutron absorber in nuclear reactors and is a constituent of some silver polishes.

Many of the workers exposed in the past to cadmium fume developed progressive dyspnoea, and some died after years of incapacity. Respiratory function tests have shown the presence of obstructive airways disease, and severe emphysema has been found in the lungs at necropsy.^{3 4} Most cadmium workers who become breathless have also developed proteinuria, while others have developed proteinuria without respiratory symptoms. While damage to the kidneys usually causes less incapacity than that to the lungs, some patients develop renal colic owing to the formation of stones, and a few patients have died from renal failure. The proteinuria in these patients is chronic, mild, and indistinguishable on electrophoretic examination from that occurring in patients with various renal tubular disorders.⁵ Concentration of the urinary colloids shows a relatively low albumin fraction with a predominance of α_2 and β globulins. Most of the proteins present are of relatively low molecular weight, and their appearance in the urine is believed to result from impaired reabsorption of proteins that are normally present in the glomerular filtrate.⁶ Other abnormalities described have been glycosuria, aminoaciduria of renal origin, hypercalciuria, and impairment of concentrating ability and of acid excretion, all of which point to dysfunction of the renal tubules.⁷

The manifestations of acute cadmium poisoning may be delayed some hours after exposure, whilst chronic poisoning usually develops insidiously at least two years after the start of exposure and occasionally some time after its cessation. The effects of this toxic element are irreversible at present, so that it is important to be aware of this uncommon condition and how to prevent exposure to cadmium fume and dust.

Autotransfusion

Recently in Nigeria A. Akinkugbe¹ has reported on the use of autotransfusion as a resuscitative measure in cases of ruptured ectopic pregnancy. Applicable to any case of severe intraperitoneal bleeding, autotransfusion is the simple expedient of collecting the patient's own shed blood from the peritoneal cavity, straining off any clot, and injecting it back into the veins. Especially in remote areas where blood donor or blood bank facilities are not available it can be a valuable adjunct to recovery.

Autotransfusion is not a new method of treatment, but, chiefly because donor's blood is usually available, it has lost favour. F. Stabler,² in 1934, reported the treatment of 13 patients with it, and F. P. Logan,³ in 1948, in South Africa, recorded 40 cases, with recovery in all. In recent years Chassar Moir^{4 5} has repeatedly advocated autotransfusion. He believes it is a valuable method of emergency resuscitation, especially in regions where blood banks are unknown. He states that he has used it "on many occasions" and "feels it a duty to demonstrate the method" to his students. He

quotes B. Markowski, who in British Honduras used autotransfusion in 100 cases, with survival in all but one.

A practical advantage of autotransfusion is the ease and speed with which it can be carried out, without the time-consuming grouping and cross-matching of blood. The blood which has collected in the abdominal cavity is simply ladled out with a galley pot or other suitable vessel and collected in a sterile flask containing 3.8% sodium citrate solution to make a proportion of nine parts of blood to one of solution. (If available, the "standard" bottle for blood donors containing 2 g. disodium citrate and 3 g. dextrose in 120 ml. water is convenient for the purpose.) When full the bottle is connected to the apparatus for intravenous infusion of saline set up before operation.

Only recently shed fluid blood is suitable for autotransfusion. But the method is intended for the collapsed patient with sudden intraperitoneal haemorrhage—the "acute" type of ruptured tubal pregnancy in which such blood is available in abundance. It is less suitable, and indeed it is not necessary, in the commoner "subacute" case (tubal mole), when the blood is old and firmly clotted.

Naturally, when donor blood is readily available, autotransfusion is apt to be forgotten, but this method of resuscitation, so simple and at the same time so effective, deserves to be better known and more widely used.

Construction of Vagina

The vagina may be congenitally absent or it may have been totally or partially excised for cancer of the uterine body, uterine cervix, or the vagina itself.

When the vagina is congenitally absent the uterus is usually absent also, and these patients have primary amenorrhoea. The ovaries are present and the girls are normally feminized. Occasionally—about 30 cases have been reported—a functional uterus is present which is capable of carrying a baby to term, and in fact has actually done so after a vaginal construction. Five to ten new cases of congenital absence are seen annually in the London hospitals which offer specialized surgery for this condition. Investigation of the patient should include intravenous pyelography to exclude associated abnormalities of the urinary tract, an examination under anaesthesia, and sometimes an exploratory laparotomy.

Congenital absence of the vagina is best treated by a McIndoe type of epithelial inlay operation.¹ A gynaecologist incises the perineum and dissects a retroperitoneal cavity. A plastic surgeon then introduces a rather thick partial-thickness skin graft from a thigh into this cavity round a prosthesis, and the labia minora are closed over it. When hard prostheses were used, fistulae, haematomata with loss of graft, urinary infections, renal stones, and extrusion of the vaginal mould were fairly common complications.² Soft prostheses are now available, the simplest of which is a cylindrically shaped piece of polyurethane plastic sponge, and with it postoperative complications are rare.³ As with all epithelial inlays, regular dilatation or constant maintenance of the original cavity size is necessary at first. The operation is therefore best performed three months before marriage, and once the graft has "taken"

¹ Akinkugbe, A., *J. Nigeria med. Ass.*, 1966, 3, 379.

² Stabler, F., *J. Obstet. Gynaec. Brit. Emp.*, 1934, 41, 768.

³ Logan, F. P., *S. Afr. med. J.*, 1948, 22, 793.

⁴ Moir, J. C., *Lancet*, 1960, 1, 60.

⁵ ———, *Operative Obstetrics*, 1964, 7th ed., p. 764. London.

¹ McIndoe, A. H., and Banister, J. B., *J. Obstet. Gynaec. Brit. Emp.*, 1938, 45, 490.

² Simmons, C. A., *Proc. roy. Soc. Med.*, 1959, 52, 953.

³ Jayes, P. H., *Ann. roy. Coll. Surg. Engl.*, 1966, 38, 210.

⁴ Pratt, J. H., and Smith, G. R., *Amer. J. Obstet. Gynec.*, 1966, 96, 31.