The reason for this preference for barbiturates is not clear, but it is noteworthy that the majority of the popular analgesics and salicylates are either wrapped as individual powders or supplied as tablets separated from one another between sheets of Cellophane or tinfoil. It is possible, therefore, that this method of packaging acts as a brake on those making impulsive suicidal attempts. By the time a sufficient dose of the analgesic drug has been extracted from its wrappings some of the emotional heat might have been dissipated, allowing time for wiser counsels to prevail.

It is so very much easier to open a bottle of tablets and swallow the lot in one impulsive gesture than it is to unwrap laboriously each pill carefully embedded in tinfoil. We would like to suggest, therefore, as a contribution to lessening the self-poisoning problem, that the drug firms should be asked to consider whether it would be possible to market all sedative and hypnotic drugs in this individually wrapped fashion. We do not know how such packaging compares in cost with the more conventional method of supply in cartons and bottles, but it is possible that any additional cost would be more than compensated for by the saving of lives and expensive hospital treatment.—We are, etc.,

F. A. WHITLOCK.
J. E. EDWARDS.

Department of Psychological
Medicine
Royal Brisbane Hospital,
Queensland, Australia.

REFERENCES

- James, I. P., Derham, S. P., and Scott-Orr, D. N., *Med. J. Aust.*, 1963, 1, 375.
- Buckle, R. C., Linnane, J., and McConachy, N., *ibid.*, 1965, 1, 754.
- Kessel, N., *Brit. med. J.*, 1965, 2, 1265, 1336.

Marihuana

SIR,—I am sorry that in his interesting review of *The Book of Grass* (22 April, p. 228) Dr. Griffith Edwards suggests that if one were in a position to start again from scratch it might be best to legalize marihuana rather than alcohol. I would not have thought it justifiable to consider them under the same category.

To the best of my knowledge no one smokes "reefers" for the intrinsic pleasure of the smoking as they do tobacco. The purpose of marihuana smoking is categorically that of intoxication, whatever form of dissociation this may take. If it did not intoxicate marihuana would not be used at all. On the contrary, sensible people take alcohol entirely for the urbane pleasure, in a social setting, of enjoying its taste in the various forms that it is presented. This is why, whether at table or in a pub, such sensible people take food at the same time as the alcohol—to minimize any possible intoxicant effect. I hope, then, that marihuana will never be legalized, alcohol never prohibited.—I am, etc.,

SEYMOUR SPENCER.
Headington,
Oxford.

Compulsory Treatment of Addicts

SIR,—I was pleased to read that Dr. M. M. Glatt (29 April, p. 309) confirms the opinion which I expressed to the Research Group of the Society of Medical Officers of

Health in June 1964 that "a kindly compulsion will be the first step in treatment and cure."

More than 30 years ago two great authorities on addiction, Sir William Willcox and Dr. Wolff, stressed the danger of addiction. Neglect of their advice throughout all these years has led to the appalling soul-slaughter and mind-slaughter of today, with the deaths of many precious lives intervening. Our community has been softened. We have been placed in jeopardy.

Dare we criticize our administrators for the £25 fine on a pedlar, who is thereby encouraged to sell his filthy drugs in greater quantity and to more victims in order to pay his fine?

Dare we commend to our administrators the 20–40 years' imprisonment, which is compulsory for pedlars in Ohio and which has almost eliminated drug addiction from that State?

Is there no one in this country who believes this to be the kind of "compulsion" which is needed?—I am, etc.,

Great Shelford,
Cambridge.

N. C. LONDON.

Benzodiazepines

SIR,—In your section "Today's Drugs" (1 April, p. 36) you refer to the addiction potential of the barbiturates, meprobamate, and the benzodiazepines. In vulnerable unstable personalities, such as the more psychogenic type of alcoholic, and in the individual who has previously been dependent on other drugs, the possibility of their becoming psychologically and physically dependent also on barbiturates¹ and meprobamate^{2,3} is very real.

As regards the benzodiazepines, you state that "so far, apparently no cases of addiction have been reported with the benzodiazepines, though, as with meprobamate, there may be some slight risk." In our experience the risk of the development of dependence in "vulnerable" personalities in the case of meprobamate seems certainly much greater than with chlordiazepoxide. We have, for example, given the latter drug (and diazepam to a somewhat lesser extent) to many hundreds of alcoholic inpatients and outpatients for short periods without coming across any clear-cut cases of dependence, whereas in a relatively short period we noticed a number of cases of psychological dependence with meprobamate within a short time of patients having started to take the drug,² so that, as with the case of barbiturates, we soon gave up its use for alcoholics. On the other hand, the risk of psychological and, rarely, of physical dependence cannot be completely excluded if chlordiazepoxide and diazepam are used for longer periods or in high doses, and cases of physical "addiction" have been reported from the U.S.A., where abrupt cessation led to severe abstinence symptoms.^{4,5} In a recent Austrian study among alcoholics treated as outpatients for a minimum period of four weeks Kryspin-Exner⁶ found cases of abuse and dependence in 6 patients (7.7%) among 78 patients treated with meprobamate, and in 4 patients (3.6%) among 111 patients treated with chlordiazepoxide, and in 7 patients (2.3%) among 302 alcoholics treated with diazepam. The author concluded from his study that cases of psychological and physical dependence undoubtedly occur both

with chlordiazepoxide and diazepam, but in a much milder form than in the case of barbiturates.

As regards nitrazepam, we have used the drug over the past two years in outpatient and inpatient practice as our routine hypnotic for psychiatric patients, finding it a reliable, effective drug, acceptable to patients, including those who had in the past relied on barbiturates. Although the drug has been given to a great many alcoholics, so far no cases of dependence have occurred, and as a rule there was little difficulty in getting these people off the drug. Of a number of non-barbiturate hypnotics investigated by us over the past 14 years⁷ nitrazepam seems so far the most satisfactory. On the other hand, it would be most surprising if such an effective drug as this will not sooner or later be found to produce dependence, at least in psychologically vulnerable personalities.—I am, etc.,

St. Bernard's Hospital,
Southall, Middlesex.

M. M. GLATT.

REFERENCES

- Glatt, M. M., *Bull. Narcot.*, 1962, 14, No. 2, 19.
- *Brit. med. J.*, 1957, 1, 164.
- Haizlip, T. M., and Ewing, J. A., *New Engl. J. Med.*, 1958, 258, 1181.
- Lesse, S., *J. Neuropsychiat.*, 1962, 3, 154.
- Hollister, L. E., Motzenbecker, F. P., and Degan, R. O., *Psychopharmacologia (Berl.)*, 1961, 2, 63.
- Kryspin-Exner, K., *Brit. J. Addict.*, 1966, 61, 283.
- Glatt, M. M., *Psychiat. et Neurol. (Basel)*, 1966, 152, 28.

Pulmonary Artery Thrombosis and the Nephrotic Syndrome

SIR,—I was very interested in Dr. I. S. Menon's finding (8 April, p. 110) that the fibrinolytic activity of renal-vein blood is high. This accords with histochemical demonstrations that plasminogen-activator activity is high in the kidney. I believe this to be of teleological importance in that the kidney may be required to protect itself against fibrin which, having been formed in the general circulation, has been swept into the glomeruli, whose filtration may thus be blocked.

In conjunction with Dr. G. Taylor, I have been studying serum taken from patients with acute renal failure or shock by immunoelectrophoresis against a split fibrin anti-serum. Fifteen of 17 cases of acute renal failure were positive for fibrin products, as were 5 of 7 cases of shock. None of the 10 cases of chronic uraemia were positive, but 3 of 20 normal sera were. The finding in the normal patients is not discouraging but rather indicates the sensitivity of this qualitative method, and agrees with the quantitative method of Merskey, Kleiner, and Johnson¹ with regard to normal patients who have increased fibrinolytic activity. Since when uraemia is established fibrinolysis is depressed, the findings lend support to the theory that disseminated intravascular coagulation has occurred. We hope to report the full details elsewhere.

That any type of "ischaemic" renal failure might be the result of defibrination can be expected from the well-known associations in obstetrics of renal failure with defibrination, from the fact that the list of causes of acute renal failure is the same list as the causes of defibrination, and from the evidence that in the experimental Schwartzman reaction renal cortical necrosis is induced.