(13 January, p. 109). I wanted and still want to warn the others. I have my suspicions, too, about you, Mr. Editor, as the result of your putting out a special supplement on the royal malady complete with vivid purple-coloured cover!—I am, etc.,

C. E. DENT.

University College Hospital
Medical School,
London W.C.1.

REFERENCE

- ¹ Waldenström, J., and Vahlquist, B., *Acta med. scand.*, 1944, 117, 1.

Steroids and Nephrotic Syndrome

SIR,—Dr. I. G. Wickes (16 March, p. 706) claims that the regimen he uses will identify those children with nephrotic syndrome who are "likely to relapse." He instructs the mothers to test their child's urine with Esbach's reagent and to record the result. However, this does not predict anything; it merely records the relapse or the degree of proteinuria. We know of no way of predicting which nephrotic child will run a relapsing course; neither renal biopsy nor the selectivity of proteinuria will do this. Dr. Wickes then advocates that the mother should titrate the dose of steroid against the amount of proteinuria, maintaining that in this way the increasing proteinuria that may be associated with colds and other infections is kept in check. We do not doubt that the proteinuria goes during this regimen, but believe that it would often disappear without any therapy. Our experience accords with that of Arneil and Lam.¹ Very often the proteinuria goes as the child recovers from the respiratory infection, and therefore it is better to withhold steroids for one or two weeks, provided that oedema does not develop. Withholding therapy avoids needless steroids, and parents agree to it willingly, since the child does not necessarily feel ill at the time of the proteinuria.

As he suggests, a simple record of proteinuria kept by the parent can be very helpful for outpatient supervision. However, Esbach's test seems a rather cumbersome one to inflict on parents, particularly as it takes 24 hours to read. Albustix is a simple and reliable test² which a parent or child can use readily at home, recording the result in a few seconds. When demonstrating the method we warn them that trace and transient one-plus readings are unimportant.—We are, etc.,

S. R. MEADOW.

R. H. R. WHITE.

Department of Paediatrics
and Child Health,
Birmingham University.

REFERENCES

- ¹ Arneil, G. C., and Lam, C. N., *Lancet*, 1966, 2, 819.
² Rennie, I. D. B., and Keen, H., *Lancet*, 1967, 2, 489.

Detection of Bacteriuria

SIR,—Dr. J. M. Bradley and others in their paper (16 December, p. 649) have misquoted the results we obtained in our assessment of the triphenyl tetrazolium chloride reduction test for the detection of significant bacteriuria.¹ Their statement that in our series "only 44% of infections in the 10⁴–10⁹ organism per ml. of urine range were

detected" referred only to the ability of the test to detect infections with Gram-positive organisms, and this was made clear in our paper. As we also made clear, when patients on antibiotic therapy were excluded, the triphenyl tetrazolium chloride (T.T.C.) test picked up 12 of 24 infections with Gram-positive organisms, and these included, besides a number involving *Streptococcus faecalis*, infections with diphtheroids, micrococci, and *Candida albicans*, infections which the method described by Dr. Bradley and colleagues, depending as it does on the use of one MacConkey plate only, must inevitably miss.

When all urines with significant bacteriuria (10⁵ or more bacteria per ml.) encountered in a general hospital are considered the detection rate for the T.T.C. test was 182 of 258 specimens or 70.5%. Our paper further pointed out that in a selected population, such as in obstetrical patients, where 79% of all infections are due to *E. coli*, the detection rate of the T.T.C. test can be bettered, and can detect as many as 90% of cases of significant bacteriuria.

No chemical test, of course, can replace accurate quantitative culture of urine, but, as Dr. Bradley and colleagues point out, there is a real need for a simpler, less expensive screening method. We found, and reported, the T.T.C. test to be a much better screening method than the quotation of our paper would suggest. A direct comparison between the T.T.C. test and the method of Dr. Bradley *et al.* is not possible. Their graph (Fig. 2) shows a reasonable correlation between the number of colonies counted by their method, and a true quantitative culture in 20 specimens, none of which apparently had more than 10⁵ bacteria per ml. The graph (Fig. 4) showing the correlation between counts obtained with a standard loop and their screening method also shows a correlation, but in fact their paper does not state how many of a given number of positive urines can be screened out, nor how many false-positives are obtained, and the comparative values of the T.T.C. method (which is simpler to perform) and their screening method remain to be assessed.—I am, etc.,

ELLEN VAN DER HOEVEN.

Hôtel Dieu Hospital,
Kingston,
Ontario,
Canada.

REFERENCE

- ¹ Hinton, N. A., and van der Hoeven, E., *Canad. Med. Ass. J.*, 1965, 93, 639.

Acute Iron Poisoning in Children

SIR,—Drs. D. G. D. Barr and D. K. B. Fraser (23 March, p. 737) report further tragedies from swallowing ferrous sulphate. This is the salt of a weak base and strong acid. Its solution has a pH of 3.7 and readily damages any mucosa with which it is long in undiluted contact. Many neutral salts are now available which small children may swallow by the dozen without harm. Often these are damned because they cost 2d. a week more than ferrous sulphate, but with increasing use of desferrioxamine even this miserable economic argument may fail. Is it not time to abandon and to ban the use of ferrous sulphate, which is a lethal poison?—I am, etc.,

T. H. HUGHES-DAVIES.

Downton, Wilts.

Q Fever

SIR,—The account of Q fever in Northern Ireland by Dr. J. H. Connolly (2 March, p. 547) prompts us to report our own experience in the period 1960–7. During these eight years we have made a laboratory diagnosis of Q fever by serological tests or by isolation of *Rickettsia burneti* from 24 patients (and confirmed it in two additional cases from other laboratories). They resembled closely those reported by Dr. Connolly in age and sex (see Table) and in seasonal distribution.

Age and Sex Distribution

Age Group	Male	Female	Total
10–29	4	1	5
30–49	6	4	10
50–69	5	2	7
70+	2	0	2
All ages	17	7	24

The acute illness of two patients began in the first quarter of the year, of 10 in the second, one in the third quarter, and six in the fourth. The types of illness were: pneumonia, 10 cases (one chronic with probable endocarditis); fever, eight cases (one severe with vomiting, dehydration, and rash; one post-partum); subacute endocarditis, four fatal cases, all male; pericarditis, two cases, one with pleuropneumonia, one with aortitis and syphilitic aneurysm. The last patient, the four with subacute endocarditis, and the patient with pneumonia and probable endocarditis all had high complement fixing antibodies to both phase 1 and phase 2 antigens, as had also an additional case of endocarditis in which we made confirmatory tests for another laboratory. Serious cardiovascular diseases have thus been more prominent in our series than in Dr. Connolly's, but pericarditis was found in both series. Tests for Q fever as well as for Coxsackie virus infection therefore seem appropriate in cases of acute pericarditis.

Contact with farm animals (mostly bovine) or their products was traceable in most cases. Six were farmers, three had driven trucks loaded with dung, one had cleared dung from cattle wagons, and three used dung as gardeners; three had consumed unpasteurized milk and one of these lived beside a farm; one office worker frequently visited a byre; an electrician developed pneumonia with serological evidence of both Q fever and a psittacosis-group infection after working in an abattoir¹; another case of pneumonia developed after working in fields near lambing sheep. No contact was traced in three Glasgow patients or three from the counties of Renfrew, Dumfries, and Kirkcudbright; in each of these counties Q fever is known to infect cattle. Infection of sheep and cattle in the west of Scotland has been recognized for more than a decade² and continuing surveillance tests in collaboration with Dr. T. F. Elias-Jones continue to reveal *R. burneti* in samples of unpasteurized milk. We agree, however, with Dr. Connolly that inhalation of particles derived from farm animals or their products seems a more important source of infection than ingestion of infected milk.

The finding of false positive complement fixation antibody responses to Q fever in some children with infections due to adenoviruses types 7 and 21 was reported at the meeting of the Society for General Microbiology, Janu-