High-dosage Fluphenazine

SIR,—Comment in your correspondence column has drawn attention to side-effects of phenothiazines,¹⁻³ and there is current interest in the use of fluphenazine enanthate.⁴ I wish to draw attention to the value of high-dose oral fluphenazine administration in acute schizophrenic illness. This therapeutic approach to a broad cross-section of psychotic patients has been studied by some French psychiatrists^{5,6} but has not been recorded in publications in English. The French authors comment on the remarkable tolerance of doses up to and even exceeding 500 mg. daily. They observed that side-effects of medication were definitely fewer in high doses as compared with those in doses of less than 50 mg. daily. In their opinion the clinical response to high doses was excellent. When evaluating dosage of fluphenazine one should bear in mind that it is 10 to 20 times as potent as chlorpromazine.⁷

Because of these favourable reports I have recently treated newly admitted patients who are in my opinion suffering from an acute schizophrenic illness by administering an initial dose of oral fluphenazine 100 mg. daily, and this is increased to 200 mg. daily within 48 hours and in one case to 400 mg. daily, the dose level being increased until there is evidence of response by psychiatric interview and observation of ward behaviour. Routine anti-Parkinsonian drugs have been given, either benzhexol 5 mg. three times daily, procyclidine hydrochloride 5 mg. three times daily, or benztropine methanesulphonate 2 to 4 mg. daily. I have observed 25 patients given doses of fluphenazine 100 mg. or in excess of this daily. There have been 12 females and 6 males given 100 mg., 4 females and 2 males given 200 mg., and 1 female given 400 mg. The only side-effects noted as possibly being due to the fluphenazine were excessive salivation in one female on 200 mg. daily, drowsiness and motor retardation in one female on 100 mg. daily, and slight jaw stiffness in one female on 100 mg. daily. The absence of any serious side-effects has been most impressive, and the lack of any persistent sensations of discomfort from the patients. In my observations of the therapeutic response I have been impressed by the anti-psychotic, socializing, and energizing properties of the drug and by the rapidity of improvement as compared with other phenothiazines.

I would like to note one illustrative example: a female aged 40 years with previous psychiatric inpatient admissions was brought to hospital by the police under the provisions of section 136 of the Mental Health Act. On examination she was thought-disordered, showed incongruity of affect, was deluded, excited, and aggressive. Fluphenazine was given, increasing to 400 mg. by the fourth day, and she was by then much improved and was discharged after three weeks. This response was more rapid and complete as compared with that recorded on previous admissions.

In conclusion fluphenazine appears to have this unusual property of being well tolerated in very large doses, provided anti-Parkinsonian agents are given, and the therapeutic response is such that further critical evaluation is indicated in planned controlled trials.—I am, etc.,

P. W. RICHMOND.

Banstead Hospital,
Sutton,
Surrey.

REFERENCES

- 1 Kimerling, J. J., and Patel, S. R., *Brit. med. J.*, 1967, 4, 554.
- 2 Rogers, J. B., *Brit. med. J.*, 1967, 4, 235.
- 3 Keat, E. C. B., and Toghiani, P. J., *Brit. med. J.*, 1967, 3, 867.
- 4 Capstick, N., *Brit. med. J.*, 1968, 1, 181.
- 5 Charriot, G., Lambert, P. A., Delauney, J., and Marcon, G., *Congrès de Psychiatrie et de Neurologie de Langue Française*, Marseilles, 1964.
- 6 Bouillier, H., *Congrès de Psychiatrie et de Neurologie de Langue Française*, Marseilles, 1964.
- 7 Ayd, jun., F. J., *J. med. Soc. N.J.*, 1960, 57, 4.

Basilar Syndrome

SIR,—If Dr. A. J. Macleod (23 March, p. 767) cares to glance at p. 174 of the fourth edition of my *Textbook of Orthopaedic Medicine*, volume I, he will find a discussion on the dangers of cervical manipulation. My calculation is that death is brought about in not more than 1 in 10 million manipulations of the neck. On the following page, under the heading of "Danger of Not Manipulating," the disadvantage is emphasized of leaving a central disc protrusion where it is, to draw out the very osteophytes that eventually compress the spinal cord.

Provided manipulation is carried out using the techniques of orthopaedic medicine, and is confined to patients selected on the criteria set out in volume I, no untoward results are to be expected, and neither I nor any of the thousand physiotherapists I have trained in my time have (so far) caused lasting damage, let alone death.—I am, etc.,

JAMES CYRIAX.

London W.1.

REFERENCE

- 1 Cyriax, J., *Textbook of Orthopaedic Medicine*, 1962, Vol. I, p. 174, 4th ed., London.

Screening Tests for Phenylketonuria

SIR,—I would like to comment on your leading article "Screening Tests for Phenylketonuria" (16 March, p. 656) and on the report in the same issue from Dr. James Farquhar and others in Scotland (16 March, p. 674).

The latter report has set out in clear terms the advantages offered by the Guthrie bacterial inhibition test over other screening procedures. Perhaps the most important point made by this report is the simplicity of the administrative system and its fluidity which would enable it to be applied to other tests which may in time supersede or supplement the proposed tests. A note of foreboding is, however, struck in the last paragraph of your leading article, which seems to me to advocate following the American system of making newborn testing for phenylketonuria a mandatory procedure. You cite Connecticut as having taken this step, whereas in fact the procedure has been enjoined by law on physicians and hospitals in at least 14 of the United States.¹ It would have been well to state that much of the criticism of mass newborn screening which has developed in America—some of which your leading article alludes to—has stemmed from this legal enforcement. One hopes that authorities on this side of the Atlantic will be slow to follow suit and that the medical profession will give careful thought to the alternatives and likely sequelae before advocating this step.

Perhaps the experience gained in the Republic of Ireland may show that there is more to be gained by seeking rather than demanding the co-operation of hospitals, doctors, and nurses. For the past two years a scheme set up by the Department of Health has been in operation by which the Guthrie test for phenylketonuria has been made available on all newborn infants. The scheme has been free of charge and without emolument to the individuals collecting specimens. The results of the first 16 months of testing are shortly to be published² and will not be anticipated here except to point out that since testing began on 1 February 1966 a coverage amounting to 80% of the newborn population has been achieved to date. Tests are carried out in a small central laboratory in this hospital with a small staff, and local organization and co-ordination are in the hands of the county medical officers of health. The experience gained has prompted the hope that when the results of the initial period are disseminated they will encourage those concerned with the newborn to achieve a higher and possibly full coverage.—I am, etc.,

SEAMUS F. CAHALANE.

Department of Pathology,
Children's Hospital,
Dublin.

REFERENCES

- 1 *State Laws Pertaining to Phenylketonuria*, U.S. Department of Health, Education, and Welfare, 1965.
- 2 Cahalane, S. F., *Arch. Dis. Childh.*, in press.

Lower Limb Injuries

SIR,—Dehne and Immermann¹ collected from the literature and from their own experience a total of 42 cases of fracture of the shaft of the femur combined with dislocation of the ipsilateral hip. In only 15 of these was the diagnosis of the hip dislocation made at the initial examination. In the majority of cases in which the dislocation was not discovered at the initial examination the diagnosis was not made until four to six months later.

Increasing attention has been paid in recent years to the importance of diagnosing proximal injuries in the region of the hip and pelvis in the presence of a more distal injury to one of the lower limbs.² The patient may not make specific complaint of pain in the region of the injured hip. The surgeon directs his attention to the treatment of the distal injury—commonly fractures of the tibia and fibula—and the proximal injury is missed, often for weeks and sometimes for many months. The proximal injury may be a fracture or fractures of the pelvis, a fracture-dislocation of the hip joint, an intracapsular fracture of the neck of the femur, or a trochanteric fracture. The distal injury is usually fracture of the tibia and fibula, often compound, but it may be a foot or ankle injury, an injury to the knee such as compound fracture of the patella, or, as in Dehne and Immermann's series, a fracture of the shaft of the femur. Combined injuries of this sort are commonly seen in motorcyclists and pillion passengers. Failure to diagnose the proximal injury within a matter of hours or days of the accident often results in quite a marked avoidable disability in addition to any disability inherent in the diagnosed distal injury. The aggrieved patient may sue his surgeon for negligence and the

claim may well be indefensible. The cost of cases of this sort to the medical defence organizations is considerable; the reputation of the surgeon is tarnished; and, most important of all, the patient suffers an avoidable permanent disability.

The remedy is simple. Routine x-ray examination of the hips and pelvis should be made before any treatment is given for a sizable injury to a lower limb sustained in a violent accident such as a road traffic accident. The surgeon must not expect that the patient will identify for him the nature and positions of his injuries. The proximal injury in the region of the hip or pelvis is commonly painless or relatively painless in comparison with the pain of the more obvious distal injury. The lesson should have been learned by now, but it has not. This letter has been written to encourage the idea that routine x-rays of the pelvis and hips should be taken to exclude "the proximal injury."—I am, etc.,

Liverpool 1.

NORMAN ROBERTS.

REFERENCES

- ¹ Dehne, E., and Immermann, E. W., *J. Bone & Surg.*, 1951, 33A, 731.
- ² Roberts, N. W., *Modern Trends in Orthopaedics*, 1962, 3, chapter 7, p. 147. Edited by J. M. P. Clark. London.

Human Heart Transplantation

SIR,—In reviewing the record "Human Heart Transplantation" by Professor C. N. Barnard and his team (2 March, p. 566), Sir John McMichael seems either to have approached his subject with preconceived prejudice or to have missed the point of heart transplantation entirely.

As has been repeatedly stated by Professor Barnard, heart transplantation is only considered when the recipient is already dying and when further medical or more conservative surgical treatment will be to no avail. Professor V. Schrire stated the indication for heart transplantation very clearly on the record—namely, irreversible, intractable myocardial failure, with the patient "literally at the end of the road" and "dying in front of your eyes." Whatever the cause of the myocardial failure (Coxsackie myocarditis, coronary artery disease, etc.), the indication of extreme pump failure remains the same. The reviewer's question: "Is his life, with continuing dependence on drugs, going to be better than could be achieved with ordinary treatment?" is quite superfluous, as there is no alternative for the patient except death. The problems he envisages "surrounding living donors" are even more inappropriate. If the reviewer does not realize that the donor is of necessity already dead, then the whole question of heart transplantation must indeed pose for him incomprehensible and insuperable problems.

In the case of any transplant patient, whether heart or kidney, the consideration is not only how long the patient will live, but also whether the remaining lifespan is made more tolerable. So far, Professor Barnard's second heart transplant patient, who was a complete invalid facing imminent death, is reported to be gradually returning to normal life. Perhaps he is the best judge of whether the operation was worthwhile or not.—I am, etc.,

Pretoria,
South Africa.

E. J. SCHULZ.

F

Hypermobility Joints

SIR,—In your leading article on Hypermobility Joints (9 March, p. 596), the familial incidence of this condition was mentioned.

During a recent investigation we examined a large number of patients with simple joint hypermobility. Many of them had affected relatives, and it is our impression that familial joint laxity is a definite entity which is transmitted as an autosomal dominant. One striking feature was the consistency of the condition within any kindred. For example, in one family with 28 affected members in five generations, there were orthopaedic problems ranging from pes planus to congenital dislocation of the hip. On the other hand a professional contortionist from an old circus family claimed that none of her affected relatives ever had joint troubles.

At the present time it is impossible to state whether these kindreds have two different forms of familial joint hypermobility, or whether they are at the opposite ends of the spectrum of a single entity.—We are, etc.,

St. Thomas's Hospital,
London S.E.1.

P. H. BEIGHTON.

St. Bartholomew's Hospital,
London E.C.1.

F. T. HORAN.

Injury from Liquid Propane

SIR,—I wish to report an unusual and unexpected industrial accident, one of my patients being involved.

Two men were moving a cylinder of propane. One, the patient, was holding the valve wheel, while his colleague clasped the bottom of the cylinder. The outlet was directed upwards. In the course of the removal the man clasping the bottom of the cylinder changed his grip, rotating the cylinder about the valve stem, with the result that the valve opened and ended up pointing downwards. The patient's right thigh was drenched in liquid propane, which produced a large area of second degree frostbite. Had the valve outlet initially been facing downwards the resulting accident might well have been a disaster.—I am, etc.,

Great Bedwyn,
Marlborough, Wilts.

H. J. FENN.

Volvulus of the Sigmoid Colon in Africa

SIR,—I was pleased to see your reference to the concept of geographical pathology in the leading article "Volvulus of the Sigmoid Colon" (3 February, p. 264). However, you, as many others, have erred in considering the Negro population of Africa as a single group. You correctly note the reports from several East Africa countries in support of your statement regarding the high incidence of sigmoid volvulus among the Negro population of Africa. There is, however, considerable variation in the incidence of many diseases among the Africans in different parts of this huge and diverse continent.

At the University College Hospital, Ibadan, Nigeria, sigmoid volvulus is very uncommon. A recent review of 794 cases of intestinal obstruction at this hospital showed only eight cases (1%) were due to sigmoid volvulus,¹ a figure similar to that you quote for the incidence in the British Isles. The leading causes

of intestinal obstruction here are hernia and intussusception. In contrast to these observations, a report from Ghana² indicates that they have a high incidence of sigmoid volvulus. To confuse the problem even more, personal experience in over 1,000 barium-enema examinations in Ibadan has shown that a redundant, mobile sigmoid loop is very common here. Perhaps this suggests that a combination of several factors is responsible for this condition.

In summary, this letter is to record the low incidence of sigmoid volvulus, not in Nigeria, not even in all of Western Nigeria, but at University College Hospital, Ibadan, drawing patients mainly from in and around the city of Ibadan. Only by being specific about the populations concerned can data relating to geographical pathology help in solving some of the riddles we face in medicine.—I am, etc.,

STANLEY P. BOHRER.

Department of Radiology,
University College Hospital,
Ibadan, Nigeria.

REFERENCES

- ¹ Solanke, T. A., personal communication.
- ² Cole, G. J., *Gut*, 1965, 6, 151.

Panhypopituitarism after Cured Tuberculous Meningitis

SIR,—Is it not possible that the patient described by Dr. V. K. Summers and others (10 February, p. 359) at the age of 27 could have suffered from acute infectious cavernous sinus thrombophlebitis, with hypopituitarism as a complication? This complication was described, for instance, by Walsh¹ and Williams.²—I am, etc.,

J. A. M. FREDERIKS.

Binnenziekenhuis,
Eindhoven, Holland.

REFERENCES

- ¹ Walsh, F. B., *Arch. Ophthalmol.*, 1937, 17, 46.
- ² Williams, E., *Proc. roy. Soc. med.*, 1956, 49, 827.

Overseas Service

SIR,—“Go West, young man,” is advice followed by young doctors for many obvious reasons. May I try to interest them in a less crowded professional world in the South—in the sun? The developing countries have medical needs described by two statistics of shortage: health expenditure varies between 3s. and 20s. per head per annum, compared with 600s. for England and Wales; the doctor-patient ratio is between 1:20,000 and 1:100,000; 1:1,200 in England and Wales.

Why go South? Humanitarian reasons? Dr. Maurice King, of Makerere University, Kampala, in his preface to *Medical Care in Developing Countries* makes out the practical case for short-term service after intern jobs and before the children require secondary school education. Two major factors in any doctor's mind are keeping a place in the professional competition for good jobs and the financial needs of a new family. Finance need not be a problem. Overseas contract workers are better off than their equivalents at home in housing, transport, recreation, and salary. Opting out from the competition is a problem, but there is a partial solution already in existence if only it were applied more often. Whenever some