old, and both she and her sister-in-law were clear that difficulty in swallowing was her main problem.

The patient looked ill and wasted, and was admitted as an emergency, with a probable diagnosis of carcinoma of the oesophagus. Alternatively, it was considered that she may have had enlarged nodes compressing her oesophagus and causing the dysphagia. In the ward she admitted to frequency, as she had been drinking a lot to try to wash her food down. She also had nocturia. Her pulse was regular at 92 and her blood pressure was 150/80. Apart from wasting there were no abnormal signs. A barium meal report stated, "the oesophagus showed no evidence of obstruction and had normal peristaltic movements, but there is a slight irregularity of the margin extending throughout the whole length—? significance. The duodenal cap appeared normal. There is a small pouch arising from the lesser curve border of the antrum just proximal to the pyloric canal; this may represent ectopic pancreatic tissue," and the chest x-ray was normal.

There was no evidence of diabetic retinopathy or neuropathy. However, she was found to have glycosuria and ketonuria on routine ward testing. Blood sugars were 350 mg./100 ml. and 540 mg./100 ml., and her E.S.R. was 12 mm. in one hour. The rest of her blood picture was normal, as was a midstream urine specimen. Her blood urea was 41 mg./100 ml., bicarbonate 24 mEq/l., potassium 3.9 mEq/l., and sodium 134 mEq/l. A diagnosis of diabetic ketosis was made, and she was started on soluble insulin. Her dysphagia disappeared within hours of controlling her diabetes.

This case is therefore reported as a most unusual presentation of diabetes mellitus.

I am grateful to Dr. H. E. Webb for allowing me to report this case.

—I am, etc.,

L. HERZBERG.

St. Thomas's Hospital,
London S.E.1.

Androgens and Growth

SIR,—The caution displayed by your author in his review of androgens and anabolic steroids (19 April, p. 165) is well founded. I do not share Dr. R. Greene's (10 May, p. 379) optimistic view of the effects of such treatment. That androgens promote skeletal maturation to a greater extent than linear growth has been oft repeated but is not a fallacy. How would Dr. Greene explain the reduction in adult height which is commonly seen in constitutional sexual precocity, or in the simple virilizing form of congenital adrenal hyperplasia which is untreated or inadequately treated?

Many eminent paediatric endocrinologists¹⁻⁴ have expressed reservation about the benefit of androgen therapy in growth disorders. Laron⁵ finds that the degree of response of children to various anabolic steroids varies in an unpredictable manner. Children with primordial or constitutional dwarfism, chondrodysplasia, and gonadal dysgenesis respond poorly. While psychological benefit may be produced by inducing a growth spurt or signs of sexual maturation in a short male with delayed puberty, therapy

should be given only in short 3–6-month courses and treatment should be interrupted for periods of 6–12 months to allow an adequate period of time in which to observe the effects on skeletal maturation.—I am, etc.,

P. H. W. RAYNER.

Department of Paediatrics
and Child Health,
University of Birmingham,
Birmingham 16.

REFERENCES

- Wilkins, L., in *Current Therapy*, edited by H. F. Conn, 1952, p. 367. Philadelphia, Saunders.
- Prader, A., and Illig, R., in *Symposium on Protein Metabolism*, 1962, p. 383, edited by F. Gross. Springer, Berlin.
- Hubble, D. V., in *Recent Advances in Paediatrics*, 3rd ed., edited by D. Gairdner, 1965, p. 211. London, Churchill.
- Blizzard, R. M., in *Human Growth*, edited by D. B. Check, 1968, p. 53. Philadelphia, Lea & Febiger.
- Laron, Z., in *Somatic Growth of the Child*, edited by J. J. Van der Werff ten Bosch and A. Haak, 1966, p. 287. Leiden, Stenfort Kroese; Springfield, Illinois, Thomas.

Sexual Mores and V.D.

SIR,—The reviewer of *Sex and Society in Sweden* (31 May, p. 564), after noting with approval the "enlightened" and "liberal" sexual mores of Swedish society, seems perplexed by the increasing incidence of syphilis and gonorrhoea, especially in teenagers, and also by the increase in illegitimate births and requests for abortion. As an explanation he suggests that "some sort of devil" must be frustrating the good work, or that the above and other disturbing features of Swedish society, such as the high incidence of alcoholism and the highest suicide rate in the world, are merely external manifestations of "hidden internal conflicts." Is not the simpler explanation of some causal interrelation between the various phenomena worthy at any rate of consideration?—I am, etc.,

London W.1.

D. H. PATEY.

Disodium Cromoglycate and Induced Asthma

SIR,—In their letter (10 May, p. 382) Professor J. Pepys and Dr. F. E. Hargreave criticize the (mostly negative) results of Miss Helga Bewersdorff and myself (26 April, p. 220) in our attempt to protect patients against allergen-provoked asthma with disodium cromoglycate. They say our cases F and G should not have been included according to our own standards. We have included attacks lowering the vital capacity by 10 to 20% or the peak flow by 30 to 40%, not "and," as was said erroneously in our paper. We felt justified including these patients, particularly as they became audibly wheezy during their attacks. The variations in our cases I, J, and L, which Professor Pepys and Dr. Hargreave mistakenly regard as evidence of protection, are often seen in bronchial provocation tests. I have carried out such tests since 1948 and regard these variations as normal. The difficulty of judging this is illustrated by our patient D, who seemed to show definite partial protection when given 16.8 mg. cromoglycate, but none when given 8.4 or 33.6 mg. Despite this rather doubtful result we listed him as partially protected.

In the matter of the F.E.V. as a standard measurement of lung function we had referred to an earlier paper.¹ My scepticism is based on the fact that in the asthmatic every forced expiration starting from maximum inspiration is likely to increase bronchial spasm. I have illustrated this phenomenon with a spirometric recording.² Moreover, as is well known, in an asthmatic the forced expiration of coughing usually increases bronchial spasm. The same phenomenon occurs when the forced vital capacity is recorded, and also, to some extent, with the F.E.V. Both are therefore often reduced in asthmatics after the first measurement. Thereafter, they may seem well reproducible, and this is presumably the example Professor Pepys and Dr. Hargreave refer to. But the trouble is that the degree of spasm provoked may change from subject to subject and from day to day. I therefore regard the F.E.V. as a poor lung function test in asthmatics, almost as poor as the maximum ventilatory capacity which was fashionable some years ago but has now been abandoned in the main. The F.E.V. fashion deserves the same fate. In any case, Professor Pepys and Dr. Hargreave would have done well to use a second method as a safeguard—for instance, the peak flow.

Their statement that the coarse-particle powder of cromoglycate they used had, in their hands, a potent inhibitory effect (and therefore must have been working) begs the question and needs no discussion. Both Professor Pepys and Dr. M. C. S. Kennedy (24 May, p. 511) ask for more details of our observations on the five chronic asthmatics. In these patients we attempted to improve lung function with single or repeated doses of cromoglycate aerosol, but had completely negative results in three or four experiments with each subject. We reported this negative result very briefly in order to save space. Whatever results may be obtained with cromoglycate used for several weeks, it seems clear that a beneficial effect cannot be demonstrated in an acute experiment in chronic asthma.

To sum up: The reasons for the differences between our results and those of Professor Pepys and his colleagues are not clear. They found "complete" protection in nine out of 10 patients and partial protection in the tenth. We found complete protection in only two out of 13 and partial protection in one and possibly in two others. There were important differences in the methods of administration, methods of testing, and (except in one patient) different allergens. Other sources of error may be hidden in our ignorance about the mechanism of action of the cromoglycate. It may have been a little disconcerting for Professor Pepys and his colleagues to find that we have had different results, but in the circumstances we must agree to disagree. I hope that this disagreement will lead to a more sober appreciation of this new and interesting substance.

To Dr. J. M. Smith (31 May, p. 574) we reply that patients with mild asthma, who form a great proportion in his investigation, are not suitable for drug trials, as their asthma too often spontaneously improves. As to the coarseness of the cromoglycate particles (Dr. J. S. G. Cox and others, 7 June, p. 634), we have sucked the powder through the spinhaler on to a cover slip and have found microscopically about the same amount of very large conglomerates as when we examined the contents of the cromoglycate capsules. The spinhaler therefore breaks up hardly any of these conglomerates which cannot penetrate into the finer airways. I am not surprised that only 8% of the total amount inhaled is excreted. In contrast, the