The study was performed while in receipt of a thrombosis research grant from Manchester Regional Hospital Board. We are also indebted to Manchester Family Planning Clinics, who sent the women for testing, and Dr. W. Lye, of Dista Products Ltd., for supplying the Dista Sequential preparation used in this study.

—We are, etc.,

L. POLLER.

JEAN M. THOMSON.

Withington Hospital,
Manchester 20.

REFERENCES

- 1 Committee on Safety of Drugs, *Adverse reactions series*, 1969, No. 8.
- 2 Thomson, J. M., and Poller, L., *British Medical Journal*, 1965, **2**, 270.
- 3 Poller, L., and Thomson, Jean M., *British Medical Journal*, 1966, **2**, 23.
- 4 Poller, L., Tabiowo, Anne, and Thomson, Jean M., *British Medical Journal*, 1968, **3**, 218.

Asparaginase

SIR,—In your leading article (24 May, p. 465) you state: "Asparaginase is now known also to affect normal cells, particularly those of the thymus and lymphoreticular system. Whether this effect, which has so far been seen only in animals, will prove an added hazard to therapy in man is not yet known."

I recently treated a patient with acute lymphoblastic leukaemia who developed a profound leucopenia following asparaginase treatment. The findings in life and the post-mortem examination strongly suggested that the normal cellular elements of the bone marrow, as well as leukaemic cells, had been depressed. A detailed report on this patient by Dr. M. J. Oehlers, Dr. W. Fetwadjeff, and myself will be published elsewhere.—I am, etc.,

H. J. WOODLIFF.

Department of Pathology,
Brompton Hospital,
London S.W.3.

Acute Infective Gastroenteritis

SIR,—I should like to comment on the letter by Dr. Mary J. Wilmers and others (31 May, p. 573) advocating the wholesale use of neomycin and other antibiotics in their gastroenteritis ward with the expressed purpose of preventing cross-infection with pathogenic *Escherichia coli*.

They give as their rationale the failure of careful barrier nursing techniques to contain cross-infection with these organisms. We have developed a system of barrier nursing at this hospital which has proved highly effective,^{1,2} but it must be emphasized that in order to operate the system efficiently it is essential to have properly constructed and equipped cubicles and a sufficiency of nurses for both day and night duty.

There are a number of possible disadvantages and dangers in the indiscriminate use of suppressive antibiotic treatment in babies. (1) It may engender in the minds of doctors and particularly nurses a false sense of security. Dr. Wilmers and her colleagues continue to get periodic bouts of cross-infection despite the approach they are recommending. (2) This practice encourages the emergence of resistant strains of *E. coli*. (3) Neomycin and antibiotics such as oxytetracycline may have undesirable and occasionally serious side-effects in small babies.

The question of whether or not to use antibiotics for babies seriously affected with gastroenteritis is a separate and debatable

matter. For the last four years or more we have seriously restricted the use of antibacterial preparations in our gastroenteritis unit, and have found rather to our surprise that very ill infants not given antibiotics generally respond promptly and well to accepted measures directed at restoring and maintaining electrolyte and fluid balance coupled with skilled nursing.

Needless to say, faced with an outbreak of hospital-acquired gastroenteritis in a paediatric unit or maternity department, it would be folly not to use an appropriate antibiotic in an attempt to stem the spread of infection.—I am, etc.,

TREVOR P. MANN.

Royal Alexandra Hospital for Sick Children,
Brighton 1, Sussex.

REFERENCES

- 1 Gibson, M., and Mann, T. P., *Nursing Times*, 1965, **61**, 1309.
- 2 Gibson, M., *Nursing Times*, 1965, **61**, 1334.
- 3 Cushing, A. H., *Pediatrics*, 1967, **40**, 656.

Contaminated Chloroform

SIR,—For at least 15 years we have used analytical grade chloroform for the extraction of the dithizone complex with lead when estimating this metal in biological fluids. During the last few weeks these estimations have given anomalous values which have appeared to be too high. Careful investigation ruled out contamination of the apparatus as a cause. The source of the error has been traced to certain batches of chloroform labelled as being suitable for use with dithizone and free from metallic impurities. Some batches of chloroform gave a red colour with dithizone which was equivalent to a positive error of 20 µg. of lead/100 ml. whole blood.

The impurity can be removed from the contaminated chloroform by washing with 10% sulphuric acid. The washings give an apparent lead content of 1.9 µg. of lead in 100 ml. of contaminated chloroform, in the presence of citrate/cyanide buffer. The metallic impurity is not lead, as by taking the same amount of wash liquor and omitting the cyanide buffer, the complex gives an optical density of 1.75 at 520 mµ in 0.5 cm. cells in 10 ml. of chloroform. It should be noted that if the contaminated chloroform is used to "delead" cyanide buffer solution then the cyanide becomes contaminated with the impurity which is not removed by dithizone.

Similar high optical densities can be obtained in the presence of thiosulphate, of iodide, and of citrate buffers. Thus, severe interference can be expected in the measurement of copper, cadmium, cobalt, and zinc when contaminated chloroform is used in the dithizone methods for these elements.

It is strongly recommended that laboratories which use chloroform for the extraction of dithizone metal complexes should test each batch of every consignment of chloroform before it is used in micro-analytical methods. It appears that the chloroform became contaminated during the filling process owing to a faulty filling line.—We are, etc.,

R. C. BROWNE.

J. STEEL.

R. W. ELLIS.

Nuffield Department of Industrial
Health,
University of Newcastle upon Tyne.

Relapsing Amoebic Colitis

SIR,—The report by Drs. S. R. Kanari and R. Knight (7 June, p. 613) will, I hope, stimulate awareness of the possibility of amoebic disease in patients who have lived or served abroad.

Army personnel have been particularly at risk in certain campaigns. In 1965 I diagnosed amoebic colitis in over a dozen soldiers who had been infected at the same source in Borneo, but there were many others, both British and Australian, who were at risk. Should any of these attend doctors, military or civil, with a complaint of piles, or recurrent diarrhoea, or rectal bleeding, it is to be hoped that amoebiasis will be considered. My patients were frequently referred with "piles." In the acute case I noticed that the abundant mucus always contained myriads of tiny bubbles; it was almost effervescent. I have not seen this described and it may not be of significance, but in that particular outbreak led me to repeated searching and examination of the specimens myself in the laboratory if a negative report was received from a technician.

As has been pointed out many times, the geographical history is the most important clue to the possibility of "tropical disease."—I am, etc.,

R. SCOTT.

Military Hospital,
Catterick Camp, Yorks.

Neonatal Serum Bilirubin Estimation

SIR,—Dr. T. Hargreaves (17 May, p. 448) indicates that false "low" results could occur using a diazo-coupling method during serial estimations of neonatal bilirubin levels and be reported as "normal" results, thus leading to complications arising over any proposed treatment by exchange transfusion.

Recently, while comparing two methods for the estimation of serum bilirubin levels in neonates, using a micro spectrophotometric method,¹ and a diazo-coupling method,² both methods were found to be in close agreement, the spectrophotometric method tending on average to be 1 mg.% higher than the comparable diazo results. However, in one case, a premature jaundiced infant, there was a marked discrepancy between results obtained using both methods, the diazo method producing consistently higher values than the spectrophotometric method over a period of several days (see Table). After careful rechecking of both methods, further inquiries resulted in the information that the baby was being treated with the antibiotic ampicillin (50 mg. twice daily). During a subsequent exchange transfusion samples of blood before and after transfusion were obtained and the serum separated, 2 ml. was acidified and extracted with ethyl acetate. The extract was evaporated to dryness and the residue subjected to two-dimensional chromatography, first in isopropyl alcohol, ammonia, water (80:10:10) overnight, and secondly in Anisole/acetic acid (70:30) for five hours. The chromatograms were dried, sprayed with 10% sodium carbonate, redried, and developed in diazotized *p*-nitro aniline to detect phenolic compounds. Both pre- and post-transfusion chromatograms developed an intensely staining orange/red spot which did not appear to correspond, position-wise, to any known phenolic acid and was presumed to be either ampicillin or a metabolite of ampicillin. On stopping treatment with ampicillin the marked discrepancy between the diazo and spectrophotometric values for the bilirubin levels which had occurred reverted to the expected level. Blood was also taken from full-term in-