SIR,—I was saddened to read the report of the proceedings of the Special Representatives Meeting on the Todd report (*Supplement*, 23 May, p. 139). The Representative Body has again shown lack of vision. In its consideration of the recommendations of Council, it copied Council in failing to consider the Royal Commission's proposals for postgraduate training in the light of undergraduate education as it will have evolved in, say, 20 years' time. This mistake has been made in almost every discussion I have heard on the subject in the past year.

As I understood the Todd report, the Royal Commission on Medical Education considered that, in view of the changing basis of medicine from the artistic to the more scientific, the needs of the potential doctor will change. These needs must be met as closely, yet as efficiently, as possible: by careful long-term planning and perhaps by radical changes in the type of initial training offered—that is, the undergraduate would be given a basic education in medical science (in its widest sense). In many medical schools plans for such changes are almost off the council table. (There is no suggestion that this will not include "meeting patients" at an early stage, as feared by your correspondent Dr. J. S. Ross (23 May, p. 479)). As if in a game of Chinese whippers, the meanings of many words and phrases coined in the Todd report have changed and are giving rise to confusion. It is understandable that, faced with reams of reports and repetitive debate, the representatives have lost track of the original concept. That a speaker as well informed as Mr. J. S. Elkington (*Supplement*, 23 May, p. 141) should speak as eloquently as he does on a completely false premise is symptomatic—and tragic.

I would appeal to those involved in debate and negotiation to remember that they are probably blessed with above average initiative. On the other hand, some of the well-prepared but uninitiated medical graduates of the future may flounder and disappear without some (flexible) guidance to further training in the continuing rat-race. This would be utterly detrimental to any health service this country may hope to sustain.—I am, etc.,

ALISON P. HILL.

Guy's Hospital Students' Club,
Guy's Hospital,
London S.E.1.

Mobile Pressure Chamber

SIR,—Mr. J. N. Norman and his surgical colleagues are to be congratulated on the design, safety, and simplicity of their one-man hyperbaric pressure chamber (9 May, p. 333). It is unfortunate that the indications for the use of the chamber are not described. They state that "... the benefit of using hyperbaric oxygen in carbon monoxide poisoning is twofold, the rapid dissociation of carboxyhaemoglobin being secondary to the immediate correction of tissue hypoxia." Their patients with a carboxyhaemoglobin level on admission of the order of 5%—a level achieved by smoking one cigarette—clearly cannot benefit from the former effect. The tissue hypoxia which exists after the carboxyhaemoglobin has fallen as low as 5% is a phenomenon second-

dary to capillary damage and oedema. The most important clinical manifestation of this is cerebral oedema. This responds in minutes to the intravenous infusion of hypertonic mannitol.¹ Hyperbaric oxygen would seem to be unnecessary in this circumstance.

The major benefit of hyperbaric oxygen is the correction of the primary anoxia and the rapid dissociation of the high levels of carboxyhaemoglobin. This can only be achieved by the early exposure of the patient to hyperbaric oxygen, and it is in this instance that the mobile unit described would be of great value.

Caution is indicated in regarding "most of our cases" as accidentally poisoned. Despite age, impaired sense of smell and vision, together with confusion in operating modern gas taps, Kessel² has stated that the poisoning is unlikely to be accidental and is in fact intentional self-poisoning.—I am, etc.,

HENRY MATTHEW.

Regional Poisoning Treatment Centre,
Royal Infirmary,
Edinburgh.

REFERENCES

- Matthew, H., and Lawson, A. A. H., *Treatment of Common Acute Poisonings* 2nd edn., 1970, E. & S. Livingstone, Edinburgh.
- Kessel, N., *British Medical Journal*, 1965, 2, 1265 and 1336.

Underpayment and Overwork

SIR,—I am in complete agreement with the views expressed by Drs. R. D. and Mary Catterall (25 April, p. 234). There has been no effective rectification of the fundamental problems which exist. Should this not be done fairly soon, continued emigration (particularly of those specialists who are universally scarce) and a decline in the standards in practice in British medicine is inevitable.

Less than one year ago as a senior registrar in a large British teaching hospital I was thoroughly dissatisfied with conditions of the hospital junior doctor and had serious intentions of emigrating. At present I am the holder of a travelling scholarship which has allowed me to spend time in France, Denmark, and the U.S.A. This opportunity has allowed me to re-evaluate Britain and British medicine in the light of other cultures and systems. It has been pleasing to observe that the best of British medicine and medical training compares favourably with that in Europe, Scandinavia, and North America. But during this time I have been nowhere (or met anyone who has) where the conditions and salaries of hospital junior doctors are worse than they are in Britain.

It now appears to me that all the multitude of complaints and grievances are merely different ways of expressing the fact that salaries are too low.

To say that the trainee specialist would not have all his present problems if only he or she were to delay marriage may well be true, but in present-day society is an anachronism and neither fair comment nor a feasible solution. The hospital junior doctor in his late twenties does not have to look far to realize that society demands more and gives less to him than to most other groups in present-day British society. Most of us would agree with Socrates that

"Human nature cannot know the mystery of an art without experience." We grumble about the long hours we have to work really because this allows us insufficient time to do those things which enable other people to exist reasonably well on low salaries. The young married man working for his examinations cannot afford the time to paint the house or fix his car, neither can he afford to pay someone to do it for him. I complain about the numerous times either my husband (also a doctor) or I arrive home at 10 p.m. instead of 6 p.m., because our budget does not allow us to get into the car and drive to the nearest restaurant it demands that I must now cook a meal. Because it is easier and more socially acceptable, in the days of the 38-hour week, these grievances tend to be expressed as a desire for shorter hours rather than for better salaries.

No one can deny that it is possible to obtain a postgraduate education and experience in Britain second to none in the world, but the fact that books and courses are so expensive puts the best out of reach of the large percentage of British graduates, most of whom are compelled to follow the longer and less efficient path of self-education.

It seems to me that it is unreasonable to expect that in our materialistic society young men and women should continue to tolerate the situation indefinitely. If they fail to change the status quo they will take one of two courses: they will choose general practice as a career (for exactly the same reasons as in North America radiology now attracts a higher percentage of the top medical graduates than ever before), or they will emigrate.

I personally, only because I am fortunate enough to belong to a household where there are two professional salaries, have decided that I do not wish to exchange the things I value in British medicine and European culture and way of life for a larger salary. My hope in expressing this viewpoint is that it might cause those members of the profession with real influence to re-evaluate the situation of the hospital junior doctor in relation to the present rather than the past as they knew it.—I am, etc.,

MARGARET BARK.

University Hospital,
Ann Arbor, Michigan,
U.S.A.

Advice and Contraception

SIR,—“Crisis in the F.P.A.” is described in your news pages (9 May, p. 372). The dissension is between the authorities who wish to serve the greatest number of people with contraceptive advice with the minimum of necessary consultation and the doctors who see the need for “adequate” consultations.

Surely the evidence from all over the world goes to show that the doctors are right. The majority of those most desperately concerned with problems of mating and parenthood need to know not more about how, but more about why to undertake fertility control.

In Britain alone the number of family planning clinics has just about doubled in the past six years. But has there been any reduction in the numbers of unwanted babies, illegitimate pregnancies, incidence of