and child psychiatrist are also needed for the regular assessment of the child's developmental progress and to help with the emotional problems which arise commonly in children requiring a special and rather unpalatable diet for many years.

The greatest danger in treatment arises from brain damage, and even death, resulting from too severe restriction of dietary phenylalanine, and we have had children referred to us in whom it seemed likely that severe mental retardation was the result of such dietary mismanagement. A regular prompt laboratory service for phenylalanine estimations is thus essential.

We, therefore, suggest that centralization of the processing of the screening procedure, confirmation of the diagnosis, and management of the established case should all take place in one of a small number of special centres. Such centralization of management does not necessarily involve frequent visits to the centre after the initial inpatient stay. An appropriate sample of blood or plasma for the estimation of phenylalanine can be sent by post. Since the dietitian will be known already to the mother, any dietary change required in the light of the result can be made without delay by 'phone or by post. We have controlled the management of many children in this way and, in general, have encountered excellent co-operation from the local paediatrician and general practitioner.

Phenylketonuria is rare; only about 90 patients are born in the United Kingdom each year. If our management is to be adequate and if, in addition, research is to be carried out to improve methods of diagnosis and management, centralization is essential.

For complicated surgical procedures, such as open heart surgery, the profession is beginning to realize that centralization leads to better results and is more economical. Similarly, we should recognize that there are certain rare medical disorders whose management is complicated and that patients suffering from them benefit from being referred to a special centre. Phenylketonuria is one of them.—We are, etc.,

BARBARA E. CLAYTON.

O. H. WOLFF.

Hospital for Sick Children,
London W.C.1.

REFERENCES

- 1 Medical Research Council Working Party on Phenylketonuria, *British Medical Journal*, 1968, 4, 7.
- 2 Department of Health and Social Security, *Screening for early detection of phenylketonuria*. Circular H.M. (69) 72. London, H.M.S.O., 1969.

Liquorice and Addison's Disease

SIR,—The report by Dr. D. I. M. Siegler concerning a patient suffering from autoimmune Addison's disease with hypercalcaemia (30 May, p. 522) is incomplete in that it omits to mention that the patient had a liking not only for salt but also for liquorice. The sodium-retaining properties of this sweet protected her from clinical adrenal insufficiency in the heat of Aden, where she lived as a soldier's wife, and also masked many of the physical signs of this disease—for example, pigmentation. This is

the second patient I have encountered in which the ingestion of large amounts of liquorice has enabled a patient with Addison's disease to escape from the clinical results of adrenal failure for a long period of time. One can speculate as to whether her penchant for liquorice, as well as her appetite for salt, arose from some subconscious intuition as to the therapeutic value of these substances in her disease.

Addison's disease is indeed a rare cause of hypercalcaemia. The doctors in the metabolic department of this hospital, who have a considerable experience of hypercalcaemia, had not previously encountered Addison's disease as a cause. Indeed the real cause of the hypercalcaemia of this patient would have escaped recognition had she not been shown at the hospital's weekly staff meeting.—I am, etc.,

E. J. ROSS.

Medical Unit,
University College Hospital
Medical School,
London W.C.1.

Susceptibility to Aspirin Bleeding

SIR,—Your leading article (23 May, p. 436) on susceptibility to aspirin bleeding emphasizes a number of important points, notably the problem of defining the causes of susceptibility to aspirin-induced bleeding. However, the initial claim that 25% of cases of haematemesis and melaena in the London area are caused by aspirin is unjustifiable. In the source paper¹ there were 56% of recent aspirin-takers among those with acute bleeding compared with 32% in controls. The excess of 24% could represent the effect of aspirin if the control and bleeding series were comparable, but here there is cause for doubt. The control series used by Valman and his colleagues would appear to be identical with that of Parry and Woods,² and the latter authors emphasize that the proportions of aspirin-takers in their control series varied from 25 to 50% according to the diagnostic grouping used. If 50% were appropriate, and there are no means of judging, then the excess of aspirin-takers in the test group would fall to 6%.

The problems of ensuring comparability between test and control groups in investigating the role of aspirin in inducing acute gastrointestinal bleeding have been commented upon by others,³⁻⁵ and it seems clear that the complete comparability has yet to be achieved. The dyspeptic patient used as a control may have given up aspirin because it made his abdominal symptoms worse, while the patient with a stroke or coronary thrombosis may have forgotten the, to him, insignificant event of taking aspirin shortly before admission. On the other hand, the patient with haematemesis or melaena may have taken aspirin because he mistook for a 'flu-like illness the faintness initially induced by the bleeding.

The difficulties could be solved if patients with so-called aspirin-induced bleeding could be proved to be consistently susceptible but in fact, as you say, when they have been re-challenged with aspirin they appear to be no more liable to occult bleeding than control subjects.³ This anomaly could be explained if a combination of aspirin and another factor were needed to induce acute bleeding. Alcohol would seem to be an obvious accessory factor, but though it ac-

centuates occult bleeding there is no clear evidence that it alters liability to acute haemorrhage.

The general epidemiological evidence suggests that aspirin may be a causal factor in inducing bleeding but it is not unequivocal. In a paper recently published by you, Dr. K. F. R. Schiller and his colleagues⁶ found that during three successive quinquennia the rate of admission to the Radcliffe Infirmary for haematemesis and melaena remained constant. During that time the proportion of individuals recorded as taking drugs, predominantly aspirin, beforehand was found to have quadrupled. This disparity might only reflect the recent interest in aspirin as a cause of bleeding; however, it is also noteworthy that the proportions of patients with gastric and duodenal ulcer did not fall as might have been expected if aspirin-induced acute ulceration had become an increasing problem.—I am, etc.,

M. J. S. LANGMAN.

Department of Medicine,
University of Nottingham.

REFERENCES

- 1 Valman, H. B., Parry, D. J., and Coghill, N. F., *British Medical Journal*, 1968, 4, 661.
- 2 Parry, D. J. and Wood, P. H. N., *Gut*, 1967, 8, 301.
- 3 Allibone, A., and Flint F. J., *Lancet*, 1958, 2, 1121.
- 4 Davis, J. S., *Lancet*, 1958, 2, 1121.
- 5 Salter, R. H., *American Journal of Digestive Diseases*, 1968, 13, 38.
- 6 Schiller, K. F. R., Truelove, S. C., and Williams, D. G., *British Medical Journal*, 1970, 2, 7.

Thrombopoietin

SIR,—Thrombopoietin being a "bee in my bonnet" since 1957, I read with great interest the preliminary communication by Dr. D. G. Penington (7 March, p. 606). He states that in the experiments of my colleagues and myself increases in the platelet counts of animals receiving different plasma samples are "relatively small, of the order of 20 to 25% over the resting platelet count." In fact, our papers show that only increases of 30% or more over the resting counts were considered as "positive." In addition, we demonstrated that elevation of the circulating platelet count in several conditions is preceded, in man as in animals, by the appearance of higher levels of this apparently specific factor. This plasma factor cannot be dialysed, runs with beta-globulins, and is easily destroyed by trypsin.¹ It is likely that other plasma factors function in the later stages of thrombocytopoiesis. Availability of thrombopoietin alone does not, therefore, guarantee the undisturbed maturation of cells of the megakaryocyte series and, conversely, if a certain plasma factor is able to induce thrombocytosis, this factor may or may not be thrombopoietin as such.

We never stated that the plasma thrombopoietic activity is elevated in "essential" thrombocythaemia. On the contrary, we stated that seven out of seven apparently "essential" cases gave negative responses in our mouse test.² The simplest way to demonstrate elevated thrombopoietin levels is to test plasma samples from severely thrombocytopenic patients. More than 100 cases of thrombocytopenia were tested in our laboratories, and the fact that some samples of thrombocytopenic plasma exhibit elevated thrombopoietic activity has already been