carcinoma patients the serum calcium was raised to around the 16 mg/100 ml level and that the classical symptoms of hypercalcaemia were present; there were no skeletal metastases.

It would appear that the low incidence of the hypercalcaemic syndrome in Locks's series and its absence in the Jersey study would tend to support the comments of Drs. von Wichert and Mitchell-Heggs.—I am, etc.,

JOHN CRAGG

General Hospital,
Jersey, C.I

¹ Locks, M. O., *Journal-Lancet*, 1962, 82, 165.

New Look at Tennis Elbow

SIR,—A study of the relationship between tennis equipment and tennis elbow, on the rare occasions in which it actually affects tennis players, may shed some light on the aetiology and management of the condition in general.

The pathogenesis of the condition is not well understood. Most of the extensor muscles of the wrist arise from ligaments, fascia, and intermuscular septa which are relatively mobile. This may make it difficult to co-ordinate the contraction of individual fibres when the hand is gripping and the forearm rotating as in tennis. This lack of easy co-ordination may predispose to minor repeated unrecognized injury near the origins of the muscles, and may also make healing more difficult as the mobility of the origins may be affected by adhesions.

In many of the tennis players affected by this condition the equipment used is faulty. Either the racket lacks resilience and flexibility, or the gut is too "boardy," and the arm is jarred causing pain. The handle may be too big or too small, putting excessive strain on the forearm muscles when they are rotating. A top-heavy racket has a similar effect. It is difficult to attribute a cure to any special measure in a condition like tennis elbow which can cure itself, but in many cases when the "faulty" racket is replaced the symptoms are relieved more quickly than by other means. The most dramatic instance is that of an experienced player with this condition who over two years had tried everything except operation (including indigenous treatments) without relief. He changed to another racket of the same weight and handle size but with a better weight distribution and a shock absorbing string suspension system; his symptoms disappeared in two days and have not returned for nine months. Relating these observations to the known theories about the aetiology of tennis elbow it is our impression that the use of faulty equipment can in tennis players lead to excessively inco-ordinated muscle action and damage which produces tennis elbow. Correction of the tools can relieve the condition.

It is possible that in non-tennis players (who form the majority of the sufferers from this condition) there are similar "faults" in equipment—for example, badly balanced shears—or, of course, bad technique in the use of faultless equipment. We would suggest that early consideration should be given to modifying equipment or its use, in addition

to the traditional methods of management in all cases of tennis elbow.—We are, etc.,

S. CHAWLA
A. P. R. ALUWIHARE

Diana & Co. Ltd.,
Colombo 1, Ceylon

Waiting Room Problems Solved

SIR,—One of the intractable problems facing hospital administrators today is the out-patient queue. Everybody talks about it, many complain about it, but here at last is someone who is doing something about it.

On a recent busy Monday morning here an elderly Indian gentleman was causing great delight to a large crowd of suffering humanity waiting to see a doctor. For a modest fee of sixpence, he would swallow the most remarkable objects—his recent meal had consisted of iron nails, broken glass and pieces of old torch batteries. He is well known in Fiji as an entertainer wherever crowds gather, but this was apparently the first occasion that he had chosen the clinic waiting room for his stage. He was persuaded to submit to radiological examination, and an abdominal x-ray is reproduced without comment.

In spite of his austere diet, our visitor clearly enjoys the best of health and temper, so there is no indication for medical interference.—I am, etc.,

G. HOLMES

Lautoka Hospital,
Lautoka, Fiji

Termination of Life

SIR,—Your remarks (23 January, p. 187) on this subject are tinged with current prejudice. You talk of legal euthanasia "tragically undermining the patient's endurance," thus revealing the commonly held philosophy that illness, however incurable or protracted, is something society has ordained to be suffered to the end. To ask for and obtain release can neither be antisocial nor immoral, and the burden of denying it must lie heavily on the conscience of an ultraconservative profession.

Doctors have daily experience of appalling torture suffered by their patients, only very inadequately relieved by the resources of medicine. To talk of "killing patients" is a

confusion of semantics. But the 1961 Suicide Act should be amended to allow a doctor to help his patient yield up his life if he so desires, accepting death openly and fearlessly as a gift from a merciful Providence. Our present attitude is irrational and primitive. It mirrors the ultra materialism of contemporary society.—I am, etc.,

S. L. HENDERSON SMITH

Huddersfield, Yorks

N.H.S. Superannuation

SIR,—I am writing to say how much I support the Association in its efforts to obtain an improved N.H.S. superannuation scheme and how much I welcomed your leading article (20 March, p. 622) on this subject.

There is, however, one important point which you appear to have overlooked as there is a further way in which these N.H.S. pensions compare badly with other government pensions. I am advised that N.H.S. pensions are paid less British income tax even if the recipient becomes resident abroad and no longer liable for British tax. In this way N.H.S. pensions seem to be separately discriminated against, as this is not the case with Civil Service and Army pensions.

I think it is particularly unfortunate that this discrimination is made because many doctors, especially hospital ones, might decide to retire at the age of 60 in order to do service in one of the underdeveloped countries. This would be one way in which these countries could obtain the services of very experienced specialists for a number of years, but this special discrimination must seriously discourage this.—I am, etc.,

REGINALD J. CARR

Royal Infirmary,
Bradford, Yorks

Hospital Practitioner Grade

SIR,—It was reported in the *Supplement* (6 March, p. 64) that it has been proposed that the General Medical Services Committee should be responsible for making submissions to the Review Body and for negotiations with the Department of Health and Social Security on the above grade. This is a welcome development. Occupational health for hospital staff is a field in which this new grade may be of particular value. At present, doctors wishing to specialize in occupational health within the hospital service are graded clinical assistants.

In new and large district hospitals the responsibilities and training necessary to initiate and run an occupational health department are considerable. A doctor of sufficient experience is unlikely to pursue a career in the clinical assistant grade. Unless realistic salaries and gradings are applied to this new specialty within the hospital service, doctors of the right calibre and training will not be available. We are now witnessing new hospitals inadequately staffed, with wards and departments failing to open. If occupational health is to make a contribution to improving this situation, doctors with specialized experience must be recruited.

Acceptance of the hospital practitioner grade may be the means to achieve this.—I am, etc.,

J. A. LUNN

Northwick Park Hospital,
Harrow, Middx