undermine the whole basis of the doctor-patient relationship and shake the confidence of the public, especially the adolescent girl who so needs our help.—I am, etc.,

ERIC TOWNSEND

Camborne, Cornwall

Pathogenesis of Myasthenia Gravis

SIR,—Your most interesting leading article on myasthenia gravis (3 April, p. 1) prompts me to refer to earlier work indicating the presence of a hormone in the thymus influencing neuromuscular transmission. Torda and Wolff¹ showed plasma in myasthenia gravis inhibited acetylcholine synthesis, and in 1944 we² published a paper showing that normal thymus increases acetylcholine synthesis and thymus tumour in myasthenia gravis inhibits synthesis of acetylcholine. We concluded that the replacement of stores of acetylcholine in the neuromuscular junction was the deficiency in myasthenia gravis, and you indicate the same with the production of lower quanta of acetylcholine at the motor endplate.³ I also pointed out⁴ this correspondence in muscle weakness in thyrotoxicosis with "lymphorrhages" in muscle and myasthenia gravis, and suggested these lymphoid collections produced a similar substance.—I am, etc.,

E. R. TRETHEWIE

University of Melbourne,
Victoria, Australia

¹ Torda, C., and Wolff, H. G., *Science*, 1943, **98**, 224.

² Trethewie, E. R., and Wright, R. D., *Australian and New Zealand Journal of Surgery*, 1944, **13**, 244.

³ Thesleff, S., *Annals of the New York Academy of Sciences*, 1966, **135**, 195.

⁴ Trethewie, E. R., *Medical Journal of Australia*, 1956, **1**, 13.

Jejunal pH and Folic Acid

SIR,—In a recent report Dr. Ann Benn and others (16 January, p. 148) showed that the jejunal pH in epileptics on longterm anticonvulsant therapy was more alkaline than in controls, which might cause malabsorption of folic acid.

This explanation of anticonvulsant megaloblastic anaemia is fully in agreement with my results of in vivo perfusion experiments in rats.¹

These experiments showed that the absorption of folic acid depends on the intraluminal pH. Introduction into the jejunum of barbiturates in unbuffered solutions caused a rise of pH in the intestinal contents and consequently the absorption of folic acid was reduced by $\frac{1}{2}$ – $\frac{3}{4}$ of the total. In buffered solutions barbiturates caused neither alterations of pH nor decreased absorption of folic acid.

In one patient on treatment with phenytoin 150 mg twice daily for epilepsy the absorption of folic acid was reduced by approximately half by phenytoin (Fig.). As betainehydrochloride, which in the stomach splits off hydrochloride, had an antagonistic effect on phenytoin it is assumed that in the gastrointestinal tract phenytoin causes a disturbance of the pH regulation, resulting in malabsorption of folic acid. I have also found malabsorption of folic acid in five out of six patients with anticonvulsant megaloblastic anaemia.²

Dr. Ann Benn and her colleagues claim that increasing alkalinity above pH 5.0 would cause a fall in the concentration of folic acid in the neutral form, in which condition folic acid crosses the intestinal wall. In my experiments it was found that the maximum of absorption for folic acid was about pH 6.0. Higher as well as lower pH decreased the absorption rate of folic acid. However, 100 mg barbitone (or phenytoin) taken by mouth might cause a concentration of barbitone of approximately 5 mmol/l, in the proximal jejunum. If P_{CO_2} is 40 mm Hg, intraluminal pH would be 6.8, which is well away from the maximum of absorption.—I am, etc.,

LARS ELSBORG

University Department of Medicine II,
Aarhus, Denmark

¹ Elsborg, L., *Abstract of XIII International Congress of Haematology*, 1970, p. 378

² Elsborg, L., in press.

Screened Fibrinogen for Prophylaxis of Venous Thrombosis

SIR,—Your recent leading article (6 February, p. 305) on the "Prophylaxis of Venous Thrombosis" stresses the usefulness of the ¹²⁵I labelled fibrinogen technique for the detection of thrombosis in the leg. Our own studies using this technique confirm its usefulness and that the findings correlate very well with those obtained by venography. The real value of the technique lies in the fact that it allows the investigator to observe the evolution of venous thrombosis in the legs of patients in whom there may be no clinical evidence of the complication, and to do so without disturbing the patient. Hence, the technique may be used to screen relatively large numbers of patients and as a tool for monitoring the effect of prophylactic measures it is unrivalled. Venography is expensive, uncomfortable, and not without risk and is therefore only warranted when there is no other evidence of thrombosis.

The fibrinogen used is from human donors and it is known that the blood fraction containing fibrinogen may transmit the infecting agent of serum hepatitis. This is a risk that has been known to be associated with blood transfusion and the giving of most blood

products for many years. Such risks may be acceptable in therapeutic and diagnostic situations but may not be in clinical trials. The recent advent of techniques which permit the recognition of the hepatitis associated antigen has made it possible to screen selected donors and donations in depth and to reduce considerably the risk of transmitting this infection. This being so, we have reluctantly decided that it would be unethical for us to continue to use fibrinogen for a clinical trial unless we could be sure that it came from donors and donations that had been adequately screened. To date, we have been unable to find any source of such material and have felt obliged to discontinue our clinical trials.

We feel that there is an urgent need to provide screened fibrinogen for this very valuable test and hope that suitable material will be available in the very near future.—We are, etc.,

N. C. ALLAN
R. A. MILNE
C. V. RUCKLEY
J. K. WOOD

Department of Haematology,
Western General Hospital,
Edinburgh

Nocturia during Steroid Therapy

SIR,—Three cases of nocturia associated with hypertrophic osteoarthropathy were recorded in the correspondence columns (12 December, p. 686). In Case 3 the patient is now well and symptom free. Serum sodium level 138 mEq/l. Serum potassium level 3.8 mEq/l. In conditions similar to those at preoperative assessment the urinary output of water, sodium, and potassium five months after lobectomy for carcinoma were:

Average over 5 days	7 am to 3 pm	3 pm to 11 pm	11 pm to 7 am
Urine Vol. (ml)	790 (340)	830 (520)	513 (900)
Total Sodium (mEq)	53 (16.5)	55 (35)	36 (65)
Total Potassium (mEq)	17 (10)	16 (15)	10.5 (17)

Figures in parentheses are the preoperative values.—I am, etc.,

LESLIE DOYLE

Baguley Hospital,
Manchester

Ulcerative Colitis

SIR,—The widely authoritative account of ulcerative colitis (20 March, p. 649) would be just that more complete if the very small section on psychological treatment had been amplified and a good reason for this had been given.

The reason is that, investigated in the way their cases were, the findings of the physicians who wrote on ulcerative colitis between 1930 and 1960¹ are still valid. Thus, in 44 cases seen in series in the Radcliffe Infirmary, Oxford, between 1962 and 1964 all were found to have temperamental or character abnormalities present before the onset of physical symptoms, as evidenced by careful history taking. Also the diarrhoea or

bleeding of well-defined colitis was found to have been credibly precipitated by emotion in 17 of these patients (39%). Psychotherapeutic sessions were few; no deep or long treatments were attempted. But short-term psychotherapy directed to both the character problem and the precipitating emotional factors (most commonly dependence, ambivalence, and loss in a sensitive "immature" personality) did appear to assist recovery or ward off relapse. Accurate assessment was impossible since nearly every patient was also on active drug treatment.

It has been suggested that the key word here is "sensitivity."^{2,3} The autoimmune interpretations of the present-day should not be held to be in conflict with the emphasis upon emotional sensitivity so regularly reported by observant physicians of the past. Does not emotional sensitivity frequently run with protein sensitivity? But I think it wrong to regard the main disturbance of emotional life as being resultant upon the colitis. More thorough studies place it antecedent to the physical symptoms, and I think that adequate techniques of investigation—including the psychoanalytical—will continue to place it there.—I am, etc.,

RANYARD WEST

Shelton Hospital,
Shrewsbury

¹ Daniels, G. E., *Psychosomatic Medicine*, 1940, 2, 276.

² West, R., *Advances in Psychosomatic Medicine*, 1959, 1, 270.

³ West, R., *Nursing Mirror and Midwives Journal*, 1968, 127, No. 5, p. 19.

Estimation of Cholesterol

SIR,—It is a great pity that in his otherwise excellent article (24 April, p. 187) Dr. D. S. Fredrickson omitted to include specific reference to the method of cholesterol estimation. This, to some extent, makes nonsense of such a statement as "... men with cholesterol above the 5% limit (about 290 mg/100 ml) may account for about a quarter of the myocardial infarcts." The use of blood cholesterol levels also occurs in other parts of the article.

It has been my experience that many clinicians do not seem to realize that the normal limits of cholesterol vary not only with age, etc., but also with the method of estimation employed. I am sure that this has caused unnecessary worry to many patients.

I received in this morning's post a copy of the publication issued by the Department of Health and Social Security on *Using the Laboratory*,¹ for the use of medical practitioners. I was delighted to find that this booklet points out the variation in levels of cholesterol, and other substances, with the analytical method used.—I am, etc.,

A. L. LATNER

Department of Clinical Biochemistry,
University of Newcastle upon Tyne

¹ Department of Health and Social Security, *Using the Laboratory*, London, 1971.

E.E.G. and Anticonvulsants

SIR,—The abnormal E.E.G. reported by Dr. W. I. M. Dow (24 April, p. 207) in a patient receiving a grossly excessive amount of anticonvulsants (primidone 1.5 g, pheneturide

800 mg, methylphenobarbitone 120 mg, and sulthiame 400 mg/day) and its subsequent improvement upon reduction of the dosage of primidone are to be expected.

The reduction of the drug level (hence its toxicity) and the changing state of awareness are the correlates of the E.E.G. change. The effect (if any) of folate deficiency and the effect (if any) of folate supplement on the electroencephalogram have not been demonstrated.—We are, etc.,

COLLEEN DAFBY
DAVID FUNG

Department of Electroencephalography,
Runwell Hospital,
Near Wickford,
Essex

Migrating to Saskatchewan?

SIR,—The physicians of Saskatchewan have welcomed many colleagues from Great Britain who have come to this province to practise. Each one that has come has made his mark in one way or another. I am writing to thank you for assisting us in this way.

I am also writing to advise your readers that on occasion they may be hired to come here on the basis of an unreasonable contract. It is usual for one of your people to come here and work as an assistant on salary for one year or so and then become a partner in a fee-for-service practice. Others enter solo fee-for-service practice on arrival.

Those entering a solo practice need no contract beyond one which would bind them to paying for any debts incurred from the move. They may, of course, be purchasing equipment or a house on a loan basis. Those entering group practices, which are common here, will want to know how receipts and expenses are shared and what happens on the death of a partner or on dissolution of the partnership. Beyond that, there is not much really necessary in a contract.

I would like to suggest to anyone contemplating migration to Saskatchewan and who is confronted with an odd contract that he or she contact me. I will be pleased to offer whatever advice I can.—I am, etc.,

E. H. BAERGEN
Executive Secretary,
Saskatchewan Medical Association

211 Fourth Avenue South,
Saskatoon, Sask.,
Canada

Medical Postgraduate Study Tour

SIR,—I would like to draw your attention to the postgraduate study tour in the U.S.A. which takes place from 25 September to 4 October inclusive, with an optional extension to Toronto immediately afterwards. The return flight is from New York on Friday 15 October, and those participants not going on the extended tour to Toronto will have approximately ten days for privately arranged sight-seeing or other visits in America.

Bristol Local Medical Committee is organizing this tour to give doctors an opportunity of gaining an insight into American medicine in a short time. General practitioner participants will have the advantage of qualifying for the postgraduate training allowance and two years' credits for seniority payments.

The party is limited to 24 (including wives). Any doctors who are interested are invited to communicate with the Secretary,

Bristol Local Medical Committee, 11 Westmorland House, Durham Park, Bristol BS6 6XH, telephone number 0272-52762.

Any doctors with limited time who are prepared to make their own way to and from New York may apply to join the tour so long as places are booked in advance through the organizers.—I am, etc.,

JOHN T. ALDREN
Chairman,
Bristol Local Medical Committee

Bristol

Family Doctor Booklets

SIR,—I write to support Dr. Ian Oswald's warning (17 April, p. 171).

My concern is based upon my recent experience of providing information for the press and television from a maternity department. I have nothing but praise for the numerous reporters and cameramen with whom my colleagues and I had to deal. They were discreet, considerate, patient, intelligent, and accurate. Furthermore, a five-minute television film was a model of pertinent editing. On the other hand the newspapers which used our "story"—and these included all but one of the national dailies—did so with some degree of sensationalism, varying across a spectrum from a hint to a splash. Finally, a specific instruction about the description of a fetal monitoring instrument was ignored in three instances and the expressions "birth machine," "metal midwife," and "mechanical midwife" were used. Unsolicited apologies were received from the reporters concerned, who regretted that they had been over-ruled by subeditors.

I feel that my experience is similar to that of Dr. Oswald in that it represents a use of clinical material in a manner not authorized by its originator. I would, finally, comment that TV appears to be making a better job than the press of reaching the less informed sections of the community, at any rate in documentary programmes, without resorting to sensationalism or disregard of professional anonymity, when this is requested.—I am, etc.,

W. KEITH SUTTON

Rochford, Essex

Rickettsial Antibodies in Multiple Sclerosis

SIR,—I should like to make certain comments on the article "Rickettsial Antibodies in Multiple Sclerosis" by Professor E. J. Field and Miss Mavis Chambers (3 January 1970, p. 30). It is a matter of elementary scientific fairness that any worker wishing to test a theory should mention the initial text expounding this theory and should strictly apply the original research methods: the authors have not done this for my work.

(1) In 1960, in three communications to the Paris Academy of Sciences¹⁻³ I demonstrated a rickettsial aetiology of multiple sclerosis, underlining the usual "weak positivity" of sero-diagnoses—and not "their high levels"; I specified that my sole guide was epidemiology—and not a theoretical "antigen-antibody" concept, as the authors assert, thus distorting the genesis of my work and masking its epidemiological importance.

(2) I recommended "reactivation" in all cases of "negative sero-diagnosis"—and not "weakly positive sero-diagnosis," as the authors write; they did not understand that,