amantadine did not cause alpha-receptor blockade within the dermal vessels. Intradermal injections of isoprenaline 0.1 ml of 1 mg/ml caused accentuation of the vascular pattern; however, the response to isoprenaline showed marked variability, and in weaker dilutions a reddening of the network pattern of skin vessels and a pallor of the surrounding skin occurred.

Comment

Livedo reticularis is a common finding in both control subjects and untreated patients with Parkinson's disease. The blood supply of the skin is through arterioles which each supply a cone-shaped area with a base from 1 to 4 cm across (Unna, 1896). Where the bases anastomose there is some dilatation of the venules and capillaries with relative stasis giving a cyanotic discoloration. Livedo is accentuated during amantadine treatment, and the diffuse continuous pattern suggests a generalized vascular change of a physiological nature due to a widespread dilatation of the dermal vessels. (A patchy distribution is seen in livedo occurring as a result of underlying arteritic lesions.)

In some patients livedo was accompanied by ankle oedema. This ankle oedema was not associated with any increase in total body water or salt retention (Parkes *et al.*, 1971) and was

perhaps due to increased vascular permeability in the cutaneous blood vessels. However, oedema does not commonly accompany either physiological or pathological livedo, though in the latter definite changes in the blood vessels occur.

Amantadine will cause depletion of catecholamine stores in peripheral nerve terminals (Grelak *et al.*, 1970). In normal dosage it has little or no direct action on smooth muscle (Vernier *et al.*, 1969). Thus livedo produced by amantadine treatment may be caused by an effect of this drug on physiologically active substances in nerve terminals of the sympathetic nervous system, though this possibility requires to be further explored. The absence of any evidence of drug-induced systemic disease during amantadine treatment is in keeping with a physiological rather than a pathological cause of livedo.

References

- Grelak, R. P., Clark, R., Stump, J. M., and Vernier, V. G. (1970). *Science*, **169**, 203.
 Parkes, J. D. *et al.* (1971). *Lancet* In press.
 Shealy, C. N., Weeth, J. B., and Mercier, D. (1970). *Journal of the American Medical Association*, **212**, 1522.
 Unna, P. G. (1896). In *Histopathology of Diseases of the Skin*, translated by N. Walker. Edinburgh, Clay.
 Vernier, V. G., *et al.* (1969). *Toxicity and Applied Pharmacology*, **15**, 642.

MEDICAL MEMORANDA

Diagnosis of Haemorrhagic Illness

A. DAVID GINSBURG, HERMAN A. GODWIN,
RICHARD H. ASTER

British Medical Journal, 1971, 2, 628-629

Disseminated intravascular coagulation is being diagnosed with increasing frequency in a variety of clinical circumstances. The spectrum of its manifestations is broad, ranging from a rapidly fatal haemorrhagic illness to situations in which the condition is detectable only by laboratory testing. The following case report illustrates difficulties that may be encountered in differentiating this disorder from idiopathic thrombocytopenic purpura. It also emphasizes the dramatic response which may be obtained with heparin therapy, the need for gauging the dose correctly, and the hazards that may be associated with the use of this drug.

Case Report

An 85-year-old woman was admitted to the Boston City Hospital because of rectal bleeding. A full history and physical examination showed the only other significant findings to be the presence of ecchymoses and petechiae over the legs and soft palate. Sixteen months previously severe trauma and operation had been tolerated without excessive bleeding.

Investigations showed haemoglobin 6.6 g/100 ml; reticulocyte count 1.7%; and W.B.C. 7,500/mm³ (normal differential). Red cells seemed mildly hypochromic and microcytic. Fragmented red cells

were not apparent. Platelets appeared severely decreased (6,900/mm³). The bone marrow was cellular, with increased megakaryocytes and absent stainable iron. The plasma fibrinogen level (ammonium sulphate precipitation method) was 420 mg/100 ml and the prothrombin time 15.5:13.0 seconds.

In view of a negative history, physical examination, and further laboratory investigations idiopathic thrombocytopenic purpura was diagnosed. Therapy with prednisone 40 mg by mouth daily was instituted. The response is shown in the Chart. Steroid administration on several occasions was associated with a rise in the platelet count, while a reduction in the dose was rapidly followed by a fall in platelet levels. Splenectomy was therefore considered and her coagulation status was re-evaluated. Plasma fibrinogen concentration was found to be reduced to 40 mg/100 ml (both Ellis-Stranski and ammonium sulphate precipitation techniques). The prothrombin time was 16.2:12.8 seconds and the thrombin time 22.6:15.6 seconds. Partial thromboplastin time was 39:39 seconds. The factor VIII level (one-stage technique) was 200% of normal. Cryofibrinogen was present and a few fragmented red blood cells were noted on the peripheral smear.

Disseminated intravascular coagulation was diagnosed. Prednisone 40 mg daily was continued and heparin 40 mg intravenously every six hours was begun with little effect. The heparin dose was therefore increased to 60 mg every six hours. There was then a prompt increase in platelet and fibrinogen levels, with disappearance of detectable cryofibrinogen. On the third day at the higher heparin dose the patient bled spontaneously into her left thigh. Heparin was discontinued, and within 24 hours the fibrinogen concentration had fallen from 165 to 30 mg/100 ml, and the platelet count from 70,000 to 42,000/mm³. Cryofibrinogen reappeared. Heparin therapy was reinstituted. Within 24 hours fibrinogen rose to 270 mg/100 ml and platelets to 62,000/mm³. Two days later bleeding into the thigh and flank recurred. Heparin was again discontinued, and both the platelet count and the fibrinogen again decreased.

A series of complications then supervened, including abscess formation in the thigh, Gram-negative septicaemia and shock, myocardial infarction, probably secondary to hypotension, congestive heart failure, ill-defined liver functional derangement, and mild acute tubular necrosis. Without apparent explanation platelets and fibrinogen then steadily increased to levels of 180,000/mm³ and 500 mg/100 ml respectively (not shown in Chart). Thrombin time was 8.4:7.4 seconds, prothrombin time 12.5:12.5 seconds, and cryofibrinogen was absent. The patient appeared to be improving clinically and she was allowed up. Twenty-seven days after the heparin was discontinued, however, she died suddenly and for no apparent reason. Necropsy

Thorndike Memorial Laboratory and Harvard Medical Service,
Boston City Hospital, Boston, Massachusetts

A. DAVID GINSBURG, M.D., F.C.P.(S.A.), Assistant Professor of Medicine,
(Present address: Queen's University Medical School, Kingston, Ontario,
Canada)

HERMAN A. GODWIN, M.D., Assistant Professor of Medicine, Harvard
Medical School.

RICHARD H. ASTER, M.D., Professor of Medicine, (Present address:
Milwaukee Blood Center, 763 North 18th Street, Milwaukee, Wisconsin)

failed to show any reason for her initial haematological problems but confirmed the presence of an infected haematoma in the thigh. There were no features of disseminated intravascular coagulation, but by this stage she had recovered clinically from this disorder.

Comment

This case illustrates the need to consider the diagnosis of disseminated intravascular coagulation in any patient with unexplained thrombocytopenia. There were no apparent precipitating factors or clinical features to suggest this disorder and, moreover, none were found at necropsy.

Diagnosis was dependent on the performance of routine laboratory investigations. A low fibrinogen concentration was particularly helpful, yet it was normal initially. That this should have been so emphasizes the need for serial assays of coagulation factors when a diagnosis of disseminated intravascular coagulation is being considered. Furthermore, a method measuring thrombin-clottable fibrinogen is to be preferred in these circumstances. The prothrombin time was slightly prolonged from the beginning, but did not alone raise the suspicion of disseminated intravascular coagulation. A long thrombin time was noteworthy. It is of interest that the partial thromboplastin time and the factor VIII level were normal. This re-emphasizes the observation that in disseminated intravascular coagulation a one-stage technique for measuring factor VIII levels may be misleading (Nossel *et al.*, 1969). Red cell fragmentation, absent at first, never was a major finding.

Confirmation of the diagnosis was strongly suggested by the response and sensitivity to heparin administration (see Chart). It is striking that a dose of 40 mg every six hours caused only a transient minor increase in fibrinogen and platelet levels, while a modest increase in dosage to 60 mg six-hourly was correlated with a significant rise in both (see Chart). A dilemma that is highlighted by this patient's illness is whether heparin should have been used at all and what indeed its role is in the treatment of disseminated intravascular coagulation.

The pathogenesis of the illness remains unknown. While idiopathic thrombocytopenic purpura is rare at this age—and hence not a diagnosis which should be accepted with equanimity

Platelet count, fibrinogen, and cryofibrinogen responses to steroid and heparin administration and withdrawal.

—the response to the administration and withdrawal of steroids seemed to support the diagnosis initially. It is probable, however, that the problem was one of disseminated intravascular coagulation *ab initio*. The steroid dependency is difficult to account for on this hypothesis and was certainly misleading. A fascinating alternative is that antigen-antibody reactions responsible for an initial idiopathic thrombocytopenic purpura-like illness precipitated subsequent intravascular coagulation.

This study was supported in part by grants from the National Institutes of Health (AM-00795, AM-05391, AM-05413, HE/AM-07652, AM-09115, and HE-12860-01).

References

- Nossel, H. L., Niemetz, J., Waxman, S. A. and Spector, S. L. (1969). *American Journal of Medicine*, **46**, 591.

Traumatic Diaphragmatic Transpericardial Hernia

H. BANK, H. SINKOVER, M. OPHIR
M. SHULEMSON

British Medical Journal, 1971, **2**, 629-630

Diaphragmatic tears due to blunt or penetrating wounds have been well documented since the original description by Ambroise Paré in the sixteenth century (quoted by Lindskog, 1961). We here report a case of traumatic rupture of the diaphragm and pericardium with herniation of abdominal viscera through the pericardial sac into the pleura.

Case Report

A 62-year-old writer was referred to the medical department for progressive anaemia. He had been admitted to hospital after a road accident 10 months previously with a fracture of the right acetabulum and blunt injury to the chest wall. The electrocardiogram showed ST elevations consistent with pericarditis, and in the case summary it was noted that "myocardial infarction cannot be excluded." He complained of weakness, effort dyspnoea, chest discomfort after meals, and intermittent constipation. He was pale, pulse 104, blood pressure 130/70. The chest was barrel-shaped and heart sounds were distant. Peristalsis could occasionally be discerned over the precordium. The abdomen appeared scaphoid in the prone position. The patient said that his abdomen had been protuberant before the accident and that he had noticed the sudden change and had even drawn attention to this fact.

Haemoglobin was 6.6 g/100 ml, haematocrit reading 26%. Occult blood was present in the stools. E.C.G. was normal. Chest x-ray film showed a large air bubble anterior to the heart shadow. On barium-enema examination part of the left colon was seen in the chest. Barium-meal examination showed volvulus of the stomach, with the fundus and pyloric region in the chest anterior to the cardiac shadow (Figs. 1 and 2).

When the lung was retracted at thoracotomy the heart was seen outside the pericardial sac, completely eviscerated. Most of the stomach and part of the large bowel together with the omentum were in the left pleural space, with a remnant of a torn mediastinal pericardium running across them (Fig. 3). With the abdominal viscera replaced a tear extending 15 cm could be seen through the base of the diaphragmatic pericardium and the diaphragm beneath it, running from the sternum posteriorly and to the left and through the muscular and tendinous layers of the dia-

Tel-Hashomer Government Hospital, Postgraduate Faculty of Tel-Aviv University Medical School, Israel

H. BANK, M.B., CH.B., Chief of Department of Medicine F
H. SINKOVER, M.D., Chief of Department of Surgery
M. OPHIR, M.D., Registrar in Surgery
M. SHULEMSON, M.D., Assistant in Surgery