CORRESPONDENCE

Correspondents are asked to be brief

Obstetric Appointments and the Abortion Act

Act	
C. K. Vartan, F.R.C.O.G.	767
Follow-up after Abortion	
L. A. Pike, M.B.	767
Control of Paraquat	
A. A. H. Lawson, M.D.	767
Accidental Vaccination with Paraquat	
P. J. Barber, M.B.	768
Treatment of Acute Poisoning	
W. O. Robertson, M.D.	768
Fats and Multiple Sclerosis	
G. Dean, F.R.C.P.	768
Aspirin Allergy	
G. M. Halpern, M.D.	768
Aortic Aneurysm and Peptic Ulcer	
A. P. Barabas, F.R.C.S., and others	768
Theatre Fire	
A. R. Elkington, F.R.C.S.	769
Cystic Degeneration of the Popliteal Artery in a Female	
J. C. Milliken, F.R.C.S.I.	769
Lymphocyte Sensitivity in Cancer	
D. Hughes, D.Sc., and D. W. Paty, M.D.	770
Adaptation of Houses for Home Dialysis	
S. G. Phillips, D.P.H.	770
Chloroquine Myopathy	
A. Ebringer, M.D.	770
Mutants, Hyperlipoproteinaemia, and Coronary Artery Disease	
L. M. Dalderup, M.D.	771

Pathogenesis of Myasthenia Gravis

G. A. Macgregor, M.D.	771
Accident and Emergency Services	
J. M. Potter, F.R.C.S.	771
Flying Squads for Road Accidents	
K. Little, M.B.	772
Infectiousness of Glandular Fever	
R. M. Whittington	772
Eczema and Detergents	
F. A. Ive, M.R.C.P.	772
Deaths from Tuberculosis	
R. A. L. Agnew, M.D.	772
Serum Transaminases during Salicylate Therapy	
M. Lee-Jones, M.R.C.P.	772
E.E.G. and Anticonvulsants	
W. I. M. Dow, M.B.	773
Diet and Diverticulosis	
R. C. Garry, F.R.C.P. GLASG.	773
Comparison of Drugs for Asthma	
L. Tivenius, M.D.	773
Analgesics in Terminal Disease	
P. B. Schofield, F.R.C.PATH.	773
Levodopa and Anticholinergic Drugs in Parkinsonism	
J. H. Burn, M.D., F.R.S.	773
Registrar Cover for Obstetric Unit	
J. J. Shipman, F.R.C.S.	774
False Economy	
M. B. Hay, M.B.	774

Treatment of Syringomyelia

B. N. Williams, CH.M.	774
Calcium Carbonate, Hypercalcaemia, and Peptic Ulcer	
D. W. Piper, F.R.A.C.P.	774
Care of Chronic Psychotics	
D. Tidmarsh, D.P.M.	774
Laryngeal Tuberculosis	
A. G. D. Maran, F.R.C.S.ED., and I. A. Stewart, F.R.C.S.	775
Closed Injuries to the Duodenum	
C. T. Hewetson, M.R.C.S.	775
Radiological Training	
E. Samuel, F.R.C.P.ED., F.F.R.; K. C. Simpkins, M.R.C.P., F.F.R.	775
Schistosomiasis after Bathing	
E. Agius, M.D.	775
Surgeons and Technical Skill	
M. A. Simpson, M.B.	775
Prescribing on Clinical Grounds	
G. Teeling-Smith, B.A.	776
Dental Emergencies	
M. P. Coplans, F.F.A. R.C.S.	776
Interpretation of Therapeutic Trials	
D. A. K. Black, F.R.C.P., and G. A. Rose, D.M.	776
Is Fetal Monitoring Worthwhile?	
P. N. Holberton, M.R.C.O.G.	776
Liver Damage and Methotrexate	
H. Baker, M.D.	776

Obstetric Appointments and the Abortion Act

SIR,—Mr. H. P. Dunn (22 May, p. 464) poses an important question. We have now had an Abortion Act for three years which in this permissive age permits doctors to relieve women of the very great suffering of bearing an unwanted child. Whether or not one avails oneself of this permission is left to the individual to decide, and every potential gynaecologist must face this question, as we have had to.

Those who will not terminate should realize, however, that they are passing the patient to another who may already be doing more than he or she likes, and since no one enjoys this work such practice is not going to be welcomed.

Of course one accepts that many with

deep convictions against termination are caught up in this situation and this is dealt with amicably, but taking on new recruits who will not help is a different matter. At my hospital we do not ask candidates if they will co-operate but we do insert a clause stating that terminations are carried out and we hope the message will get across.

As Mr. Dunn says: "The implications for future candidates for the M.R.C.O.G. are obvious." If they are obvious then should not those who will not terminate train for some other branch of medicine?—I am, etc.,

C. K. VARTAN

Memorial Hospital,
London S.E.18

Follow-up after Abortion

SIR,—Recently a 16-year-old patient became pregnant and, taking her boy friend with her, consulted an abortion agency. Neither her parents nor her doctor were informed, at the girl's request, but the mother became aware of the situation the day before the operation.

The family doctor became aware of the situation when, one month after the event, he was called by the mother to see the patient. The request for a visit was for fever, due to a tonsillitis, but the mother was worried in case the fever was a complication of the termination. Taking a history elicited the information that the termination had been said by the surgeon to have been opportune, because something was wrong

with the fetus, and that some follow-up was necessary. The doctor suspected that the patient had had a hydatidiform mole, and a request for information to the abortion agency elicited a reply, some two weeks later, that this was the case, and that there were no apparent arrangements for follow up, though the surgeon would now be asked to contact the doctor.

Now that the situation has been revealed, it is obviously the doctor's responsibility to arrange for the investigation and care of this patient. Who would have been responsible if chance had not brought the situation to his notice?—I am, etc.,

L. A. PIKE

Birmingham

Control of Paraquat

SIR,—In the past few years since the general introduction of paraquat as a weedkiller there have been recurrent reports of patients with acute poisoning due to this substance in both the medical and lay press. Although the numbers have not been unduly great this poisoning has caused concern because of the marked toxicity of paraquat and an associated high mortality. In contrast to the vast majority of other poisonings intensive supportive therapy, even when instituted at an early stage by the most experienced medical teams and using the most sophisticated types of equipment, does not save the patient from the inexorable deterioration which is rather characteristic of this condition. The fatal dose may be very small, and these deaths are all the more tragic as the poisonings are virtually all accidental and all too often the victim is a child.

The powdered form of paraquat is readily available to the public by sale over the counter and although the more commercial concentrated liquid form is controlled by schedule, there seems no doubt as events have proved that these controls are less than effective. It may of course be argued that instances of accidental poisoning are always the result of carelessness, irresponsibility, and even stupidity in the handling of the substance and that one cannot legislate against these individual failings. In view of the great dangers of paraquat ingestion I think that this provides a very good argument for withdrawal of paraquat from general use. Such action would of course