stages of cancer is likely to die. Not that such predictions are meaningless; the fact that 83% of these patients died within 12 weeks of admission is itself a confirmation that the patients were rightly regarded as being in a "terminal" state.

These patients cannot, of course, be regarded as completely typical of cancer patients because of the means of selection of cases for admission, but neither are they highly atypical. We have no reason to believe that cancer patients dying in south-east London are different from those dying elsewhere. Most cancer patients die in hospital (61% according to figures for England and Wales in 1968²), and although patients in the older age bracket are under-represented the sample did include many elderly patients.

One possible explanation for the consistent optimism which must be considered is the possibility that the treatment which is provided for patients at St. Christopher's Hospice actually shortens their lives. If this were the case we would expect that the physicians and nurses at the hospice, whom one would expect to adopt the institutional "norm" of survival, would be more realistic in their expectations than those who work in other settings. But this is not the case, nor was there any evidence that patients given diamorphine (the principal drug given for pain in that institution) died sooner than those who did not receive this drug. It seems, then, that this explanation must be set aside.

Because uncertainty is hard to bear relatives and even patients sometimes press their doctors to give them a precise estimate of prognosis. We often hear the phrase "The doctor gave him x

weeks to live," usually followed by a "but" and a disclosure of the magnitude of the doctor's error.

More often than not such anecdotes refer to pessimistic errors, and the patient's powers of survival despite the doctor's gloomy predictions are cited as evidence of his courage, determination, or laudable stubbornness. Our data suggest that doctors and nurses are more likely to be optimistic than pessimistic in their estimates at such times, a finding which suggests that their judgment is clouded by their hopes for the patient's survival or by their wish to reassure patients and relatives. It is almost as if, by doubling our realistic expectations, we can will a little more life into the patient.

Whatever the value of these speculations it does seem that until reliable indicators of prognosis have been established precise estimates of life expectancy in terminal cancer patients should be most cautiously given and interpreted.

This study was planned and initiated by the late Dr. Ronald Welldon shortly before his own unexpected death in 1969. It was carried out with the assistance of grants from the Department of Health and Social Security and the Sir Halley Stewart Trust. Thanks are due to Dr. Ann Cartwright for helpful comments on the draft.

References

- ¹ Glaser, B. G., and Strauss, A. O., *Time for Dying*. Chicago, Aldine, 1968.
- ² General Register Office, *Statistical Review of England and Wales for the Year 1968, Part 1, Tables, Medical*. London, H.M.S.O., 1970.

Scientific Basis of Clinical Practice

Control of Blood Pressure

IAN R. GRAY

British Medical Journal, 1972, 2, 31-33

Four mechanisms are recognized by which the blood pressure is maintained in normal circumstances. They are (1) catecholamine release, (2) the renin-angiotensin mechanism, (3) aldosterone-sodium chloride retention, and (4) the autonomic nervous system. The first three are hormonal and are concerned mainly with sustaining a background level of vascular tone, while the fourth is neurological and is responsible for fine adjustment of the blood pressure to transient changes in the condition of the body. Each of these mechanisms may be disturbed in pathological states so as to affect blood pressure, and any of them may be interfered with by medical or surgical means in the treatment of high or low blood pressure.

Catecholamine Release

The catecholamines, adrenaline and noradrenaline, are secreted by the adrenal medulla.

A functioning tumour of the adrenal medulla, the phaeochromocytoma, liberates excessive amounts of catecholamines into the blood stream causing hypertension which is often

paroxysmal. Other manifestations of sympathetic over-activity such as pallor, sweating, glycosuria, and increased metabolic rate also occur. The diagnosis of phaeochromocytoma is usually established by estimation of catecholamines in the urine, though an assay of blood levels is possible.

The manifestations of a phaeochromocytoma depend on the relative amounts of adrenaline and noradrenaline it produces. Adrenaline increases the pulse rate, the systolic blood pressure, and the cardiac output, while noradrenaline increases peripheral vascular tone and blood pressure. These different actions correspond with those mediated by beta-sympathetic and alpha-sympathetic transmission. Thus the effects of excessive circulating catecholamines in a patient with phaeochromocytoma can be blocked by using alpha-sympathetic and beta-sympathetic-blocking drugs.

Alpha receptors, which are responsible for vasoconstriction, hypertension, sweating, and contraction of the erector pilae muscles, may be blocked by such drugs as phenoxybenzamine or phentolamine. Beta receptors are blocked by drugs like propranolol. Alpha-sympathetic and beta-sympathetic-blocking drugs play an important part both in the diagnosis of phaeochromocytoma and particularly in the management of the patient during surgical treatment.

Another aspect of the differing effects of catecholamines is in their administration for therapeutic purposes. The administration of adrenaline makes good sense in patients with bronchial asthma or heart block, but is clearly contraindicated in hypo-

Coventry Hospitals, Warwickshire

IAN R. GRAY, M.D., F.R.C.P., Consultant Cardiologist

tensive states except when a positive inotropic effect on the myocardium is wanted. Noradrenaline, as an alpha-adrenergic drug, will effect vasoconstriction with a corresponding rise in blood pressure without any direct effect upon the heart.

Sympathetic-blocking drugs also have a place in the treatment of hypertension of other causes. Alpha-sympathetic-blocking drugs, such as phenoxybenzamine, are not used in the treatment of hypertension because they have to be given parenterally and their action is short-lived. However, beta-sympathetic-blocking drugs, notably propranolol, are effective antihypertensive drugs by virtue of their effect on the heart.¹ The bradycardia and the diminished force of myocardial contraction cause a significant fall in cardiac output. If the peripheral resistance remains constant the blood pressure falls correspondingly. The dosage of propranolol required for the treatment of hypertension is often large, and possibly some other pharmacological mechanism plays a part in its antihypertensive role.

Renin-angiotensin Mechanism

Many renal diseases can cause hypertension. They include acute and chronic nephritis, pyelonephritis, polycystic disease, irradiation nephritis, and renal artery stenosis.

Renal artery stenosis has given a particular impetus to research in renal hypertension because it is often unilateral and may respond to surgical treatment. Furthermore, experimental renal artery constriction has been a particularly fruitful research procedure since the pioneer work of Goldblatt.²

The mechanism by which renal ischaemia causes hypertension is primarily hormonal. Renin, which is secreted by the juxta-glomerular apparatus and which is present in the normal kidney, reacts with a precursor in the blood to form an active pressor agent, angiotensin. The mechanism by which angiotensin raises the blood pressure seems to be neurologically mediated,³ probably both locally and through the vasomotor centre of the brain.

Unfortunately the assay of renin and angiotension is difficult, and renin is best estimated from renal vein blood. A straightforward test by which renal hypertension can be detected does not therefore exist. An indirect method, measuring the blood pressure response to infused angiotensin,⁴ depends upon the lesser response in patients whose hypertension is caused by a high level of circulating endogenous angiotensin. The results of this test, however, have been of doubtful value, and it is not without risk. Other indirect methods exist by which renal ischaemia can be detected, and cases of unilateral renal disease can be selected in which nephrectomy or renal arterial surgery will probably be beneficial.

Angiotensin is available for the treatment of hypotensive states. Like noradrenaline, its value in therapy is difficult to assess but it is unquestionably effective as a pressor agent.

Aldosterone-sodium Chloride Mechanism

The relationship of adrenocortical steroids to the level of blood pressure is exemplified clinically by the hypertension of Cushing's syndrome and of Conn's syndrome and the hypotension of Addison's disease. The adrenal hormone responsible for the abnormal blood pressure in these conditions is aldosterone, a substance mainly concerned with the control of sodium and potassium excretion by the kidney.⁵ Excessive secretion of aldosterone causes an increase in excretion of potassium with retention of sodium, giving rise to hypokalaemia and hypernatraemia. There is an associated alkalosis which is probably due to a further effect of aldosterone interfering with urine acidification. While the biochemical disturbances usually dominate the clinical situation, aldosterone excess also gives rise to a hypertension which tends to be no more than moderate. This mechanism of hypertension was recognized experimentally

with the hypertension and fluid retention which could be produced by administration of desoxycorticosterone acetate together with salt.⁶ Clinically it occurs when there is adrenocortical hyperplasia, an aldosterone-secreting tumour, or stimulation of the adrenals with ACTH or its simulation with cortisone. The effect of excessive circulating aldosterone can be modified by aldosterone antagonists such as spironolactone or by giving a diet abnormally high in potassium or low in sodium.

Assay of aldosterone is difficult, and the diagnosis of primary aldosteronism tends to be reached indirectly from the biochemical findings or by the radiological detection of an adrenal tumour. When there are excessive amounts of other adrenocortical steroids which can be more easily estimated, as in Cushing's syndrome, the diagnosis can more readily be made. Likewise the deficiency of adrenocortical steroids in Addison's disease seldom presents a difficult diagnostic problem.

The antihypertensive effect of sodium depletion has been widely used in the treatment of hypertension. It was the basis of the Kempner rice diet,⁷ which was a most effective if disagreeable form of treatment. It was possible with this diet to reduce the daily intake of sodium chloride to less than one gramme, and even severe hypertension responded to this. Another application of the interference with the aldosterone mechanism was the treatment of severe hypertension by bilateral adrenalectomy and minimal replacement therapy.⁸ This form of treatment, like the rice diet, has been superseded with the development of the modern range of antihypertensive drugs.

When effective oral diuretics which could cause significant sodium loss were introduced sodium depletion was re-established in the treatment of hypertension. The benzothiadiazine diuretics are now important in antihypertensive therapy. Nevertheless, their mode of action seems less straightforward than was at first thought. During the first few weeks of diuretic therapy there is increased sodium excretion with a reduction in plasma volume and a fall in blood pressure. The plasma volume then returns to normal but the blood pressure remains at the lower level, so that in the long term these diuretics seem to affect the peripheral vascular resistance as well as the excretion of sodium and water.⁹

Autonomic Nervous System

The three endocrine mechanisms of blood pressure control have interesting clinical reflections in certain diseases associated with hypertension but have only limited relevance to the treatment of hypertensive patients. The autonomic nervous control of the blood pressure, however, has no recognized relevance

FIG. 1
C.S.=carotid sinus. A.A.=aortic arch. V.M.C.=Vasomotor centre. H.C.=higher centres. V.=vagus. C.S.—cardiac sympathetic. S.—sympathetic nerves.

to hypertensive disease, but is crucial in the therapy of hypertension.

The nervous pathways responsible for minute to minute control of the blood pressure (Fig. 1) begin with the baroreceptors in the carotid body and aortic arch. When these are stretched by an increase in blood pressure afferent stimuli pass from them by the sympathetic nerves to the vasomotor centre in the midbrain, from which efferent impulses pass to the heart and blood vessels, restoring the status quo. The efferent flow is by the vagus and the cardiac sympathetic nerves to the heart and by the sympathetic nerves to the blood vessels (Fig. 2). The pathway is via the spinal cord to cells in the grey matter and

SYMPATHETIC CONTROL OF VASCULAR TONE

FIG. 2
V.M.C. = Vasomotor centre. G.B. drug = ganglion blocking drug.

from there to the effector cells of the sympathetic ganglia. The synaptic transmission in the sympathetic ganglia is mediated by acetylcholine, as in the synapse of the parasympathetic. The nerve endings of the effector cells in the walls of the blood vessels are, by contrast, adrenergic—the alpha-adrenergic receptors responsive to noradrenaline.

GANGLIOPLEGIC DRUGS

The first step in the treatment of hypertension by interfering with the autonomic control of the blood vessels came with the discovery by Paton and Zaimis in 1948¹⁰ of the autonomic-blocking properties of the hexamethonium and pentamethonium compounds. These substances have a ganglioplegic effect on the acetylcholine mediated synapses of the autonomic nervous system. Hence they have blocking effects on both the sympathetic and parasympathetic outflows. They cause a reduction of blood pressure through a fall in vascular tone which is more pronounced in the erect position, for in this position the neural response to baroreceptor afferents makes a greater contribution. Other effects are an increase in heart rate owing to vagal block, dry-mouth, paralysis of accommodation, constipation, and urinary retention from parasympathetic blockade. Though a succession of more satisfactory ganglioplegic drugs appeared they all had the same troublesome side effects, and a drug with a selective sympathetic blocking action was obviously desirable.

SYMPATHETIC BLOCKING DRUGS

The sympathetic blocking drugs provided the advance which eliminated most of the untoward side effects of the autonomic ganglioplegic drugs. Their exact site of action is unknown, but it is either in the sympathetic effector cell or its endings in the blood vessel, a point of adrenergic rather than cholinergic transmission.¹¹ Two drugs of this category, guanethidine and bethanidine, have continued to be widely and successfully used. They may cause postural hypotension, but most of the other

unpleasant and sometimes dangerous complications of ganglion-blocking drugs are eliminated.

METHYLDOPA AND RESERPINE

Two other drugs widely used in the treatment of hypertension, methyldopa and reserpine, may interfere with the vascular tone by their effect on the noradrenaline transmission at the sympathetic nerve endings. Methyldopa is a decarboxylase inhibitor, thus causing a decrease in concentration of noradrenaline in the tissues¹² and its displacement by a "false" transmitter, alpha-methylnoradrenaline. This suggests that its antihypertensive effect might be due entirely to deficiency of noradrenaline at the sympathetic nerve endings. However, this hardly seems likely in view of the differences between its effect and that of guanethidine—for example, postural hypertension is seldom caused by methyldopa. A central component to its effect has been demonstrated, but this too seems inadequate to explain its antihypertensive effect. Reserpine causes depletion of catecholamines in various parts of the body,¹³ including the postganglionic sympathetic nerves, and its antihypertensive effect is probably through this mechanism.

The various mechanisms in the body which control the blood pressure, the levels of circulating catecholamines, the secretion of renin by the kidney, the aldosterone control of electrolyte

Mode of Action of Antihypertensive Agents

Type of Drug	Examples	Site and Mechanism of Action
Ganglion-blocking drugs	Pentolinium Mecamylamine	Cholinergic autonomic ganglia
Sympathetic-blocking drugs	Guanethidine Bethanidine	Postganglionic sympathetic neurone—adrenergic
Beta-adrenergic-blocking drugs	Propranolol	Beta-adrenergic neuromuscular transmitters in the heart
Decarboxylase inhibitor	Methyldopa	Causes tissue depletion of noradrenaline and its displacement by "false transmitter"
Benzothiadiazine diuretics	Bendrofluazide	Increased sodium excretion causing reduced blood volume, also reduction in peripheral vascular resistance
Catecholamine depletor	Reserpine	Reduction of postganglionic sympathetic activity by depletion of transmitter substance

metabolism, and the nervous pathways to and from the heart and blood vessels may be disturbed in disease states to give rise to hypertension or hypotension. Most forms of antihypertensive therapy can be understood in terms of interference with one or other of these mechanisms (see Table). Nevertheless, there are still doubts about the means by which some clinically effective drugs have their effects.

This article is based on a lecture given in the Birmingham course under the title "The Scientific Basis of Clinical Practice" (see B.M.J. 27 November 1971, p. 510).

References

- Prichard, B. N. C., *British Medical Journal*, 1964, 1, 1227.
- Goldblatt, H., Lynch, J., Hanzal, R. F., and Summerville, W. W., *Journal of Experimental Medicine*, 1934, 59, 347.
- Peart, W. S., *Pharmacological Review*, 1965, 17, 143.
- Kaplan, N. M., and Silah, J., *Journal of Laboratory and Clinical Medicine*, 1963, 62, 890.
- Ross, E. J., *Aldosterone in Clinical and Experimental Medicine*, Oxford, Blackwell Scientific Publications, 1959.
- Selye, H., in *Hypertension*, ed. E. T. Bell, p. 119. Minneapolis, University of Minnesota Press, 1951.
- Kempner, W., *North Carolina Medical Journal*, 1944, 5, 125.
- Thorn, G. W., et al., *Annals of Internal Medicine*, 1952, 37, 972.
- Conway, J., and Palermo, H., *Archives of Internal Medicine*, 1963, 111, 203.
- Paton, W. D. M., and Zaimis, E. J., *British Journal of Pharmacology*, 1949, 4, 381.
- Page, I. H., and Dustan, H. P., *Journal of the American Medical Association*, 1959, 170, 1265.
- Gillespie, L., Oates, J. A., Crout, J. R., and Sjoerdsma, A., *Circulation*, 1962, 25, 281.
- Bein, H. J., *Pharmacological Review*, 1956, 8, 435.