CORRESPONDENCE

Correspondents are asked to be brief

Holiday Cholera and Other Vibrios G. I. Barrow, M.D.	403	Diabetics and Motorway Crashes G. E. Leyshon, D.P.H., and others	405	Bran and Diverticular Disease T. L. Cleave, M.R.C.P.	408
Digoxin Dosage R. G. Simpson, F.R.C.P.ED.	403	Total Body Potassium and Chronic Renal Failure Y. K. Seedat, M.D.	405	Varieties of Memory T. D. Culbert, M.B.	409
Survival of Gonococci Outside the Body T. Elmros, M.D. and P.-A. Larsson, M.D.	403	Urinary pH and Excretion of Gentamicin C. Mariel, M.D., and others.	406	New Thoughts on Essential Hypertension G. J. Sophian, F.R.C.O.G.	409
Battered or Pigmented? I. G. Wickes, F.R.C.P., and Z. H. Zaidi, M.R.C.P.ED.	404	Smoking in Pregnancy J. W. B. Forshaw, F.R.C.P.	406	Sabbatical Year T. J. Wilmot, F.R.C.S.	409
Presenile Dementia H. M. Hodkinson, M.R.C.P.	404	Truncal Vagotomy and Pyloroplasty A. Mitra, F.R.C.S.ED.	407	Family Doctors and Contraception B. L. Furman, PH.D., and others.	409
Asthma and a Lump in the Breast S. G. Spiro, M.R.C.P., and others; M. Harris, M.R.C.PATH.	404	Value of the M.C.V. N. K. Shinton, M.D., and P. C. Elwood, M.D.	407	Fair Wage for Auxiliaries R. Brooks, F.I.M.L.T.	409
Alpha Receptors and Control of Blood Pressure A. V. P. Mackay, M.B.	404	Vaccination against Rubella J. S. Dodge, D.P.H.	407	Future of Mental Health Services J. T. Leyburg, M.D., D.P.M., and Marion E. Mackay, M.R.C.P., D.P.M.	410
Asystole after Verapamil D. M. Kirkler, F.R.C.P.ED., and R. A. J. Spurrell, M.R.C.P.	405	Pantie-girdle Syndrome T. K. Davidson, M.B., D.PHYS.MED.	407	Revised Regulations for G.P.s G. Murray Jones, M.R.C.S.	410
Azathioprine in Connective Tissue Disorders A. J. Richards, M.R.C.P., and others.	405	Inexpensive One-way Screen Elizabeth S. Barton, B.A.	407	G.M.C. Annual Retention Fee P. J. Waddington, M.B.	410
		Unbalanced Deoxyribonucleotide Synthesis Caused by Methotrexate M. H. N. Tattersall, M.R.C.P.	408	Hospital Staffing G. I. B. da Costa, F.R.C.S.ED.	410
		Removal of Extruded I.U.D. R. R. Macdonald, F.R.C.O.G.	408	G.P.s in District Hospitals P. A. Lawrence, M.B.	410

Holiday Cholera and Other Vibrios

SIR,—With the approach of the summer season, your warning (8 April, p. 62) to all concerned about the hazards of holiday cholera is timely. As you rightly indicate, the clinical picture may vary from acute gastroenteritis to mild diarrhoea, or even symptomless excretion of the organisms. *Vibrio cholerae* is not, however, the only vibrio which may cause such illness in travellers returning from abroad. The actual diagnosis depends on isolation of the causative organism, and other vibrios, particularly *V. parahaemolyticus*, should also be considered.

This halophilic marine vibrio was first identified in Japan, where it is associated with the consumption of raw fish and other sea-food products, and is now the commonest cause of food poisoning in summer.¹ It is almost certainly widely distributed throughout the world and is becoming more generally recognized as an important or potential pathogen. Its significance as an endemic cause of food poisoning in Britain is not yet known, although its presence in various marine products has been confirmed recently.² It may also be imported not only by travellers but also commercially in sea-foods. *V. parahaemolyticus*, like cholera and other enteric pathogens, should therefore be sought in all cases of gastroenteritis in persons who have recently returned from abroad, and also from anyone who has consumed sea-foods. The possibility of cross-contamination to other foods should not be forgotten. Indeed, for the first time in Britain, *V. parahaemolyticus* has just been identified as the cause of acute gastroenteritis among airline passengers who had eaten cooked crab-meat served as hors d'oeuvres during a flight from Bangkok to London.³ Apart from the Far East, food poisoning caused by this organism has recently been reported also in Australia⁴ and the U.S.A.⁵

As in cholera, *V. parahaemolyticus* is ex-

creted in large numbers during illness, but the vibrios diminish rapidly with clinical recovery. The role of possible carriers is not known, but direct transfer from one person to another does not seem to be important. Adequate clinical and epidemiological information is essential to ensure that suitable cultural techniques are used for appropriate specimens. Fortunately, *V. parahaemolyticus* can be isolated by the same media and methods used for cholera organisms, though salt-colistin or glucose-salt-teepol broth may also be employed for enrichment culture.¹ *V. parahaemolyticus* forms large characteristic green colonies on thiosulphate citrate bile-salt sucrose (TCBS) medium in contrast to the yellow sucrose-fermenting colonies of *V. cholerae*. Whatever media used, however, any Gram-negative organisms which are motile, oxidase- and catalase-positive, and sensitive to methylene blue⁶ and vibriostatic agent 0/129 (2, 4-diamino-6, 7-di-isopropyltetridine phosphate)⁷ should be regarded as vibrios and further identified. If this were done, infection with non-cholera vibrios, including *V. parahaemolyticus*, might be recognized more often.—I am, etc.,

G. I. BARROW

Public Health Laboratory,
Royal Cornwall Hospital (City), Truro

¹ Sakazaki, R., in *Foodborne Infections and Intoxications* (ed. H. Riemann), p. 115. New York, Academic Press, 1969.

² Barrow, G. I., and Miller, D. C., *Lancet*, 1972, **1**, 485.

³ *British Medical Journal*, 1972, **1**, 701.

⁴ Bartey, Y. M., Wallace, R. B., Allan, B. C., and Keeffe, B. M., *Medical Journal of Australia*, 1970, **1**, 430.

⁵ *Morbidity and Mortality*, 1971, **20**, 356.

⁶ Yan, W. K., *Journal of Medical Laboratory Technology*, 1969, **26**, 90.

⁷ Shewan, J. M., Hodgkiss, W., and Liston, J., *Nature*, 1954, **173**, 208.

Digoxin Dosage

SIR,—The need for individual adjustment of digoxin dosage in elderly patients is now widely recognized. It is particularly impor-

tant to reduce maintenance dosage to a minimal effective level. Even the conventional dose of 0.25 mg daily may be excessive for certain patients, and it is here that one runs into practical problems. I can prescribe either a half tablet dose—that is, 0.125 mg—or a Paediatric-Geriatric ("P-G") tablet of 0.0625 mg strength. Such complex decimals and unwieldy numbers are inconvenient and faintly ludicrous, not to say potentially hazardous. Such awkward fractions may also lead to a reluctance to prescribe sufficiently small doses of digoxin whenever indicated.

Why can we not have digoxin tablets in two distinct, sensible, and easily-written strengths—0.2 mg and 0.1 mg? Tablets of 0.1 mg would be scored so as to be easily divided into doses of 0.05 mg. The drug would then be prescribed in three strengths for adults—0.2 mg, 0.1 mg, and 0.05 mg—instead of the present clumsy doses of 0.25 mg, 0.125 mg, and 0.0625 mg, respectively. The former doses are more succinct and more sensible; admittedly they are fractionally smaller than the present standard doses but probably the better for that.

Lastly, it must always be remembered that a long-accustomed dose of digoxin in an elderly person may at any time become excessive—for example, in the presence of potassium deficiency, as may be caused by potent diuretics, or defective renal function. It used to be said, "once digitalis, always digitalis," but is prolonged maintenance treatment always really necessary? Very often it is, albeit sometimes in an infinitesimal dose.—I am, etc.,

R. G. SIMPSON

Geriatric Unit,
Burghmuir Hospital,
Perth, Scotland

Survival of Gonococci Outside the Body

SIR,—The possibility of acquiring gonorrhoea by non-venereal contact has been considered.¹⁻⁴ In order to shed further light on