CORRESPONDENCE

Correspondents are asked to be brief

Future of Postgraduate Medical Centres

R. B. Hunter, F.R.C.P.; D. E. Sharland, M.D. 710

Planning the Treatment of Alcoholism

C. E. Salter, D.P.M. 710

Treatment of Early Breast Cancer

J. W. S. Rickett, F.R.C.S., and A. G. Nash, F.R.C.S.; G. J. A. Edlsten, M.D., F.F.R., and K. D. MacRae, PH.D.; E. S. Lee, F.R.C.S., and others. 711

Chromosome Banding

A. Caratzali, M.D. 712

Stress and the Schizophrenias

S. Jacobs, D.P.M. 712

Aetiology of Varicosity

G. S. Ramsay, F.R.C.S.; G. B. Wyatt, M.R.C.P.; D. H. Patey, F.R.C.S. 712

Fish Cornea for Grafting

M. Haq, M.B. 712

Total Replacement of the Hip

H. Phillips, F.R.C.S., and D. J. Dandy, F.R.C.S. 713

Coalminers' Pneumoconiosis

I. T. T. Higgins, F.R.C.P. 713

Confusion with Clofibrate

J. T. Hart, M.R.C.G.P.; J. R. O'Brien, F.R.C.PATH 713

Legal Detention of Psychiatric Patients

W. A. Heaton-Ward, D.P.M. 713

Value of Thermography

E. F. J. Ring, F.I.M.L.T. 713

New Electronic Metal Locator

H. J. Hambury, F.R.C.S., and J. Watson, PH.D.; J. A. Ross, F.R.C.P.ED. 714

Abortion Act

C. B. Goodhart, PH.D. 714

Termination of Pregnancy

H. C. McLaren, F.R.C.O.G. 714

Urinary Candidiasis after Renal

Transplantation

R. J. Holt, PH.D., and R. L. Newman, M.D. 714

Preclinical Detection of Dystrophin

Myotonia

J. E. Caghey, F.R.C.P. 715

Payment by Colour

W. P. P. Leary, M.R.C.P. 715

Treatment of Trigeminal Neuralgia

S. L. Drummond-Jackson, L.D.S.R.C.S. 715

Feto-maternal Transfusion

I. Kennedy, F.R.C.S.ED. 715

Hyperthyroidism

M. C. Bateson, M.B. 715

Pentazocine in Thyroid Failure

Bethan M. Evans, M.B., and others. 716

Verapamil in Cardiac Arrhythmias

H. S. Sacks, M.B., and B. M. Kennelly, M.R.C.P. 716

Vasectomy

I. Craft, F.R.C.S. 716

New Consultant Contract

M. K. Strelling, M.R.C.P. 716

Evidence to the Lane Committee

A. W. Purdie, F.R.C.O.G. 717

Scientific Basis of Clinical Practice

C. D. Swaine, M.B. 717

Hospital Staffing

C. A. Wells, F.R.C.S. 717

Future of Postgraduate Medical Centres

SIR,—I read with interest the letter addressed to you by the chairman and secretary of the National Association of Clinical Tutors about the future of postgraduate medical centres (3 June, p. 589). I may have misunderstood this letter and, if so, would ask to be forgiven for rushing into print, but it seems to me that the writers have no ideas about the development of integrated medical services for the future, and have little knowledge of the vital part played by the professions allied to medicine and the necessity for their learning to work together in what inevitably must be a team effort.

If Drs. J. Lister and D. Ferriman are correct then the University of Birmingham is quite on the wrong track, because we are planning massive developments in the field of health services management, a major feature of which will be courses for all senior members of the health services, whether doctors or not. It is proposed to link these courses closely with similar ones for local government officers. The criticism of the Department of Health and Social Security for concerning itself primarily with the development of postgraduate and postexperience centres which are shared by all would seem to me to be quite unwarranted.

The postgraduate medical centre movement is about ten years old. It has been given massive support by the Department, but I hope that those who have been so active in making this a success will recognize the new relationships which will have to be developed between the medical profession and other professions if the National Health Service is going to deliver the goods at reasonable cost.—I am, etc.,

ROBERT B. HUNTER

University of Birmingham

SIR,—As director of one of the busiest postgraduate academic centres in Britain, I would fully support Drs. J. Lister and D. Ferriman (3 June, p. 589) in that doctors should retain control of the postgraduate centres. I do, however, fully support the administrators that when the centres are not being used they should be available for other medical and paramedical uses, thus ensuring maximum return of the capital.—I am, etc.,

D. E. SHARLAND

Whittington Hospital,
London N.19

Planning the Treatment of Alcoholism

SIR,—It is unfortunate that your leading article (27 May, p. 479) has adversely criticized a sentence quoted from the King Edward Hospital Fund for London report¹ which is itself the result of an editing error arising in an early stage of the work, and which was repeated unnoticed, despite many readings by myself and others in subsequent

revisions prior to final printing and publication. The offending sentence should read "as soon as alcoholism is detected the patient should be persuaded to see an expert and if necessary advised to enter hospital." This should redress the balance and alter the emphasis in subsequent paragraphs.

The report was sponsored primarily as an

architectural study (hence the "square footage of the cleaners' room" to which you took exception) and attempted to correlate therapeutic principles with design of buildings. It is hardly surprising, therefore, if the main emphasis of the book is on the structure (both functional and architectural) of inpatient units.

That outpatients care is always considered as a possible primary treatment is more than suggested in the section on Assessment (p. 25), where, among the manifold purposes of assessment listed, is found: "to assess the best method likely to help him achieve his resolve to gain sobriety (including, possibly, inpatient group therapy), to advise on the best method . . . to offer outpatient follow-up appointments, etc. . . ."

Far from stressing the inpatient service, I have been an active advocate of community psychiatry, and since 1964 have stressed the need for outpatient services for alcoholics, both ab initio and as follow-up services to an inpatient programme with community links. Indeed a careful study of some of the diagrams in the report shows, I hope, the development of this philosophy, even if this seems not to be spelled out too emphatically in the text. Recognizing the toll taken by the family and community aspects of the illness, and attempting to do something about it has for long been an in-built part of the total alcoholism services provided from Warlingham Park Hospital by the Croydon psychiatric service.

Despite the protest that the report does not take account of the "worrisome realities" and "does not consider the community side of things," there is in fact special mention of the lack of manpower in the medical profession, the need for reliable comparative research into methods of treatment, and possible ways that these needs may be met by using skill and expertise of

other professionals together with semi-professional workers and voluntary organizations.

Far from undermining self confidence, a quiet demonstration of understanding the alcoholics' plight (even if there is no one solution) is often the first stimulus towards regaining it.—I am, etc.,

CLIFFORD E. SALTER

London S.W.1

¹ James, W. P., Salter, C. E., and Thomas, H. G., *Alcohol and Drug Dependence—Treatment and Rehabilitation*. London, King Edward's Hospital Fund, 1972.

Treatment of Early Breast Cancer

SIR,—From the evidence in the excellent paper by Sir Hedley Atkins and others (20 May, p. 423) it would appear that there is no case for the operation of "extended tylectomy" in the management of stage I and II carcinoma of the breast in patients over 50 years of age.

Before all forms of local surgery in carcinoma of the breast are abandoned several questions might be asked concerning "extended tylectomy." In some patients a more adequate margin that the stated 3 cm can be easily removed without distortion of the breast contour. Was more than this margin ever excised? If skin dimpling is present is skin removed? In the subareolar lump is the nipple removed?

Answers to these questions would help in the evaluation of "extended tylectomy." Spread of carcinoma within the breast is by periductal lymphatics¹ and therefore to eradicate the disease from within the breast excision of breast tissue on a lobular basis is logical. Support for this concept is obtained by whole organ sectioning, a laborious task. In breasts removed for primary carcinoma intraduct changes of epitheliosis, in-situ carcinoma, and frank invasion are commonly seen and are most marked in the lobule containing the carcinoma. When there is clinical evidence of skin involvement an adequate clearance can be obtained only by excising skin with the tumour.

Surely there are many factors which influence the surgeon in his assessment of how much tissue to excise around the tumour. To stick rigidly to a fixed margin clear of the tumour is not always as appropriate as adopting a more flexible policy (according to various clinical criteria). It is of paramount importance to excise an adequate margin, which should be at least 3 cm from the edge of the tumour, as measured from the macroscopic appearance of the bisected lump. This policy is more reliable than taking the measurement from the palpable tumour edge. If skin tethering is present a good margin of skin clear of the tethered area should be included in the excision. In the well-endowed breast it is easy to excise more tissue and this should be done, in each case removing the segment of the breast affected in order to remove the involved lobule in its entirety.

Not often discussed is the relative size of the lump to the breast. This ratio is often a more pertinent factor in the formation of a treatment policy, particularly with regard to the small lump in a large breast, than the overall palpable tumour size such as is used in Manchester staging. Great variation in breast size exists between in-

dividuals as between nations such as Japan and Italy. An adequate clearance in the Tokyo breast is not as easily achieved as a similarly adequate clearance in its Neapolitan counterpart.

Incidentally the word "tylectomy" combines both Greek and Latin roots and therefore is no improvement upon "lumpectomy."—We are, etc.,

J. W. S. RICKETT
A. G. NASH

Royal Marsden Hospital,
London S.W.3

¹ Gallager, H. S., and Martin, J. E., in *Annual Clinical Conference on Cancer, 1968: Breast Cancer Early and Late*. Chicago, Year Book Medical Publishers, 1970.

SIR,—It is well recognized that the treatment of what is termed "early breast cancer" is a confused morass and therefore any effort to contribute objectively to the state of knowledge must be welcomed. We were therefore interested to see your leading article (20 May, p. 417) refer to the article by Sir Hedley Atkins and others in the same issue as a model example of the work which should be done in this field. On reading the article, however, we must confess to considerable disappointment and we must disagree strongly with the leading article on at least two points. (1) The numbers entering this trial are adequate to draw only the crudest conclusions. (2) The treatments being compared are incorrectly labelled by the authors.

As the authors themselves admit, the number of cases under study for the full 10 years is so few that conclusions must be drawn only with the greatest caution. Unfortunately, Sir Hedley Atkins and colleagues do not publish the details of their life table, presenting merely graphs without stating the number of cases the points are based on; nor do they even mention the degrees of freedom of their unnamed significance test. This does not prevent them, however, stopping their trial for stage 2 patients on the basis of a significant difference in survivals found at the 10-year point. The simplest assumption the reader can make is that the patients were evenly spaced throughout the 10 years, giving an estimated 37 patients followed up for 10 years, or approximately 9 per treatment group. Is the decision to stop the trial based on the comparison of two groups of about 9 patients, in which the death of one patient would change the percentage of survivals by over 10%? Further, even if all the 370 patients had been followed up for the full 10 years, giving about 90 patients per group, the treatment difference detectable at the 5% significance level and 95% power would be in the order of 25%; that is, if one treatment produced 50% survival, the other would need to have a true survival rate of 75%. Thus, a trial based on such small numbers can detect only the crudest differences between treatments.

The second criticism is even more damning. Sir Hedley Atkins and colleagues label their treatments according to the type of surgery performed. However, as they point out, this was not the only difference between the groups as radiotherapy varied considerably. Thus the excision group in addition to treatment to the breast received 2,500 rads in 12 days to the glandular areas, while the radical group received this

same dose in 18 days. These cannot be considered comparable because of the different time factors involved. It may be estimated that the equivalent dose to the conservatively treated group would be 3,000 rads, which is considerably greater in its biological effectiveness. Even if significant differences are found in this trial it is presumptive to attribute them to the type of operation while ignoring the differences in radiotherapy. Indeed, a growing body of opinion recognizes that immunological mechanisms play a part in the tumour/host conflict and radiation is a recognized suppressant of this system.

While the authors claim that the reader should draw his own conclusions and that the trial is essentially factual, their decision to apply radical mastectomy to all stage 2 patients and to discontinue this part of the trial implies a recommendation of this procedure. Both because of the small amount of data, and, we believe, its misinterpretation, this is to be deplored, and your congratulations to these authors is, to us, regrettable.—We are, etc.,

GEORGE EDELSTYN

Northern Ireland Radiotherapy Centre,
Belfast

K. D. MACRAE

Queen's University of Belfast

SIR,—While we are in complete agreement with Sir Hedley Atkins and others (20 May, p. 423) on the importance of controlled clinical trials, we would point out that the value of any conclusions reached depends to a considerable degree upon the questions put to trial.

The authors state that their aim was to compare radical mastectomy with extended tylectomy. Adjunctive chemotherapy was discontinued after the first seven years of the trial but all patients received radiotherapy to those areas not covered by operation.

The doses of radiation quoted in this trial are on average 2,600 rads tumour dose in 14 treatments over 18 days at 300 kV, given to the lymphatic drainage areas of those patients who had a radical mastectomy. An average tumour dose of 3,650 rads in 15 treatments over 19 days at 6 MeV was given to the residual breast tissue and internal mammary chain after tylectomy. Using the formula of Ellis¹ and allowing for relative biological efficiency, the nominal standard dose (N.S.D.) of these regimens is 1,164 ret (rad equivalent therapy) and 1,378 ret respectively. Adenocarcinomas are considered to be moderately radioresistant, and doses approaching the tolerance of normal tissues are needed to sterilize them. Many radiotherapists feel that up to 6,000 rads given in six weeks (N.S.D. approximately 1,800 ret) is necessary to prevent local recurrence of carcinoma of the breast. Fletcher² has recently shown that a regimen giving a similar N.S.D. to the treatment in this trial can be expected to control subclinical disease in only 60-70% of instances while 4,500-5,000 rads in five weeks (average N.S.D. 1,494 ret) will achieve control in more than 90%. Shukovsky³ has shown that in other sites where local control has been related to dose, small differences in total dose at the upper limit of tolerance produce a significant difference in the tumour response.