tumour mass, or enlarged nodes that had been increasing rapidly. The apparently complete remission was obtained after two courses of treatment.

Solid Tumours.—The overall results are summarized in Table VI. Out of 119 patients treated, one with bladder cancer had an apparently complete regression after a seven-day course of 50 mg/day; six had incomplete regressions, four of which were in patients with bladder cancer; there were 19 partial failures and 93 total failures.

Discussion

This clinical screening has shown that epipodophyllotoxin has a pronounced cytostatic effect on certain human tumours. The effect of this drug in reticulosarcoma is of great interest, for up till now a similar percentage of objective regressions was obtained only with cyclophosphamide (Carbone and Spurr, 1968; Hyman and Cassileth, 1966) and vincristine (Carbone and Spurr, 1968; Desai *et al.*, 1970). In view of these results it is tempting to propose the use of this drug not only for the primary treatment of reticulosarcomas in stages III and IV, but as a chemotherapy complementary to radiotherapy for stages I and II. With this aim in mind, some physicians in this group are giving epipodophyllotoxin intermittently (150 mg every second week) over long periods.

The action of epipodophyllotoxin in Hodgkin's disease and in lymphosarcomas is equally notable. However, in these diseases the drug is of less importance than in the reticulosarcomas, for each of these diseases is already known to be sensitive to several drugs.

The single apparently complete regression and the three partial regressions of bladder cancers are most encouraging, and this will be examined further by complementary trials by the Urogenital Cancer Co-operative Group of E.O.R.T.C.

Finally, we draw attention to the resolution of pleural effusions in five patients which were apparently complete and fairly long lasting (four to five months); such results are only exceptionally achieved with other forms of systemic chemotherapy. We are now studying the possible mesothelial tropism of this drug.

Clearly it is the toxic action on the bone marrow that is the main factor limiting the use of this drug. This might be overcome by using the drug in combination with another agent which shows some action against the same types of tumours that are affected by epipodophyllotoxin, but which is not toxic to the bone marrow; bleomycin has these characteristics (E.O.R.T.C., 1970, 1972).

Requests for reprints should be addressed to the Institut de Cancérologie et d'Immunogénétique, Hôpital Paul-Brousse, 14 Avenue Paul Vaillant Couturier, 94 Villejuif, France.

VM 26 was supplied by Sandoz Laboratories.

References

- Carbone, P. P., and Spurr, C. (1968). *Cancer Research*, **28**, 811.
- Desai, D. V., Ezdinli, E. Z., and Stutzman, L. (1970). *Cancer (Philadelphia)*, **26**, 352.
- E.O.R.T.C., Clinical Screening Co-operative Group (1970). *British Medical Journal*, **2**, 643.
- E.O.R.T.C., Co-operative Group for Leukaemia and Reticulocytoses (1972). *British Medical Journal*, **1**, 285.
- Hyman, G. A., and Cassileth, P. A. (1966). *Cancer (Philadelphia)*, **19**, 1386.
- Stähelin, H. (1970). *European Journal of Cancer*, **6**, 211.

MEDICAL MEMORANDA

Gold-induced Thrombocytopenia and Response to Dimercaprol

J. M. ENGLAND, D. S. SMITH

British Medical Journal, 1972, **2**, 748-749

Several types of thrombocytopenia may occur during gold therapy. Commonly the patient's platelet count falls, the gold therapy is temporarily discontinued, and a rapid recovery ensues. On other occasions a more profound and persistent thrombocytopenia occurs (Meyler and Herxheimer, 1968), and these cases may have either an aplastic marrow or, more unusually, changes suggestive of toxicity to the megakaryocytes.

We report here the case of a patient who recovered from a severe and persistent thrombocytopenia. In the literature are only a few examples of successful treatment of this condition (Lockie *et al.*, 1947; Honkapohja, 1949; Kroese, 1949; Thompson *et al.*, 1954; Saphir and Ney, 1966; Stavem *et al.*, 1968).

Departments of Haematology and Rheumatology, Northwick Park Hospital and Clinical Research Centre, Harrow, Middlesex

J. M. ENGLAND, M.B., B.S., B.Sc., M.R.C. Research Fellow

D. S. SMITH, M.B., CH.B., M.R.C.P., Consultant Physician in Rheumatology and Rehabilitation

Case Report

A 56-year-old woman with a two-year history of rheumatoid arthritis gained little relief on salicylate therapy and was given a course of sodium aurothiomalate injections. The first injection of 5 mg was given without any side effects being noted (this was 34 days before Day 0 on the Chart), and further injections totalling 140 mg were given over the ensuing 28 days. During this period her platelet count remained at around 300,000/mm³, with a normal haemoglobin and white blood count. No eosinophilia was seen.

Seven days after her last injection of gold she presented with widespread purpura and a subconjunctival haemorrhage in the right eye. Pulse 76/min, B.P. 125/85 mm Hg. There was no splenomegaly or lymphadenopathy, and no skin lesions other than purpura were found. Some blood was present in the urine, but tests for occult blood in the stool were negative.

The haemoglobin was 12.9 g/100 ml, white blood count was 7,900/mm³ platelets were <5,000/mm³, three tests for L.E. cell reaction were negative, and prothrombin and kaolin-cephalin times, urea and electrolytes, liver function tests, and chest x-ray appearances were all normal. No platelet antibodies were detected when looking for platelet agglutinins or when testing for incomplete antibodies by the direct antiglobulin consumption test or the complement fixation test.

The thrombocytopenia was initially treated with infusions of either platelet concentrate (15 units/day) or platelet-rich plasma (10 units/day with frusemide) and no further injections of gold were given.

After seven days of platelet therapy there had been only a transient rise in the platelet count to 23,000/mm³ after the administration of platelet-rich plasma. At this stage the bone marrow showed many young megakaryocytes with little evidence of platelet budding. (The Chart shows the platelet count each morning. The platelet transfusions were given in the afternoon.) At other times the platelet

count was $<5,000/\text{mm}^3$ and the patient had further episodes of purpura. It was therefore decided to start her on steroids.

Prednisolone was given at a dose of 30 mg/day and the platelet transfusions were continued. Once again there was only a transient rise in the platelet count after the infusion of platelet-rich plasma. Indeed after eight days of prednisolone therapy the platelet count

Clinical course during treatment in a case of gold-induced thrombocytopenia.

was still $<5,000/\text{mm}^3$. At this point the patient had been thrombocytopenic for 15 days and there was no evidence of spontaneous recovery and no response to the course of steroids. The serum gold at this time, measured by atomic absorption spectrophotometry, was $74 \mu\text{g}/100 \text{ ml}$.

Treatment was then begun with intramuscular injections of 100 mg of dimercaprol (British anti-lewisite) twice daily for 15 days. There was a prompt and well-sustained rise in the platelet count which reached over $100,000/\text{mm}^3$ after four days of therapy. The platelet transfusions were therefore discontinued. Over the next few days there was a transient decline before the platelets rose to her previous levels of around $240,000/\text{mm}^3$.

After six days of dimercaprol therapy the serum gold had fallen to $19 \mu\text{g}/100 \text{ ml}$, and at the end of the course the serum gold was below the limit of reliable estimation ($10 \mu\text{g}/100 \text{ ml}$). Gold was detected in the urine only on the day after starting dimercaprol therapy, the excretion being $0.4 \text{ mg}/24 \text{ hr}$. The patient suffered no side effects from dimercaprol.

Finally, the prednisolone was reduced to 7.5 mg daily, and she was maintained on this therapy for her rheumatoid arthritis. The platelet count remained well maintained at around $300,000/\text{mm}^3$.

Comment

Severe thrombocytopenia has been seen many times during gold therapy but there is no general agreement about the aetiology, prevention, or treatment of the condition.

It has been suggested that the thrombocytopenia may be an immunological phenomenon. However, neither we nor Stavem *et al.* (1968) were able to show platelet antibodies by a wide variety of techniques. Indeed the rapid recovery seen after this patient was given dimercaprol might suggest that gold was exerting a direct toxic effect in her case. The bone marrow showing normal numbers of non-budding megakaryocytes is compatible with either immune or toxic thrombocytopenia.

The prevention of gold-induced thrombocytopenia appears to be difficult. Regular blood counts may be of help but the present patient showed no "warning" eosinophilia and had a normal blood count one week before the onset of purpura. It has been suggested that monitoring serum gold levels may reduce the incidence of side effects. This patient, however, had a low serum gold ($74 \mu\text{g}/100 \text{ ml}$), and it is well recognized that thrombocytopenia can occur after small doses of gold have been given (Saphir and Ney, 1966).

The treatment of gold-induced thrombocytopenia varies widely. In this case there was no evidence of spontaneous recovery after 15 days of thrombocytopenia and no response to an eight-day course of steroids. Dramatic recovery took place only after dimercaprol had been given. The rise in platelet count occurred at a time when the serum gold was being excreted in the urine, and these findings emphasize the specific role of dimercaprol in treatment. The dosage was based on the suggestion of Saphir and Ney (1966) that the total dose of dimercaprol (in grammes) should exceed by a factor of 30 the total dose of gold (in grammes of elemental gold).

We wish to thank Dr. I. Chanarin for his help and encouragement. Miss C. E. F. Tidmarsh helped with the preparation of the platelet transfusion. J.M.E. is supported by a Medical Research Council Fellowship. Agglutination and complement fixation tests were carried out by Dr. K.L.G. Goldsmith and Miss D.M. Brazier, and Mr. R. G. S. Johns performed the spectrophotometry.

References

- Honkapohja, H. (1949). *Annales Medicinae Internae Fenniae*, 38, 33.
- Kroese, W. F. S. (1949). *Nederlandsch Tijdschrift voor Geneeskunde*, 93, 1967.
- Lockie, L. M., Norcross, B. M., and George, C. W. (1947). *Journal of the American Medical Association*, 133, 754.
- Meyler, L., and Herxheimer, A. (1968). *Side Effects of Drugs*, vol. 6, Amsterdam, Excerpta Medica.
- Saphir, J. R., and Ney, R. G. (1966). *Journal of the American Medical Association*, 195, 782.
- Stavem, P., Strømme, J., and Bull, O. (1968). *Scandinavian Journal of Haematology*, 5, 271.
- Thompson, M., Sinclair, R. J. G., and Duthie, J. J. R. (1954). *British Medical Journal*, 1, 899.