by grants from the Norwegian Council for Cardiovascular Disease and Merck, Sharp, and Dohme Ltd.

References

- Aitchison, J. *et al.* (1971). *American Heart Journal*, 82, 660.
 Braren, C. H., Campbell, R. G., Hashim, S. A., and Van Itallie, T. B. (1968). *American Journal of Medicine*, 45, 480.
 Brown, J. J., *et al.* (1969). *Proceedings of the Royal Society of Medicine*, 62, 1258.
 Brown, J. J. *et al.* (1971a). *British Medical Bulletin*, 27, 128.
 Brown, J. J. *et al.* (1972). *British Medical Journal*, 2, 729.
 Brown, J. J., Davies, D. L., Lever, A. F., Peart, W. S., and Robertson, J. I. S. (1964). *British Medical Journal*, 2, 1636.
 Brown, J. J., Davies, D. L., Lever, A. F., Peart, W. S., and Robertson, J. I. S. (1965). *Journal of Endocrinology*, 33, 279.
 Brown, J. J., Ferriss, J. B., Fraser, R., Lever, A. F., and Robertson, J. I. S. (1971b). In *Medical Uses of Spironolactone*, ed. G. M. Wilson, p. 27. Amsterdam, Excerpta Medica.
 Brown, J. J., Fraser, R., Lever, A. F., and Robertson, J. I. S. (1971c). *Abstracts of World Medicine*, 45, 549, 633.
 Bull, M. B., and Laragh, J. H. (1968). *Circulation*, 37, 45.
 Crane, M. G., and Harris, J. J. (1970). *American Journal of Medical Sciences*, 260, 311.
 Davies, D. L., and Robertson, J. W. K. (1973). *Metabolism*. In press.
 Ferriss, J. B. *et al.* (1970). *Lancet*, 2, 995.
 Fraser, R., Thomson, S., and Young, J. (1973). In press.
 Lant, A. F., Smith, A. J., and Wilson, G. M. (1969). *Clinical Pharmacology and Therapeutics*, 10, 50.
 Rose, G. A., Holland, W. W., and Crowley, E. A. (1964). *Lancet*, 1, 296.
 Spark, R. F., and Melby, J. C. (1968). *Annals of Internal Medicine*, 69, 685.

Unilateral Renal Hypoplasia with Hypertension (Ask-Upmark Kidney)

J. B. ROSENFELD, L. COHEN, I. GARTY, M. BEN-BASSAT

British Medical Journal, 1973, 2, 217-218

Congenital unilateral renal hypoplasia with hypertension was first reported by Ask-Upmark (1929). Habib *et al.* (1965) reported nine more cases of this rare condition, which they called "segmental hypoplasia of the kidney." It has a typical histological picture, affects primarily children and young adults, and is sufficiently characteristic for it to be diagnosed and treated accordingly. A case is reported here.

Case Report

The patient, a 17-year-old girl, was admitted to hospital in 1966 because of severe headaches, vertigo, and general fatigue of two months' duration. Her blood pressure was 220/115 mm Hg supine and 220/150 mm Hg erect. The appearance of the retina was normal. A chest radiograph showed moderate enlargement of the left ventricle, and the electrocardiogram showed left ventricular hypertrophy. The blood urea was 45 mg/100 ml, creatinine clearance 65 ml/min, and the urinary sediment contained 3-4 leucocytes. There was no proteinuria and urinary cultures were sterile. The haemoglobin was 14 g/100 and the white blood cells 8,000 with a normal differential count. Liver function tests were normal. An intravenous pyelogram showed that the right kidney was 3.5 cm smaller than the left, the upper right calyx showed signs of clubbing, and the upper and lower poles showed lobulation. The dye concentration appeared equal on both sides. A renogram after intravenous injection of 15 μ Ci 131 I-labelled orthoiodohippurate showed diminished concentration and prolonged transit time in the right kidney. The angiotensin infusion test was negative and the Stamey test gave equivocal results. Renoangiography showed normal renal arteries on both sides but a paucity of vessels in the upper pole of the right kidney with lack of arborization and a tapering of the arterioles (fig. 1). In the nephrographic phase the effect was stronger on the right side, suggesting a paradoxical nephrographic effect (fig. 2).

A tentative diagnosis of contracted right kidney after pyelonephritis was unsatisfactory in view of the peculiar shape of the small kidney, the absence of a history of recurrent urinary tract infections, and the absence of anomaly of the urinary tract. Bilateral renal biopsy specimens showed some sclerosis of the arteries and

arterioles, more prominent on the left side. In line with current opinion in 1966 it was decided that the patient was unlikely to benefit from right nephrectomy, and she was treated conservatively.

During the next four years the patient had recurrent urinary tract infections. Despite control of her hypertension with methyl-dopa and a low salt diet grade II hypertensive changes developed in the retina, the blood urea rose progressively to 70 mg/100 ml, and her creatine clearance decreased to 40 ml/min. Traces of protein appeared in the urine. Selective renin studies in 1970

FIG. 1—Nephroangiography shows paucity of vessels to upper pole of right kidney, with lack of arborization and tapering of the arterioles.

FIG. 2—Nephrographic phase: stronger dye on right side.

Tel-Aviv University Medical School, Beilinson Medical Centre, Petah Tikvah, Israel

J. B. ROSENFELD, M.D., Head, Department of Medicine C
 L. COHEN, M.D., M.Sc., Senior Registrar, Department of Medicine C
 I. GARTY, M.D., Head, Division of Vascular Radiology
 M. BEN-BASSAT, M.D., Senior Physician, Department of Pathology

showed renin concentration from the right renal vein to be 2,715 ng/100 ml and from the left 345 ng/100 ml. In view of these findings a right nephrectomy was performed.

During the first postoperative year the patient's blood pressure dropped to normal and the hypertensive retinal changes regressed. The creatinine clearance stabilized at 55 ml/min, the blood urea was 60 mg/100 ml, and the serum electrolytes were normal. After two years the blood pressure again rose to 180/110 mm Hg. She responded well to treatment with propranolol.

PATHOLOGY

Renal Biopsy.—The arteriole walls were thickened by deposition of homogenous, acidophil, hyaline substance. The arterial arteriosclerotic changes consisted mainly of fibrous thickening of the intima, causing narrowing of the lumen. There was fraying and splitting of the internal elastic lamina, mainly in the arteries of the "healthy" kidney. Slight glomerular ischaemic changes and tubular atrophy were noted. No histological findings compatible with pyelonephritis were found.

Excised Kidney.—The excised right kidney measured 6.0 by 5.0 by 3.0 cm and weighed 80 g. Two deep surface grooves caused a coarse lobulation typical of renal segmental hypoplasia (Ask-Upmark kidney) (fig. 3). Frontal section through the kidney showed that the grooves appearing on the capsular surface marked the sites of hypoplastic segments, with underlying, calyx-like recesses of the renal pelvis. The number of the renal pyramids was reduced to four. The renal artery was thin, with a narrow lumen.

FIG. 3—Typical gross features of Ask-Upmark kidney.

Microscopically (see fig. 4) the hypoplastic segments consisted of cystically dilated tubules lined with a single layer of flat epithelium and containing a colloid-like substance. Between these thyroid-like structures a moderate amount of connective tissue and thick-walled arterioles and small arteries were seen. The thyroid-like tubules were devoid of glomeruli. There was no inflammatory infiltration.

Comment

In 1,200 necropsies and 100 nephrectomies Ljungqvist and Lagergren (1962) found only three cases of Ask-Upmark kidney. Its typical gross features are hypoplasia, an abnormally small number of pyramids, and a transverse surface groove. The syndrome occurs more commonly in girls and presents with hypertension before the age of 12. Headache is a leading symptom either in isolation or in association with hypertensive encephalopathy. The urine volume in these cases is usually normal, with a normal concentration, and seldom exceeds 2,000 ml/24 hr. Proteinuria, when present, is never more than 1 gm/24 hr. The case described here has these characteristics, except that the patient was aged 17.

FIG. 4—Hypoplastic segment consisting of atrophic, thyroid-like tubules localized between two zones of normal renal tissue. A small nest of normal renal tissue is seen on the top of groove. (H.E. $\times 30$.)

The following facts support the view that this unilateral segmental renal hypoplasia is congenital and also differentiate the condition from chronic pyelonephritis or atrophic segmentation of vascular origin. (1) The reduction in the size and weight of the affected kidney are rarely associated with inflammation or vascular changes. (2) The renal artery is small and this is not secondary to atherosclerotic changes, since the walls of the renal artery in this nephropathy are always normal. (3) The segments affected may be in the cortex and medulla of one or several renal lobes, but they do not have the fan-like distribution characteristic of chronic pyelonephritis and renal infarctions.

The thyroid-like dilatation of the tubules considered as pathognomonic for chronic pyelonephritis (Weiss and Parker, 1939) has also been observed in renal ischaemias and hypoplasias. In addition Fahr (1938) and Batzenschlager *et al.* (1962) regard the complete disappearance of the glomeruli in the cortex to be characteristic of segmental hypoplasia and not of chronic pyelonephritis. The presence of both arteriolar and glomerular sclerosis secondary to hypertension in the non-hypoplastic part of the kidney explains why nephrectomy does not usually abolish the hypertension (Bernstein, 1968). It may also explain the mild functional impairment in the healthy kidney.

The cause of unilateral small kidney with hypertension has almost always been attributed in English medical literature to pyelonephritis or primary hypoplasia, and the early descriptions of Ask-Upmark and Habib *et al.* have been consistently overlooked.

References

- Ask-Upmark, E. (1929). *Acta Pathologica et Microbiologica Scandinavica*, 7, 383.
- Batzenschlager, A., Blum, E., and Weill Bousson, M. (1962). *Annales de Anatomie et Pathologie*, 7, 427.
- Bernstein, J. (1968). *Pathology Annual*, 217.
- Fahr, T. (1938). *Virchows Archiv für pathologische Anatomie und Physiologie und für Klinische Medizin*, 301, 140.
- Habib, R., Courtecuisse, V., Ehrensperger, J., and Roger, P. (1965). *Semaine des Hôpitaux (Annales de Pédiatrie)*, 41, 954.
- Ljungqvist, A., and Lagergren, C. (1962). *Acta Pathologica et Microbiologica Scandinavica*, 56, 277.
- Weiss, S., and Parker, F. (1939). *Medicine*, 18, 221.