particularly those with HLA 8 antigen born in coeliac families, not be fed wheat cereals in their mixed feeds till, say, 6 months of age?—I am, etc.,

P. ASQUITH

Department of Experimental Pathology,
University of Birmingham

W. T. COOKE

General Hospital, Birmingham

- 1 Thompson M. W., *American Journal of Human Genetics* 1951, 3, 159.
- 2 Stokes, P. L., Asquith P., Holmes, G. K. T., Mackintosh, P., and Cooke, W. T., *Lancet*, 1972, 2, 162.
- 3 Falchuk, Z. M., Rogentine, G. N., and Strober, W., *Journal Clinical Investigation*, 1972, 51, 1602.
- 4 McNeish, A. S., Nelson, R., and Mackintosh, P., *Lancet*, 1973, 1, 668.
- 5 Shukla, A. Forsyth, H. A., Anderson, C. M., and Marwah, S. M., *British Medical Journal* 1972, 4, 507.
- 6 Visakorpi, J. K., and Immonen P., *Acta Paediatrica Scandinavica*, 1967, 56, 49.

Medical Staff Dining-rooms

SIR,—The publication of Dr. M. W. Rout's letter (28 April, p. 244) was very timely, for just recently our hospital management committee has decided against a separate medical staff dining-room for lunches (following a Ministry directive) in spite of strong opposition from the medical staff here. We have yet to receive a satisfactory reply to what we consider the most important argument in favour of such a provision—the need for a forum isolated from prying ears, for the informal discussion of medical problems across the specialties. This eliminates the need for impersonal, cumbersome, and inefficient referral forms or letters on often trivial problems, with a consequent saving of time for inpatients, more rapid institution of their treatment, and improved inter-departmental relations and co-operation—something that cannot be valued in strictly economic terms.—I am, etc.,

R. C. REDMAN

Battle Hospital,
Reading

Actinomycin D for Wilms's Tumour

SIR,—The data given in Dr. H. W. C. Ward's letter (17 February, p. 420) can be used to question the value of actinomycin D in the management of Wilms's tumour only in a negative sense. Preference must be given to positive data.

A good example is the cure rate for pulmonary metastases in Wilms's tumour. At this hospital 14 out of 21 children have survived longer than three years after the treatment of pulmonary metastases by total lung irradiation (270 kV, 2,000 rads in 28 days) combined with a single course of actinomycin D (15 µg/kg body weight daily during the first five days of irradiation). Using an exactly similar method of irradiation only one out of eight children survived without the use of actinomycin D. Since the latter experience was the rule before the introduction of actinomycin D our more recent results have to be attributed to the increased effect of the combination of these agents. The comparison of sequential retrospective data admittedly is of limited value, but in this instance the magnitude of the difference must surely dictate that the effect is real, and a random trial with and without actinomycin D cannot be justified.

If the value of actinomycin D in the

management of pulmonary metastases is accepted, its use in a single course during postoperative irradiation seems logical. The value of the combination in this situation is not so easily demonstrable, for abdominal recurrence after surgery and irradiation alone is relatively uncommon. However, our improved survival results at three years can be attributed only in part to the salvage of children with pulmonary metastatic disease. It seems reasonable to infer that actinomycin D given with postoperative irradiation is relevant to this improvement. The three-year survival results from the Toronto region, as reflected by the registry at this hospital for the years 1964-8 inclusive, are compared below with the results from the Birmingham region for 1965-9 as given by Dr. Ward.

	Toronto	Birmingham
All Stages	49/66* (74%)	13/46 (28%)
Stage I + II	25/30 (83%)	11/18 (61%)
Stage III	10/15 (67%)	2/16 (12%)
Stage IV	3/4	0/7
Stage V	3/3	0/5
Stage uncertain ..	8/12	

* Two lost to follow-up, counted as dead.

It is not possible to attribute as large a difference in overall survival as 74% versus 28% to different policies in relation to actinomycin D alone. It is quite clear that in Wilms's tumour good results are obtained only when the surgeon, radiotherapist, and paediatrician work with a common policy in a unit seeing a reasonable number of these children each year. It may be that these factors are more relevant to the results from Birmingham than the simple question of the value of actinomycin D.—I am, etc.,

R. D. T. JENKIN

Princess Margaret Hospital,
Toronto, Canada

SIR,—Dr. H. W. C. Ward's letter on this subject (17 February, p. 420) leads me to recount the experience in Melbourne which was briefly reviewed in 1970.¹

Sixty-three cases were seen in the period 1930-65 at the Peter MacCallum Clinic and the Royal Children's Hospital, Melbourne. Seven were excluded from the radically treated series because of the inadequate treatment or metastatic disease. None survived and none had actinomycin D. The remaining 56 were treated in two non-contemporaneous groups in a very similar fashion with the exception that the later cases had actinomycin D as soon as the diagnosis was suspected. Survival at three years was as follows.

	No Actinomycin	Actinomycin D
Clean nephrectomy	6/15 (40%)	10/14 (71%)
Tumour spill or definite residual disease .. .	2/10 (20%)	9/16 (56%)
Preoperative x-ray therapy .. .	0/1	—
Total	8/26 (31%)	19/30 (63%)

These results are significantly different at the 5% level using the corrected χ^2 test and it would therefore be difficult to convince any paediatric surgeon or radiotherapist in Melbourne that a controlled trial was ethically defensible.

There is one obvious difference between Birmingham and Melbourne in the method of dealing with such cases. While in Birmingham actinomycin D is given only when the diagnosis has been confirmed at operation, in Melbourne the drug is started as soon as there is a high index of suspicion—usually on the evidence of an aortogram or inferior vena cavagram—with operation following on the third day. All abdominal masses in children are treated as emergencies, and handling of the child is kept to a minimum. This policy is rendered easier by the centralization of paediatric and radiotherapy services.—I am, etc.,

T. F. SANDEMAN

Melbourne, Australia

1 Sandeman, T. F., *Australasian Radiology*, 1970, 14, 285.

Efficacy of Whooping-cough Vaccines

SIR,—A recent report of the Public Health Laboratory Service (3 February, p. 259) showed that the pertussis vaccines produced by two manufacturers, whose names were published, gave significantly different levels of protection in children. I feel, therefore, that I can now disclose the identities of some vaccines which we have tested here in Manchester over the past 10 years but which were given only code letters in the papers previously submitted to you for publication.

The P.H.L.S. report showed that vaccines produced by Wellcome Laboratories were more effective than those of Glaxo Laboratories but that the vaccines concerned were made more than—and, for some children in the survey, much more than—six years ago. The findings justify my own misgivings at that time¹⁻³ about vaccines made before 1967. Pertussis agglutinogens 1, 2, and 3 were all present in Wellcome vaccines (designated C)¹ but agglutininogen 3 was not detected in Glaxo vaccines (A and R).^{1,2} Indeed, in referring to the Glaxo product of that time I said³ that it was "unlikely that such vaccines would give satisfactory immunity against prevalent strains" of type 1, 3.

The situation since then is very different. Of vaccines made in 1967 the Glaxo product (vaccine X)⁴ gave an even better response to agglutininogen 3, in both children and rabbits, than did the Wellcome product (vaccine Y); and a similar excellent response has been found with vaccine made in 1970. It would therefore not be in accordance with the available evidence if those administering pertussis vaccine to children were to think that recent Glaxo vaccine was still inferior to Wellcome vaccine. Indeed, these results support our optimistic view of the future efficacy of British pertussis vaccines,⁵ which was based on the improved protection given to children by vaccines made during and after 1967.

Unfortunately for those who wish to ensure that suitable laboratory tests are used to monitor the protective potency of batches of pertussis vaccine this recent improvement in the efficacy of British vaccine has been accompanied by four changes that were all introduced at about the same time.⁶ In addition to the inclusion of detectable amounts of agglutininogen 3, mentioned above, the minimal acceptable mouse-protective potency of British vaccine has been raised from two to four international units per human

dose, adjuvant has been reintroduced, and the recommended schedule for the vaccination of children has been changed.

In Israel a similar improvement in the efficacy of pertussis vaccine has been noted,⁷ but with fewer accompanying changes. In that country the improvement dates from 1968, when the inclusion of agglutinin 3 in their vaccine was ensured. The earlier vaccine, which gave unsatisfactory protection, did, however, contain adjuvant and it also achieved a mouse-protective potency of four international units per human dose.⁸ Moreover, the vaccination schedule in Israel has not been altered.

The one feature common to the improvements in British and Israeli vaccines is that their manufacturers now make deliberate attempts to ensure that their products contain enough agglutinin 3. But ought our hopes for the wellbeing of vaccinated children to be based merely on the good-will of present well-informed staff, adequate though this proves to be at present? Or should we formulate requirements which would ensure that present knowledge is not forgotten?

To conclude: though there is evidence that for some years before 1967 the pertussis vaccine of Glaxo Laboratories was less effective than that of Wellcome Laboratories, there is ample reason to believe that this does not apply to vaccine made during the last six years. There is also further evidence that effective pertussis vaccines are those that contain sufficient agglutinin 3.—I am, etc.,

NOEL W. PRESTON

Department of Bacteriology and Virology,
University of Manchester

- ¹ Preston, N. W., *British Medical Journal*, 1963, 2, 724.
- ² Preston, N. W., *British Medical Journal*, 1965, 2, 11.
- ³ Preston, N. W., *British Medical Journal*, 1965, 2, 1001.
- ⁴ Abbott, J. D., Preston, N. W., and Mackay, R. I., *British Medical Journal*, 1971, 1, 86.
- ⁵ Preston, N. W., and Stanbridge, T. N., *British Medical Journal*, 1972, 3, 448.
- ⁶ Perkins, F. T., *British Medical Journal*, 1969, 4, 429.
- ⁷ Shmilovitz, M., Preston, N. W., Zaltser, H., and Cahana, A., *Israel Journal of Medical Sciences*, 1972, 8, 1936.
- ⁸ Singer, E. D., 1973. Personal communication.

Discrimination against Rhodesian Nurses

SIR,—In Personal View (28 April, p. 238) Professor Levy says that regulations governing the entry of Rhodesian students into the U.K. have recently come into effect and are manifestly discriminatory against the young Rhodesian girl who seeks to train as a nurse in the U.K. It would even appear that the discrimination is maximal against the less well-educated girl who wishes to train for the S.E.N. diploma rather than for the S.R.N. The raised educational standards demanded by the new regulations apply exclusively to applicants from Rhodesia.

Personally I have no sympathy with the present rebel Rhodesian regime and I would approve any whole-hearted action our Government might take against it. When, however, the action involves petty discrimination against trainee nurses, then surely it is time our nursing institutions and the B.M.A. united to protest that the professions of nursing and medicine have a long tradition of international co-operation and have in the past striven to rise above national disputes. Our British institutions

should protest against the new discriminatory measures.

World opinion is hardly likely to view our new regulations as evidence of our Government's firm determination to deal with the Rhodesian problem, but our colleagues abroad will note the profession's silent acquiescence as evidence that we care less about our international obligations than we did.—I am, etc.,

PETER DIGGORY

London W.1.

Resources and Needs in Africa

SIR,—In your leading article (5 May, p. 255) you commented with editorial sympathy on the first International Congress on Basic Medical Sciences in Africa, held at Lusaka in March.

Will you allow me, in the name of the trustees of the Commonwealth Foundation, to mention that the presence of many of the distinguished Commonwealth African delegates at this meeting was made possible by a Foundation grant of £4,250? This does not in the slightest detract from the imaginative and far-sighted planning of Professor Ezeilo and his colleagues, with whom the writer had the privilege of discussing plans for the conference as far back as November 1971. The main point is that you have commended the value of this meeting—a judgement which will certainly not be lost on our trustees.—I am, etc.,

JOHN CHADWICK

Director,
The Commonwealth Foundation

London S.W.1

Prescribing Mandrax

SIR,—I was interested to read the letters of Drs. G. C. Mathers and R. W. Green (28 April, p. 243) and of Dr. H. J. S. Matthew (12 May, p. 367) commenting on my letter published earlier in the *B.M.J.*, (7 April, p. 54).

I would like to stress that I did not write my letter to extol the virtues of Mandrax (methaqualone and diphenhydramine). My complaint was of a communication I had received from the senior medical officer. I can assure Dr. Mathers that I am aware of the addictive qualities of Mandrax, and if he will refer to my letter again he will note that only a little over 1% of my patients had been prescribed this drug, and all the recipients were of a mature age. I cannot ever remember prescribing the drug to a youngster—in fact I find it unusual for a teenager to come to me with a sleep problem, and when they do it is usually amitriptyline which they receive.

I completely reject the advice of Dr. Green that I should accept the S.M.O.'s communication "in the manner intended." I considered that the letter was unnecessary, and it is my interpretation, devoid of paranoia, that the "manner intended" could have had but one object, and that was to intimidate me, and I think that intimidation should be resisted. I should imagine that the S.M.O. has powers that he can exercise over erring doctors, and I am very pleased that he should have such power, but he should abandon his tactics of attempted intimidation.

I have before me the last published details of suicidal poisoning in England and Wales during the year 1971 which have been ex-

tracted by the General Register Office.¹ The list of drugs is a very formidable one and, as one would expect, deaths due to barbiturates outweigh those due to any other listed drug; contrary to Dr. Green's and Dr. Matthew's assertion there were six listed deaths attributable to Mogadon (nitrazepam).

Though as I stated earlier the intention of my letter was to draw attention to a letter I received from the S.M.O. and not to discuss the virtues or otherwise of drugs, I would like, if I may, to comment on Dr. Matthew's letter. Dr. Matthew suggests that a voluntary ban on Mandrax could help to prevent thefts from chemist's shops. Why did he not follow his suggestion to its logical conclusion and suggest a ban on barbiturates as well? It is not my experience to find that Mandrax is a drug that my patients become addicted to. There is not a demand for increasing dosage by those to whom the drug is prescribed. A half to one tablet at night are the requirements of my patients, and I could count on the fingers of one hand those who require two tablets at night. I am in no position to know the most popular drug of addiction apart from alcohol—and I'll wager that most of us have a tippie at this—but my guess would be that barbiturates would far outweigh all other prescribable drugs put together, and as to the cause of death by drug overdosage there were 286 deaths from Tuinal (quinalbarbitone and amylobarbitone) alone in 1971, and 43 from Mandrax.¹

That the S.M.O. should "politely draw" my attention to those of my patients to whom I prescribe Mandrax is what irritates me, besides the implied threat already mentioned. I regard it as an impertinence that a remote Ministry official should write and tell me what I'm prescribing for my patients. I do not prescribe sleeping tablets willy-nilly; it often happens that patients with a sleeping problem do not require hypnotics at all. I wish to remain master of my own house, and be free to practise as I see fit. If I err, then let officialdom act.

The day has drawn to a close and bedtime has arrived. Rather than risk taking half a Mandrax to soothe me off to sleep I'm going to have a nice stiff whisky instead. Good health and good night.—I am, etc.,

C. G. BROWN

Slough, Bucks

¹ General Register Office, *Pharmaceutical Journal*, 1973, 210, 77.

Behçet's Syndrome and Oral Fibrinolytic Therapy

SIR,—We were interested to read the paper of Drs. T. Chajek and M. Fainaru (31 March, p. 782) and, by reference to a similar case, would like to support their observation that impaired fibrinolysis is an important factor in the pathogenesis of Behçet's syndrome.

We have under our care a male patient now aged 32. For eight years he has had recurrent attacks of deep venous thrombosis and one episode of pulmonary embolism. During this time he has also suffered from recurrent mouth ulcers, scrotal ulcers, erythema-nodosum-like lesions, and superficial phlebitis. When referred to us he had lower inferior vena caval obstruction but there was no proteinuria, suggesting that the renal veins were not affected. For a few years he had been treated with intravenous heparin and oral anticoagulants. Though this had